

<sup>3</sup> Sobhraj Maternity Hospital, Gynaecology, Karachi, Pakistan

<sup>4</sup> South London & Maudsley NHS Foundation Trust, Psychiatry, London, United Kingdom

<sup>5</sup> Dow University of Health Sciences, Institute of Nursing, Karachi, Pakistan

<sup>6</sup> The University of Central Lancashire, Preston, United Kingdom

<sup>7</sup> Tech4Life Enterprises, Software, Karachi, Pakistan

<sup>8</sup> Queens University, Psychiatry, Kingston, Canada

\* Corresponding author.

**Introduction** Postnatal depression is known to cause disability and suffering in women and negative consequences both for their infants and their families, with huge costs globally. Several studies from low and middle income countries (LAMIC) have demonstrated that effectively delivered psychological interventions are cost effective for improving maternal and child health, but access to these interventions is limited in both the low and high income countries.

**Objective** The objective of the study is to develop and test a mobile phone-based intervention (TechMotherCare), which will include components of cognitive behavioural therapy (CBT) and child development related psychoeducation.

**Aim** The aim of the study is to examine the feasibility and acceptability of the TechMotherCare intervention.

**Methods** A total of 36 participants will be recruited from health centers in Karachi, Pakistan for this 2-arm randomized pilot study. The TechMotherCare App intervention will be based on principles of CBT and learning-through-play (LTP) a parenting intervention and will assess the real-time depressive symptoms of participants and respond, using intelligent real time therapy (iRTT) dependent on symptoms reported by participants.

**Results** Outcome assessments will be completed after 3 months (end of intervention). In-depth qualitative interviews will also be conducted with participants pre- and post-intervention. The trial is ongoing and we will present both the qualitative and quantitative results.

**Conclusions** The results of this pilot trial will inform the design of a larger randomised controlled trial using a mobile based technology platform to address the huge treatment gap in LAMICs.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2276>

## EV1292

### E-mental health: Updates on recent achievements and pitfalls

D. Hilty<sup>1,\*</sup>, A. Fiorillo (Direttore)<sup>2</sup>, K. Krysta<sup>3</sup>, M. Krausz<sup>4</sup>, D. Mucic<sup>5</sup>

<sup>1</sup> Aligned Telehealth, Telemedicine, Davis, USA

<sup>2</sup> Centro Collaboratore dell' Organizzazione Mondiale, Dipartimento di Psichiatria dell', Università SUN, Naples, Italy

<sup>3</sup> Medical University of Silesia, Department of Psychiatry and Psychotherapy, Sosnowiec, Poland

<sup>4</sup> School of Population and Public Health, Centre for Health Evaluation and Outcome Sciences CHÉOS, Vancouver, Canada

<sup>5</sup> The Little Prince Psychiatric Centre, Psychiatry, Copenhagen, Denmark

\* Corresponding author.

The patient-centered care features quality, affordable, and timely care in a variety of settings – technology is a key part of that – particularly among younger generations and child and adolescent patients. The consumer movement related to new technologies is nearly passing clinicians by, as new ways of communicating with others (text, e-mail, Twitter, Facebook) revolutionizes how we experience life and access healthcare. This paper explores a continuum with healthy, innovative behavior on one end (e.g., social media) and pathological Internet use on the other end – and the range of self-help and e-mental healthcare options being used. Specifically, it focuses on how social media adds to,

yet may complicate healthcare delivery, such that clinicians may need to adjust our approach to maintain therapeutic relationships, interpersonal/clinical boundaries, and privacy/confidentiality. We suggest planning ahead to discuss expectations about online communication between doctors and patients as part of the informed consent process, offer other do's and don't's for patients and clinicians, and review applicable guidelines. More research is needed on consumer and patient use of technology related to healthcare, as is an approach to basic and advanced measurement of outcomes.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2277>

## EV1293

### Telepsychiatry in the opinion of Polish patients and psychiatrists

M. Wojtuszek, J. Kachnic, J. Wutke, K. Krysta\*

Medical University of Silesia, Department of Psychiatry and Psychotherapy, Katowice, Poland

\* Corresponding author.

**Background** Telepsychiatry takes advantage of modern communication technologies and new forms of patient–doctor and doctor–patient contact.

**Objective** This research focuses on the knowledge of telepsychiatry and the attitude to it among patients and psychiatrists in Poland. **Subject and methods** An anonymous survey was conducted among 105 psychiatrists aged 26–74, including 74 women and 31 men and 102 patients aged 21–79, including 61 women and 41 men.

**Results** Research reveals that the majority of patients never met with the concept of “e-psychiatry” and do not know what it means. However, more than 50% of respondents answered positively to every question considering the utility of telepsychiatry. Furthermore according to 18%, it is possible to replace an eye-to-eye conversation by videoconferencing. Only 15% of doctors claim to have an extensive knowledge on telepsychiatry, and 10% do not know what it means. The vast majority of physicians perceive positive aspects of this method of medical care, but 63% would not want any general insertion of telepsychiatry. Doctors are apprehensive of losing personal data and medical confidentiality, and of the necessity of legal changes.

**Conclusions** The obtained results allow us to conclude that Polish patients and doctors, regardless of their knowledge, age, gender or disease, perceive advantages of telepsychiatry. In connection with this, implementing this method into the Polish medical market makes sense and is in accordance with both patients' and doctors' opinion. Based on our research, we confirmed that there is a necessity of wider popularisation of telepsychiatry among Polish therapists.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2278>

## EV1294

### MindfulSpot: A mindfulness mobile app for people dealing with infertility

B. Monteiro<sup>1,\*</sup>, A. Galhardo<sup>1,2</sup>, M. Cunha<sup>1,2</sup>, M. Couto<sup>1</sup>, F. Fonseca<sup>3</sup>, L. Carvalho<sup>4</sup>

<sup>1</sup> Instituto Superior Miguel Torga, Psychology, Coimbra, Portugal

<sup>2</sup> University of Coimbra, CINEICC, Faculty of Psychology and Educational Sciences, Coimbra, Portugal

<sup>3</sup> Instituto Superior Miguel Torga, Multimedia, Coimbra, Portugal

<sup>4</sup> Instituto Superior Miguel Torga, Technology and programming of information systems, Coimbra, Portugal

\* Corresponding author.