

**Results:** Patients were randomized to receive DVS 100-400mg (N=1048; 18-39 years [n=451]; 40-55 years [n=457]; >55 [n=140]) or placebo (N=718; 18-39 years [n=306]; 40-55 years [n=310]; >55 [n=102]). The final SDS, WHO-5, and VAS-PI overall pain change from baseline for the total population were significantly greater in the DVS vs the placebo group ( $P < 0.001$ ). Mean differences (adjusted ANCOVA) from placebo in women were: SDS: 18-39: -2.1, 40-55: -2.1, >55: -3.2; WHO-5: 18-39: +1.1, 40-55: +1.8, >55: +3.3; VAS-PI 18-39: -1.4, 40-55: -5.2, >55: -9.1. In men, results were consistent across age groups studied (SDS: 18-39: -2.9, 40-55: -3.4, >55: +2.6; WHO-5: 18-39: +1.5, 40-55: +2.4, >55: -0.3; VAS-PI 18-39: -3.0, 40-55: -7.7, >55: +2.0).

**Conclusions:** DVS improves functional outcomes and pain symptoms in depressed men and women across the age groups studied.

## P0242

Depression in coronary heart disease

A. Stetkiewicz<sup>1</sup>, A. Goch<sup>2</sup>, A. Borkowska<sup>1</sup>. <sup>1</sup>Department of Medical Psychology Medical University of Lodz, Lodz, Poland <sup>2</sup>Chair of Cardiology and Cardiosurgery Medical University of Lodz, Lodz, Poland

**Background and Aims:** Current studies on Coronary Heart Disease (CHD) show significant relationship with depression, which play important role in the course and clinical pictures of the illness. Depression is connected with worse course and more pessimistic prognosis in patients with CHD.

This study is expected to lead to better understanding the etiopathogenesis of CHD and its connections with depression and to create an efficient treatment programs in these disturbances.

**Methods:** Two methods estimating intensification of depressive symptoms were used in this study; Beck Depression Inventory (BDI) and Hamilton Depression Rating Scale (HDRS)

**Results:** To explore the connection between depression and CHD the investigation has been conducted in a group of 111 patients with different course of CHD. The age of the group was 41-65 years (mean = 55; SD = 5,6).

The results of the study indicated the sub clinical state of depression in the whole investigated group. The highest values of BDI and HDRS were noticed in women as well as of men and women with disability pension or in pensioners.

**Conclusions:** We conclude that women with CHD have higher level of depression and the active employment status has protective influence on the mood.

## P0243

Depression and the competence for selection, optimisation and compensation (SOC)

G. Stoppe. University Psychiatric Hospital, Basel, Switzerland

**Background:** The meta theory of selection, optimisation and compensation is a concept of successful coping and especially "successful" aging (1,2). Since depression is the most prevalent psychiatric disorder in old age, we wanted to know whether depression is associated with impairment of SOC-competence on a state or trait basis.

**Method:** In a pilot study n=50 inpatients took part after informed consent. They were between 21 and 73years old and suffered from a depressive syndrome (F32, F43.2, F33, F43.1, F31, F41.2). Patients with organic and/or psychotic underlying condition were excluded.

We applied a standardised scale, developed by Freund & Baltes (3). Together with standard instruments (MADRS, GDS) at the beginning and the end of inpatient treatment. For this study, data from a representative German study (ALLEE) served as control.

**Results:** As to be expected, SOC-competence was significantly lower compared to normal values in the depressed state. After remission however, there were no differences left. There were also no significant differences found when comparing a group of younger (<50y) and a group of older (>50y) patients.

**Conclusion:** We draw the preliminary conclusion that impairments in SOC competence seem to be no trait marker for depression at any age. Further study should test whether there are differences in various subtypes of depression (e.g. early versus late onset, double depression.....).

## P0244

Chaotic neural response reflecting stroop incongruent task is related to serum Cortisol levels in patients with unipolar depression

M. Susta<sup>1</sup>, P. Bob<sup>1</sup>, J. Pavlat<sup>1</sup>, T. Zima<sup>2</sup>, H. Benakova<sup>2</sup>, J. Raboch<sup>1</sup>. <sup>1</sup>Department of Psychiatry, First Faculty of Medicine, Charles University, Prague, Czech Republic <sup>2</sup>Department of Clinical Biochemistry and Laboratory Diagnostics, First Faculty of Medicine, Charles University, Prague, Czech Republic

**Background and Aims:** According to recent findings detecting a cognitive conflict is related to activation of anterior cingulate cortex (ACC) and central autonomic network. Several recent findings also suggest the hypothesis that the cognitive conflict is related to specific nonlinear chaotic changes of the neural signal. This conflict related activation elicits autonomic responses which can be assessed by psychophysiological measures such as heart rate variability calculated as beat to beat R-R intervals (RRI).

**Method:** The present study used Stroop word-colour test as an experimental approach to psychophysiological study of cognitive conflict in connection with RRI measurement, assessment of serum cortisol and calculation of largest Lyapunov exponents in nonlinear data analysis of RRI time series in 32 patients with unipolar depression.

**Result:** Significant correlation  $-0.43$  ( $p < 0.01$ ) between largest Lyapunov exponents during conflicting Stroop task and serum cortisol levels has been found.

**Conclusion:** The study indicates that a defect of neural inhibition during conflicting Stroop task is closely related to decreased serum cortisol levels which probably reflect defense psychological mechanisms.

## P0245

Depressive disorders and herpes simplex virus infection

G. Taldibaeva<sup>1</sup>, L. Bardenshteyn<sup>2</sup>, M. Gomberg<sup>1</sup>, I. Shcherbakova<sup>2</sup>. <sup>1</sup>Central Research Institute for Skin and Venereal Diseases, Moscow, Russia <sup>2</sup>Department of Psychiatry and Narcology, Moscow State University of Medicine and Dentistry, Moscow, Russia

**Introduction:** The impact of HSV infection on psychiatric status of patients correlates with the disease nature and duration. Recurrent genital herpes can have a major impact on sufferers' sex lives. HSV recurrences are often accompanied by anxiety and depression, up to suicidal attempts.

Purpose of this study is to investigate clinical typology of depressive disorders of HSV patients.