

LITHOBID®

(Lithium Carbonate, USP)
Slow-Release Tablets, 300 mg

Smooth, slow release of lithium carbonate for initial or maintenance treatment of mania associated with bipolar disorder

BRIEF SUMMARY:

The following is a brief summary only. Before prescribing, see complete prescribing information in LITHOBID® Slow-Release Tablets product labeling.

WARNING

Lithium toxicity is closely related to serum lithium levels, and can occur at doses close to therapeutic levels. Facilities for prompt and accurate serum lithium determinations should be available before initiating therapy (see DOSAGE AND ADMINISTRATION).

INDICATIONS:

Lithium is indicated in the treatment of manic episodes of manic-depressive illness. Maintenance therapy prevents or diminishes the intensity of subsequent episodes in those manic-depressive patients with a history of mania.

Typical symptoms: of mania include pressure of speech, motor hyperactivity, reduced need for sleep, flight of ideas, grandiosity, elation, poor judgment, aggressiveness, and possibly hostility. When given to a patient experiencing a manic episode, lithium may produce a normalization of symptomatology within 1 to 3 weeks.

WARNINGS:

Lithium should generally not be given to patients with significant renal or cardiovascular disease, severe debilitation, dehydration, sodium depletion, and to patients receiving diuretics, or angiotensin converting enzyme (ACE) inhibitors, since the risk of lithium toxicity is very high in such patients. If the psychiatric indication is life threatening, and if such a patient fails to respond to other measures, lithium treatment may be undertaken with extreme caution, including daily serum lithium determinations and adjustment to the usually low doses ordinarily tolerated by these individuals. In such instances, hospitalization is a necessity.

Chronic lithium therapy may be associated with diminution of renal concentrating ability, occasionally presenting as nephrogenic diabetes insipidus, with polyuria and polydipsia. Such patients should be carefully managed to avoid dehydration with resulting lithium retention and toxicity. This condition is usually reversible when lithium is discontinued. Morphologic changes with glomerular and interstitial fibrosis and nephron atrophy have been reported in patients on chronic lithium therapy. Morphologic changes have also been seen in manic-depressive patients never exposed to lithium. The relationship between renal function and morphologic changes and their association with lithium therapy have not been established.

Kidney function should be assessed prior to and during lithium therapy. Routine urinalysis and other tests may be used to evaluate tubular function (e.g., urine specific gravity or osmolality following a period of water deprivation, or 24-hour urine volume) and glomerular function (e.g., serum creatinine or creatinine clearance). During lithium therapy, progressive or sudden changes in renal function, even within the normal range, indicate the need for reevaluation of treatment.

An encephalopathic syndrome (characterized by weakness, lethargy, fever, tremulousness and confusion, extrapyramidal symptoms, leukocytosis, elevated serum enzymes, BUN and FBS) has occurred in a few patients treated with lithium plus a neuroleptic, most notably haloperidol. In some instances, the syndrome was followed by irreversible brain damage. Because of possible causal relationship between these events and the concomitant administration of lithium and neuroleptic drugs, patients receiving such combined therapy or patients with organic brain syndrome or other CNS impairment should be monitored closely for early evidence of neurologic toxicity and treatment discontinued promptly if such signs appear. This encephalopathic syndrome may be similar to or the same as Neuroleptic Malignant Syndrome (NMS).

Lithium toxicity is closely related to serum lithium concentrations and can occur at doses close to the therapeutic concentrations (see DOSAGE AND ADMINISTRATION).

Outpatients and their families should be warned that the patient must discontinue lithium therapy and contact his physician if such clinical signs of lithium toxicity as diarrhea, vomiting, tremor, mild ataxia, drowsiness, or muscular weakness occur.

Lithium may prolong the effects of neuromuscular blocking agents. Therefore, neuromuscular blocking agents should be given with caution to patients receiving lithium.

Usage in Pregnancy: Adverse effects on nidation in rats, embryo viability in mice, and metabolism in vitro of rat testis and human spermatozoa have been attributed to lithium, as have teratogenicity in submammalian species and cleft palate in mice.

In humans, lithium may cause fetal harm when administered to a pregnant woman. Data from lithium birth registries suggest an increase in cardiac and other anomalies, especially Ebstein's anomaly. If this drug is used in women of childbearing potential, or during pregnancy, or if a patient becomes pregnant while taking this drug, the patient should be apprised by their physician of the potential hazard to the fetus.

Usage in Nursing Mothers: Lithium is excreted in human milk. Nursing should not be undertaken during lithium therapy except in rare and unusual circumstances where, in the view of the physician, the potential benefits to the mother outweigh possible hazard to the child. Signs and symptoms of lithium toxicity such as hypertonia, hypothermia, cyanosis and ECG changes have been reported in some infants.

Usage in Children: Since the safety and effectiveness of lithium in children under 12 years of age has not been established, its use in such patients is not recommended at this time.

There has been a report of transient syndrome of acute dystonia and hyperreflexia occurring in a 15 kg child who ingested 300 mg of lithium carbonate.

PRECAUTIONS:

The ability to tolerate lithium is greater during the acute manic phase and decreases when manic symptoms subside (see DOSAGE AND ADMINISTRATION).

The distribution space of lithium approximates that of total body water. Lithium is primarily excreted in urine with insignificant excretion in feces. Renal excretion of lithium is proportional to its plasma concentration. The elimination half-life of lithium is approximately 24 hours. Lithium decreases sodium reabsorption by the renal tubules which could lead to sodium depletion. Therefore, it is essential for the patient to maintain a normal diet, including salt, and an adequate fluid intake (2500-3500 mL) at least during the initial stabilization period. Decreased tolerance to lithium has been reported to ensue from protracted sweating or diarrhea and, if such occur, supplemental fluid and salt should be administered under careful medical supervision and lithium intake reduced or suspended until the condition is resolved.

In addition to sweating and diarrhea, concomitant infection with elevated temperatures may also necessitate a temporary reduction or cessation of medication.

Previously existing thyroid disorders do not necessarily constitute a contraindication to lithium treatment. Where hypothyroidism preexists, careful monitoring of thyroid function during lithium stabilization and maintenance allows for correction of changing thyroid parameters and/or adjustment of lithium doses, if any. If hypothyroidism occurs during lithium stabilization and maintenance, supplemental thyroid treatment may be used.

In general, the concomitant use of diuretics or angiotensin converting enzyme (ACE) inhibitors with lithium carbonate should be avoided. In those cases where concomitant use is necessary extreme caution is advised since sodium loss from these drugs may reduce the renal clearance of lithium resulting in increased serum lithium concentrations with the risk of lithium toxicity. When such combinations are used, the lithium dosage may need to be decreased, and more frequent monitoring of lithium serum concentrations is recommended. See WARNINGS for additional caution information.

Concomitant administration of carbamazepine and lithium may increase the risk of neurotoxic side effects.

The following drugs can lower serum lithium concentrations by increasing urinary lithium excretion: acetazolamide, urea, xanthine preparations and alkalinizing agents such as sodium bicarbonate.

Concomitant extended use of iodide preparations, especially potassium iodide, with lithium may produce hypothyroidism. Indomethacin and piroxicam have been reported to significantly increase steady state serum lithium concentrations. In some cases lithium toxicity has resulted from such interactions. There is also some evidence that other nonsteroidal, anti-inflammatory agents may have a similar effect. When such combinations are used, increased serum lithium concentrations monitoring is recommended.

LITHOBID® (Lithium Carbonate, USP) Slow-Release Tablets, 300 mg

Concurrent use of calcium channel blocking agents with lithium may increase the risk of neurotoxicity in the form of ataxia, tremors, nausea, vomiting, diarrhea and/or tinnitus. Concurrent use of metronidazole with lithium may provoke lithium toxicity due to reduced renal clearance. Patients receiving such combined therapy should be monitored closely.

Concurrent use of fluoxetine with lithium has resulted in both increased and decreased serum lithium concentrations. Patients receiving such combined therapy should be monitored closely.

Lithium may impair mental and/or physical abilities. Patients should be cautioned about activities requiring alertness (e.g., operating vehicles or machinery).

Usage in Pregnancy: Pregnancy Category D (see WARNINGS).

Usage in Nursing Mothers: Because of the potential for serious adverse reactions in nursing infants from lithium, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother (see WARNINGS).

Usage in Children: Safety and effectiveness in children below the age of 12 have not been established (see WARNINGS).

Usage in the Elderly: Elderly patients often require lower lithium dosages to achieve therapeutic serum concentrations. They may also exhibit adverse reactions at serum concentrations ordinarily tolerated by younger patients. Additionally, patients with renal impairment may also require lower lithium dosages (see WARNINGS).

ADVERSE REACTIONS:

The occurrence and severity of adverse reactions are generally directly related to serum lithium concentrations and to individual patient sensitivity to lithium. They generally occur more frequently and with greater severity at higher concentrations.

Adverse reactions may be encountered at serum lithium concentrations below 1.5 mEq/L. Mild to moderate adverse reactions may occur at concentrations from 1.5-2.5 mEq/L, and moderate to severe reactions may be seen at concentrations from 2.0 mEq/L and above.

Fine hand tremor, polyuria and mild thirst may occur during initial therapy for the acute manic phase, and may persist throughout treatment. Transient and mild nausea and general discomfort may also appear during the first few days of lithium administration.

These side effects usually subside with continued treatment or with a temporary reduction or cessation of dosage. If persistent, a cessation of lithium therapy may be required. Diarrhea, vomiting, drowsiness, muscular weakness and lack of coordination may be early signs of lithium intoxication, and can occur at lithium concentrations below 2.0 mEq/L. At higher concentrations giddiness, ataxia, blurred vision, tinnitus and a large output of dilute urine may be seen. Serum lithium concentrations above 3.0 mEq/L may produce a complex clinical picture involving multiple organs and organ systems. Serum lithium concentrations should not be permitted to exceed 2.0 mEq/L during the acute treatment phase.

The following reactions have been reported and appear to be related to serum lithium concentrations, including concentrations within the therapeutic range:

Central Nervous System: tremor, muscle hyperirritability (fasciculations, twitching, clonic movements of whole limbs), hypertonicity, ataxia, choreoathetotic movements, hyperactive deep tendon reflex, extrapyramidal symptoms including acute dystonia, cogwheel rigidity, blackout spells, epileptiform seizures, slurred speech, dizziness, vertigo, downbeat nystagmus, incontinence of urine or feces, somnolence, psychomotor retardation, restlessness, confusion, stupor, coma, tongue movements, tics, tinnitus, hallucinations, poor memory, slowed intellectual functioning, startled response, worsening of organic brain syndromes. Cases of Pseudotumor Cerebri (increased intracranial pressure and papilledema) have been reported with lithium use. If undetected, this condition may result in enlargement of the blind spot, constriction of visual fields and eventual blindness due to optic atrophy. Lithium should be discontinued, if clinically possible, if this syndrome occurs. **Cardiovascular:** cardiac arrhythmia, hypotension, peripheral circulatory collapse, bradycardia, sinus node dysfunction with severe bradycardia (which may result in syncope); **Gastrointestinal:** anorexia, nausea, vomiting, diarrhea, gastritis, salivary gland swelling, abdominal pain, excessive salivation, flatulence, indigestion; **Genitourinary:** glycosuria, decreased creatinine clearance, albuminuria, oliguria, and symptoms of nephrogenic diabetes insipidus including polyuria, thirst and polydipsia; **Dermatologic:** drying and thinning of hair, alopecia, anesthesia of skin, acne, chronic folliculitis, xerosis cutis, psoriasis or its exacerbation, generalized pruritus with or without rash, cutaneous ulcers, angioedema; **Autonomic Nervous System:** blurred vision, dry mouth, impotence/sexual dysfunction; **Thyroid Abnormalities:** euthyroid goiter and/or hypothyroidism (including myxedema) accompanied by lower T₃ and T₄. ¹³¹Iodine uptake may be elevated (see PRECAUTIONS). Paradoxically, rare cases of hyperthyroidism have been reported. **EEG Changes:** diffuse slowing, widening of frequency spectrum, potentiation and disorganization of background rhythm. **EKG Changes:** reversible flattening, isoelectricity or inversion of T-waves. **Miscellaneous:** Fatigue, lethargy, transient scotomata, exophthalmos, dehydration, weight loss, leukocytosis, headache, transient hyperglycemia, hypercalcemia, hyperparathyroidism, albuminuria, excessive weight gain, edematous swelling of ankles or wrists, metallic taste, dysgeusia/taste distortion, salty taste, thirst, swollen lips, tightness in chest, swollen and/or painful joints, fever, polyarthralgia, and dental caries. Some reports of nephrogenic diabetes insipidus, hyperparathyroidism and hypothyroidism which persist after lithium discontinuation have been received.

A few reports have been received of the development of painful discoloration of fingers and toes and coldness of the extremities within one day of starting lithium treatment. The mechanism through which these symptoms (resembling Raynaud's Syndrome) developed is not known. Recovery followed discontinuance.

OVERDOSAGE:

The toxic concentrations for lithium (≥ 1.5 mEq/L) are close to the therapeutic concentrations (0.6-1.2 mEq/L). It is therefore important that patients and their families be cautioned to watch for early toxic symptoms and to discontinue the drug and inform the physician should they occur. (Toxic symptoms are listed in detail under ADVERSE REACTIONS).

Treatment: No specific antidote for lithium poisoning is known. Treatment is supportive. Early symptoms of lithium toxicity can usually be treated by reduction or cessation of dosage of the drug and resumption of the treatment at a lower dose after 24 to 48 hours. In severe cases of lithium poisoning, the first and foremost goal of treatment consists of elimination of this ion from the patient.

Treatment is essentially the same as that used in barbiturate poisoning: 1) gastric lavage, 2) correction of fluid and electrolyte imbalance and 3) regulation of kidney functioning. Urea, mannitol, and aminophylline all produce significant increases in lithium excretion. Hemodialysis is an effective and rapid means of removing the ion from the severely toxic patient. However, patient recovery may be slow.

Infection prophylaxis, regular chest X-rays, and preservation of adequate respiration are essential.

SOLVAY PHARMACEUTICALS

Marionetta, GA 30062

22E0803 Rev 2/95

© 1995 SOLVAY PHARMACEUTICALS, INC.

References: 1. Grof P, MacCrimmon D, Saxena B, et al. Bioavailability and side effects of different lithium carbonate products. *Neuropsychobiology*. 1976;2:313-323. 2. Shaw DM, Hewland R, Johnson AL, et al. Comparison of serum levels of two sustained-release preparations of lithium carbonate. *Curr Med Res Opin*. 1974;2:90-94. 3. Kirkwood CK, Wilson SK, Hayes PE, et al. Single-dose bioavailability of two extended-release lithium carbonate products. *Am J Hosp Pharm*. 1994;51:486-489. 4. Cooper TB, Simpson GM, Lee JH, Bergner P-EE. Evaluation of a slow-release lithium carbonate formulation. *Am J Psychiatry*. 1978;135(8):917-922.

On the high wire
of mania...



FOR A SAFE, SMOOTH RETURN TO A MORE NORMAL LIFE...

Smooth, slow release of lithium carbonate for initial or maintenance treatment of mania associated with bipolar disorder

- Smoother blood levels may reduce side effects^{1,2}
 - Helps minimize peak-to-trough variations in serum lithium concentrations
 - Common side effects that may occur during initial therapy include fine hand tremor, polyuria, mild thirst, and transient and mild nausea. These side effects usually subside with continued treatment, temporary reduction of dosage, or cessation.
- Interchangeable with immediate-release lithium preparations on a mg-to-mg basis¹⁻⁴


- Film-coated tablets eliminate metallic taste concerns
- B.I.D. convenience may enhance patient compliance

LITHOBID[®]
Slow-Release Tablets, 300 mg

WARNING: Lithium toxicity is closely related to serum lithium levels, and can occur at doses close to therapeutic levels. Facilities for prompt and accurate serum lithium determinations should be available before initiating therapy.

Please see brief summary of prescribing information on adjacent page.

ESTABLISHED THERAPY FOR OCD



EFFECTIVE CONTROL OF OBSESSIONS AND COMPULSIONS^{1*}

LOW INCIDENCE OF AGITATION
(2% vs 1% for placebo)¹

LOW INCIDENCE OF SEXUAL DYSFUNCTION¹

- ❖ LUVOX[®] Tablets vs placebo[†]: decreased libido 2% vs 1%; delayed ejaculation 8% vs 1%; anorgasmia 2% vs 0%; impotence 2% vs 1%

FAVORABLE SAFETY PROFILE

- ❖ Relatively low incidence of anticholinergic side effects in controlled trials of OCD and depression, LUVOX[®] Tablets vs placebo¹: dizziness 11% vs 6%; constipation 10% vs 8%; dry mouth 14% vs 10%
- ❖ The most commonly observed adverse events compared to placebo were somnolence 22% vs 8%, insomnia 21% vs 10%, nervousness 12% vs 5%, nausea 40% vs 14%, abnormal ejaculation 8% vs 1%, asthenia 14% vs 6%¹
- ❖ Concomitant use of LUVOX[®] Tablets and monoamine oxidase inhibitors is not recommended¹

FLEXIBLE DOSING

Initial Dose: 50 mg once a day HS
Dose Range: 100 to 300 mg/day

COMPREHENSIVE SAFETY DATABASE

(Worldwide Exposure for Reporting Overdose[‡])¹

- ❖ Data from 40 countries
- ❖ Over 12 million patients treated
- ❖ More than 37,000 patients studied in clinical trials

LUVOX[®]
fluvoxamine maleate 50 mg & 100 mg
SCORED TABLETS

**A SELECTIVE SEROTONIN
REUPTAKE INHIBITOR**

*Effectiveness not established beyond 10 weeks in controlled trials.

[†]Parameters occurring $\geq 1\%$ with fluvoxamine maleate.

[‡]Prescribers should write the smallest tablet quantity consistent with good patient management to reduce overdose risk.

Please see brief summary of prescribing information on adjacent page.