

HALL CLASSES OF GROUPS WITH A LOCALLY FINITE OBSTRUCTION

F. DE GIOVANNI , M. TROMBETTI  and B. A. F. WEHRFRITZ

(Received 5 January 2023; accepted 19 June 2023; first published online 24 July 2023)

Communicated by Benjamin Martin

Abstract

A well-known theorem of Philip Hall states that if a group G has a nilpotent normal subgroup N such that G/N is nilpotent, then G itself is nilpotent. We say that a group class \mathfrak{X} is a *Hall class* if it contains every group G admitting a nilpotent normal subgroup N such that G/N belongs to \mathfrak{X} . Hall classes have been considered by several authors, such as Plotkin [‘Some properties of automorphisms of nilpotent groups’, *Soviet Math. Dokl.* **2** (1961), 471–474] and Robinson [‘A property of the lower central series of a group’, *Math. Z.* **107** (1968), 225–231]. A further detailed study of Hall classes is performed by us in another paper [‘Hall classes of groups’, to appear] and we also investigate the behaviour of the class of finite-by- \mathfrak{Y} groups for a given Hall class \mathfrak{Y} [‘Hall classes in linear groups’, to appear]. The aim of this paper is to prove that for most natural choices of the Hall class \mathfrak{Y} , also the classes $(\mathbf{LF})\mathfrak{Y}$ and $\mathfrak{B}\mathfrak{Y}$ are Hall classes, where \mathbf{LF} is the class of locally finite groups and \mathfrak{B} is the class of locally finite groups of finite exponent.

2020 *Mathematics subject classification*: primary 20F18; secondary 20F19, 20F50.

Keywords and phrases: Hall class, nilpotent group, locally finite group.

1. Introduction

If N is a nilpotent normal subgroup of a group G , the structure of the factor group G/N has a strong influence on the whole group G . The first evidence of this phenomenon can be traced back to a classical theorem of Hall [11] which states that if G/N is nilpotent, then G itself is nilpotent. A few years later, Plotkin [15] gave a different proof of Hall’s nilpotency criterion and proved that if G/N is locally nilpotent, then G is locally nilpotent too, a fact that is not a direct consequence of the previous theorem. In 1968, Robinson [16] showed that this kind of result is strictly connected to the behaviour of certain tensor products of G -modules; this approach makes it possible to obtain similar statements for many other relevant group properties.

© The Author(s), 2023. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.



Along these lines, we say that a group class \mathfrak{X} is a *Hall class* if it contains every group G admitting a nilpotent normal subgroup N such that G/N' belongs to \mathfrak{X} . Clearly, finite groups and locally cyclic groups form Hall classes, and it is also easy to see that any class consisting only of groups with a locally cyclic Fitting subgroup is a Hall class.

In our terminology, the theorems of Hall and Plotkin just say that the class \mathfrak{N} of nilpotent groups and the class $L\mathfrak{N}$ of locally nilpotent groups are Hall classes. Moreover, it follows from the result of Robinson that hypercentral groups, hypercyclic groups, (locally) supersoluble groups and (locally) polycyclic groups form Hall classes; also, it turns out that a group class \mathfrak{X} that is closed with respect to forming normal subgroups and extensions is a Hall class, provided that it contains all homomorphic images of the tensor product of any pair of abelian \mathfrak{X} -groups. More recently, it was proved that the class of paranilpotent groups is a Hall class (see [10]), and also that FC -nilpotent, FC -hypercentral and locally FC -nilpotent groups form Hall classes (see [1]).

Since any Hall class containing all abelian groups must contain \mathfrak{A} , it is clear that for every positive integer c , the class of nilpotent groups of class at most c is not a Hall class, so in particular abelian groups do not form a Hall class. We remark also that the class of torsion-free groups is not a Hall class, although the class of locally finite groups and the class of groups of finite exponent are Hall classes.

It is obvious that unions and intersections of collections of Hall classes are likewise Hall classes. However, if \mathfrak{X} and \mathfrak{Y} are Hall classes, the class $\mathfrak{X}\mathfrak{Y}$, consisting of all groups G containing a normal \mathfrak{X} -subgroup N such that G/N is in \mathfrak{Y} , need not be a Hall class. For instance, the class $\mathfrak{F}\mathfrak{N}$ of all finite-by-nilpotent groups is not a Hall class; actually, for many natural choices of the Hall class \mathfrak{Y} , groups that are finite-by- \mathfrak{Y} do not form a Hall class (see [7], where many examples of this circumstance can be found).

The aim of this paper is to prove that the situation is completely different if the finite obstruction at the bottom is replaced by a locally finite one. The following summarizes very briefly most of our main conclusions.

THEOREM. *The class of all (locally finite)-by- \mathfrak{Y} groups and the class of all (locally finite of finite exponent)-by- \mathfrak{Y} groups are Hall classes for the following 18 choices for \mathfrak{Y} : nilpotent, Fitting, hypercentral, Engel, Baer, Gruenberg, locally nilpotent, paranilpotent, hypercyclic, locally supersoluble, FC -nilpotent, FC -hypercentral, locally (nilpotent-by-finite), soluble, hyperabelian, locally soluble, finite (Prüfer) rank, finite abelian section rank.*

Note that (locally finite)-by-nilpotent groups and (locally finite of finite exponent)-by-nilpotent groups were already proved to form Hall classes in [9, 18], respectively, although with different proofs.

Our notation, which is mostly standard, can be found in [17]; we refer to [20] for results and terminology concerning linear groups.

2. Nilpotency conditions

As mentioned in the introduction, nilpotent groups, locally nilpotent groups and hypercentral groups form Hall classes; in addition, it is known that Baer groups, Gruenberg groups, Fitting groups and Engel groups have the same behaviour (see [7]). The goal of this section is to prove that if \mathfrak{Y} is any of these seven group classes, then $(\mathbf{L}\mathfrak{F})\mathfrak{Y}$ is likewise a Hall class, where $\mathbf{L}\mathfrak{F}$ denotes the class of all locally finite groups.

Let G be a group. If A and B are G -modules, the tensor product $A \otimes_{\mathbb{Z}} B$ can be made into a G -module via $(a \otimes b)g = (ag) \otimes (bg)$ for all $a \in A, b \in B, g \in G$. Recall that a class \mathcal{M} of G -modules is *tensorial* if, for all A and B in \mathcal{M} , every G -homomorphic image of $A \otimes_{\mathbb{Z}} B$ (and in particular $A \otimes_{\mathbb{Z}} B$ itself) belongs to \mathcal{M} . It is well known that if N is a normal subgroup of G , for each positive integer i , there exists a G -epimorphism

$$(N/N') \otimes_{\mathbb{Z}} (\gamma_i(N)/\gamma_{i+1}(N)) \longrightarrow \gamma_{i+1}(N)/\gamma_{i+2}(N),$$

where the factors of the lower central series of N are regarded as G -modules by conjugation. This is the reason for which tensorial classes play a relevant role in the context of Hall classes. For instance, Hall's nilpotency criterion rests upon the fact that the class of *polytrivial G -modules* (that is, G -modules admitting a finite series of G -submodules whose factors are trivial G -modules) is tensorial (see for instance [21, page 10]).

If G is a group and if \mathcal{M}, \mathcal{N} are classes of G -modules, we say that a G -module A is *\mathcal{M} -by- \mathcal{N}* if it has a G -submodule $B \in \mathcal{M}$ such that A/B is in \mathcal{N} . Moreover, in the following, we denote by \mathcal{T}_G the class of all G -modules whose underlying abelian groups are periodic.

LEMMA 2.1. *Let G be a group and let \mathcal{M} be a tensorial class of G -modules that is quotient-closed. Then the class of all G -modules that are \mathcal{T}_G -by- \mathcal{M} is tensorial.*

PROOF. Clearly, the class \mathcal{T}_G -by- \mathcal{M} is quotient-closed. Let A and B be two arbitrary G -modules that are \mathcal{T}_G -by- \mathcal{M} , and let T_A and T_B be the periodic parts of A and B (as abelian groups), respectively. The natural sequence

$$(T_A \otimes_{\mathbb{Z}} B) \oplus (A \otimes_{\mathbb{Z}} T_B) \longrightarrow A \otimes_{\mathbb{Z}} B \longrightarrow (A/T_A) \otimes_{\mathbb{Z}} (B/T_B) \longrightarrow \{0\}$$

is exact, where the G -submodule

$$(T_A \otimes_{\mathbb{Z}} B) \oplus (A \otimes_{\mathbb{Z}} T_B)$$

is easily seen to be in \mathcal{T}_G and

$$(A/T_A) \otimes_{\mathbb{Z}} (B/T_B)$$

is in \mathcal{M} since A/T_A and B/T_B are. Therefore, the G -module $A \otimes_{\mathbb{Z}} B$ is \mathcal{T}_G -by- \mathcal{M} and the statement is proved. □

COROLLARY 2.2. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by-polytrivial is tensorial.*

A classical theorem of Issai Schur states that if a group G is (locally) finite over its centre $\zeta_1(G)$, then G' is (locally) finite (see [17, Part 1, Theorem 4.12 and its Corollary]). For our purposes, we need the following application of Schur's theorem.

LEMMA 2.3. *Let G be a group and let N be a normal subgroup of G . If N has a finite G -invariant series whose factors are either locally finite or central in G , then the subgroup $[N, {}_k G]$ is locally finite, where k is the number of factors of this series that are not locally finite.*

PROOF. Let

$$\{1\} = N_0 < N_1 < \cdots < N_t = N$$

be a finite G -invariant series of minimal length t such that each N_{i+1}/N_i is either locally finite or central in G and at most k factors are not locally finite. Of course, we may suppose that G has no nontrivial locally finite normal subgroups and $k > 0$, so that in particular $N_1 \leq \zeta_1(G)$. By induction, we have that

$$M/N_1 = [N, {}_{k-1} G]N_1/N_1$$

is locally finite. Then $M/\zeta_1(M)$ is locally finite, so M' is locally finite by Schur's theorem and hence M is torsion-free abelian. Since $N_1 \leq \zeta_1(G)$ and M/N_1 is locally finite, it follows that M is central in G and hence $[N, {}_k G] \leq [M, G] = \{1\}$. \square

The choice $N = G$ in the above statement gives the following characterization of (locally finite)-by-nilpotent groups.

COROLLARY 2.4. *A group G is (locally finite)-by-nilpotent if and only if it has a normal series of finite length whose factors are either locally finite or central in G .*

We are now in a position to prove the first main theorem of this section.

THEOREM 2.5. *The class $(\mathbf{LF})\mathfrak{N}$ of all (locally finite)-by-nilpotent groups is a Hall class.*

PROOF. Let N be a nilpotent normal subgroup of a group G such that G/N' is in $(\mathbf{LF})\mathfrak{N}$. If T/N' is the periodic part of N/N' , then N/T has a finite G -invariant series whose factors are central in G and hence N/N' is \mathcal{T}_G -by-polytrivial as a G -module (by conjugation). Application of Corollary 2.2 yields that N has a finite G -invariant series whose nonperiodic factors are central in G . It follows that G has a normal series of finite length whose factors are either locally finite or central, and hence it is (locally finite)-by-nilpotent by Corollary 2.4. \square

Philip Hall proved that if N is a nilpotent normal subgroup of a group G and G/N' is nilpotent, then the nilpotency class of G can be bounded in terms of the nilpotency classes of N and G/N' . Note here that the above proof allows us to give a quantitative version of Theorem 2.5 too.

THEOREM 2.6. *There exists an integer-valued function $f = f(c, d)$ such that, if G is any group containing a nilpotent normal subgroup N of class c such that $\gamma_{d+1}(G/N')$ is locally finite, then $\gamma_{f(c,d)}(G)$ is locally finite.*

Recall that a group G is a *Fitting group* if it is generated by its nilpotent normal subgroups. Of course, Fitting groups are locally nilpotent and it follows easily from Hall’s theorem that they form a Hall class (see [7] for details).

COROLLARY 2.7. *The class of all (locally finite)-by-Fitting groups is a Hall class.*

PROOF. Let G be a group containing a nilpotent normal subgroup N such that G/N' is (locally finite)-by-Fitting and let T be the unique maximal locally finite normal subgroup of G . If L/N' is the subgroup of all elements of finite order of G/N' and K/L is any nilpotent normal subgroup of G/L , we have that KN/N' is (locally finite)-by-nilpotent and hence also the normal subgroup KN of G is (locally finite)-by-nilpotent by Theorem 2.5. Thus, KN/T is nilpotent, so that G/T is a Fitting group and G is (locally finite)-by-Fitting. □

The main ingredients of the proof of Theorem 2.5 are essentially the tensoriality of \mathcal{T}_G -by-polytrivial G -modules (Corollary 2.2) and the sufficient part of the characterization of (locally finite)-by-nilpotent groups given in Corollary 2.4. Actually, many of the most relevant classes of groups with a locally finite obstruction at the bottom can be proved to be Hall classes, provided that results corresponding to Corollaries 2.2 and 2.4 are obtained (it is then enough to repeat the argument of the above proof).

If G is a group and τ is an ordinal number, a G -module A is called *hypertrivial (of height τ)* if it has an ascending series of G -submodules

$$\{0\} = A_0 < A_1 < \cdots < A_\alpha < A_{\alpha+1} < \cdots < A_\tau = A$$

such that $A_{\alpha+1}/A_\alpha$ is a trivial G -module for each ordinal $\alpha < \tau$. Since the class of hypertrivial G -modules is tensorial (see [16, Lemma 1]), Lemma 2.1 has the following consequence, which corresponds to Corollary 2.2.

COROLLARY 2.8. *If G is a group, the class of all G -modules that are \mathcal{T}_G -hypertrivial is tensorial.*

In the following, if G is a group, $\bar{\zeta}(G)$ denotes the hypercentre of G , and we say that a normal section X/Y of G is *hypercentrally embedded* in G if $X/Y \leq \bar{\zeta}(G/Y)$.

LEMMA 2.9. *Let G be a group and let N be a torsion-free normal subgroup of G such that $N\bar{\zeta}(G)/\bar{\zeta}(G)$ is locally finite. Then N is contained in $\bar{\zeta}(G)$.*

PROOF. Since $N/\bar{\zeta}(N)$ is obviously locally finite, N contains a locally finite normal subgroup M such that N/M is hypercentral (see [3, Lemma 2]). However, N is torsion-free and hence it is hypercentral. Assume for a contradiction that the statement is false and let μ be the least ordinal number such that $\zeta_\mu(N)$ is not contained in $\bar{\zeta}(G)$. Then $\mu > 0$ is not a limit and $\zeta_{\mu-1}(N) \leq \bar{\zeta}(G)$. Since $N/\zeta_{\mu-1}(N)$ is torsion-free

(see [17, Part 1, Theorem 2.25]), we may replace G by $G/\zeta_{\mu-1}(N)$ and N by $\zeta_{\mu}(N)/\zeta_{\mu-1}(N)$, and so assume without loss of generality that N is abelian.

Since $N/(N \cap \bar{\zeta}(G))$ is locally finite, we may consider the least ordinal number σ such that $N/(N \cap \zeta_{\sigma}(G))$ is not torsion-free. Clearly, $\sigma > 0$ is not a limit and $N/(N \cap \zeta_{\sigma-1}(G))$ is torsion-free. Thus, a further replacement of G by the factor group $G/\zeta_{\sigma-1}(G)$ allows us to assume that $N/(N \cap \zeta_1(G))$ is not torsion-free. Then there is an element x of $N \setminus \zeta_1(G)$ such that $x^k \in \zeta_1(G)$ for some positive integer k . If g is any element of G ,

$$[x, g]^k = [x^k, g] = 1,$$

so $[x, g] = 1$ because N is torsion-free and hence $x \in \zeta_1(G)$. This contradiction completes the proof of the statement. \square

LEMMA 2.10. *Let G be a group and let N be a normal subgroup of G . If N has a G -invariant series of finite length each of whose factors is either locally finite or hypercentrally embedded in G , then N contains a locally finite G -invariant subgroup T such that N/T is hypercentrally embedded in G .*

PROOF. Let

$$\{1\} = N_0 < N_1 < \cdots < N_t = N$$

be a G -invariant series of finite length whose factors are either locally finite or hypercentrally embedded in G . Without loss of generality, we may assume that N has no nontrivial locally finite normal subgroups, so in particular $N_1 \leq \bar{\zeta}(G)$, and also that N_2/N_1 is locally finite. Then $N_2/\bar{\zeta}(N_2)$ is locally finite and hence it follows from [3, Lemma 2] that N_2 is hypercentral, and so also torsion-free. Application of Lemma 2.9 yields now that $N_2 \leq \bar{\zeta}(G)$. It follows that N has a G -invariant series of the above type whose length is strictly smaller than t and hence the statement is proved by induction. \square

The above result allows us to characterize (locally finite)-by-hypercentral groups in a way similar to that in which (locally finite)-by-nilpotent groups are described in Corollary 2.4.

COROLLARY 2.11. *A group G is (locally finite)-by-hypercentral if and only if it has a normal series of finite length whose factors are either locally finite or hypercentrally embedded in G .*

It is now possible to state the following result, whose proof runs along the same lines of that of Theorem 2.5.

THEOREM 2.12. *The class of all (locally finite)-by-hypercentral groups is a Hall class.*

We turn now to the case in which \mathfrak{V} is the class of Engel groups. Recall here that an element x of a group G is *right Engel* if for each $g \in G$, there is a positive integer $m = m(x, g)$ such that $[x, {}_m g] = 1$, and G is called an *Engel group* if all its elements are right Engel.

Let G be a group and let A be a G -module. We say that A is an *Engel G -module* if for all elements a of A and g of G , there exists a positive integer $m = m(a, g)$ such that $a(g - 1)^m = 0$. Clearly, if A is an abelian normal subgroup of a group G that is an Engel G -module by conjugation, then all elements of A are right Engel in G .

The class of all Engel G -modules is closed with respect to forming submodules, quotient modules and extensions, and it is easy to show that this class of modules is tensorial (see [7]). Thus, Lemma 2.1 above has the following further consequence.

COROLLARY 2.13. *If G is a group, the class of all G -modules which are \mathcal{T}_G -by-Engel is tensorial.*

LEMMA 2.14. *Let G be a group and let A be a G -module that is torsion-free as an abelian group. If A contains an Engel G -submodule B such that $A/B \in \mathcal{T}_G$, then A is an Engel G -module.*

PROOF. Let a and g be arbitrary elements of A and G , respectively, and let $k = k(a)$ be a positive integer such that $ka \in B$. Then there exists a positive integer $m = m(a, g)$ such that $(ka)(g - 1)^m = 0$, so $k(a(g - 1)^m) = 0$ and hence $a(g - 1)^m = 0$. Therefore, A is an Engel G -module. □

LEMMA 2.15. *Let G be a group and let N be a normal subgroup of G . If N has a G -invariant series of finite length each of whose factors is either locally finite or an Engel G -module (by conjugation), then N contains a locally finite G -invariant subgroup T such that N/T consists of right Engel elements of G/T .*

PROOF. Let

$$\{1\} = N_0 < N_1 < \dots < N_t = N$$

be a G -invariant series of finite length whose factors are either locally finite or Engel G -modules. Without loss of generality, it can be assumed that N has no locally finite nontrivial normal subgroups, so in particular N_1 is an Engel G -module that is torsion-free as an abelian group. Moreover, by induction on t , we may also suppose that N/N_1 contains a locally finite G -invariant subgroup M/N_1 such that N/M consists of right Engel elements of G/M . Let E be an arbitrary finitely generated subgroup of M and put $A = E \cap N_1$. Then E/A is finite and hence A is finitely generated. Since N_1 is an Engel G -module, for each element x of E , there exists a positive integer k such that $[A, {}_k x] = \{1\}$. It follows that the subgroup $\langle x, A \rangle$ is nilpotent and so even abelian (see for instance [17, Part 2, Lemma 6.37] or [2, Proposition 2.1.3]). Therefore, $A \leq \zeta_1(E)$ and so $N_1 \leq \zeta_1(M)$. In particular, $M/\zeta_1(M)$ is locally finite and hence M is torsion-free abelian by Schur’s theorem. Application of Lemma 2.14 yields now that M is an Engel G -module. However, each element of N/M is right Engel in G/M and so all elements of N are right Engel in G . The statement is proved. □

COROLLARY 2.16. *Let G be a group admitting a normal series of finite length each of whose factors is either locally finite or an Engel G -module. Then G is (locally finite)-by-Engel.*

THEOREM 2.17. *The class of all (locally finite)-by-Engel groups is a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N' is (locally finite)-by-Engel. Then N/N' is \mathcal{T}_G -by-Engel as a G -module and so by Corollary 2.13, we have that N has a finite G -invariant series whose nonperiodic subgroups are Engel G -modules. If L/N' is the unique maximal locally finite normal subgroup of G/N' , it follows that L has a G -invariant series of finite length each of whose factors is either locally finite or an Engel G -module. Application of Lemma 2.15 yields that L contains a locally finite G -invariant subgroup T such that L/T consists of right Engel elements of G/T . Therefore, G is (locally finite)-by-Engel and the statement is proved. \square

A closer look at the above arguments allows us to state the following result concerning bounded Engel groups; recall here that a group G is *bounded Engel* if there exists a positive integer m such that $[x,{}_m y] = 1$ for all x, y in G .

THEOREM 2.18. *The class of all (locally finite)-by-(bounded Engel) groups is a Hall class.*

Let G be a group. Recall that G is a *Baer group* if all its cyclic subgroups are subnormal, while G is a *Gruenberg group* if every cyclic subgroup of G is ascendant. It is known that G is a Baer group (respectively a Gruenberg group) if it is generated by abelian subnormal (respectively abelian ascendant) subgroups. Of course, Baer groups are Gruenberg and every Gruenberg group is locally nilpotent. Notice also that a soluble group is Baer (respectively Gruenberg) if and only if it is bounded Engel (respectively Engel); for these results, see [17, Part 2, Theorems 7.34 and 7.35]. Thus it follows from Theorems 2.18 and 2.17 that soluble groups that are (locally finite)-by-Baer and soluble groups that are (locally finite)-by-Gruenberg form Hall classes. Actually, we can prove that the solubility assumption can be dropped.

Let \mathfrak{X} be a group class. As usual, we denote by $\mathbf{S}_n\mathfrak{X}$ (respectively, $\mathbf{Q}\mathfrak{X}$) the class of all groups that are isomorphic to subnormal subgroups (respectively to factor groups) of \mathfrak{X} -groups. Of course, $\mathbf{S}_n\mathfrak{X} = \mathfrak{X}$ if and only if \mathfrak{X} is closed with respect to forming normal subgroups and $\mathbf{Q}\mathfrak{X} = \mathfrak{X}$ if and only if all homomorphic images of \mathfrak{X} -groups belong to \mathfrak{X} .

LEMMA 2.19. *Let \mathfrak{X} and \mathfrak{Y} be group classes such that $\mathbf{S}_n\mathfrak{X} = \mathfrak{X}$, $\mathbf{Q}\mathfrak{Y} = \mathfrak{Y}$ and $\mathfrak{X}\mathfrak{Y}$ is a Hall class. If G is a group containing a nilpotent normal subgroup N with $G/N' \in \mathfrak{X}\mathfrak{Y}$, then N is contained in a normal subgroup M of G such that $G/M \in \mathfrak{Y}$ and $\gamma_k(M) \in \mathfrak{X}$ for some positive integer k .*

PROOF. Let X/N' be a normal \mathfrak{X} -subgroup of G/N' such that $G/X \in \mathfrak{Y}$. Clearly, the group XN/N' is \mathfrak{X} -by-abelian, so the normal subgroup $M = XN$ is \mathfrak{X} -by-nilpotent since $\mathfrak{X}\mathfrak{Y}$ is a Hall class. Thus, there is a positive integer k such that the subgroup $\gamma_k(M)$ belongs to $\mathbf{S}_n\mathfrak{X} = \mathfrak{X}$. Moreover, G/M belongs to $\mathbf{Q}\mathfrak{Y} = \mathfrak{Y}$ and the statement is proved. \square

THEOREM 2.20. *The class of all (locally finite)-by-Baer groups and the class of all (locally finite)-by-Gruenberg groups are Hall classes.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N is (locally finite)-by-Baer. To prove that G itself is (locally finite)-by-Baer, we may suppose that G has no nontrivial locally finite normal subgroups. Thus, a combination of Theorem 2.5 and Lemma 2.19 with $\mathfrak{X} = \mathbf{L}\mathfrak{F}$ shows that there exists a nilpotent normal subgroup M of G such that G/M is (locally finite)-by-Baer and G/M is a Baer group. If g is an arbitrary element of G , the subgroup $\langle g, M \rangle$ is subnormal in G and $\langle g, M \rangle/M$ is (locally finite)-by-nilpotent. Thus, $\langle g, M \rangle$ is (locally finite)-by-nilpotent by Theorem 2.5, and so even nilpotent because G has no nontrivial locally finite normal subgroups. It follows that $\langle g \rangle$ is subnormal in G and hence G is a Baer group.

To prove that (locally finite)-by-Gruenberg groups form a Hall class, we can use a similar argument, replacing subnormality by ascendancy and Theorem 2.5 by Theorem 2.12. \square

To accomplish what we promised at the beginning of this section, it only remains to show that $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$ is a Hall class.

If G is a group and \mathcal{M} is a class of G -modules, we denote by $\mathbf{L}\mathcal{M}$ the class of all G -modules A such that for each finite subset X of A , there is an \mathcal{M} -submodule of A containing X . The G -modules in the class $\mathbf{L}\mathcal{M}$ are called *locally* \mathcal{M} .

LEMMA 2.21. *Let G be a group and let \mathcal{M} be a tensorial class of G -modules. If \mathcal{M} is closed with respect to forming G -sections, then the class $\mathbf{L}\mathcal{M}$ is tensorial.*

PROOF. Suppose that A and B are G -modules in the class $\mathbf{L}\mathcal{M}$, and let E be a finitely generated G -submodule of $A \otimes_{\mathbb{Z}} B$. Clearly, there exist finitely generated submodules A_0 of A and B_0 of B such that E is G -isomorphic to a G -section of $A_0 \otimes_{\mathbb{Z}} B_0$. Since A_0 and B_0 belong to the tensorial class \mathcal{M} , also E is in \mathcal{M} . Therefore, $A \otimes_{\mathbb{Z}} B$ belongs to $\mathbf{L}\mathcal{M}$ and hence $\mathbf{L}\mathcal{M}$ is tensorial. \square

Since the class of polytrivial G -modules is tensorial, it follows from Lemma 2.21 that also locally polytrivial G -modules form a tensorial class and therefore Lemma 2.1 has the following consequence.

COROLLARY 2.22. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by-(locally polytrivial) is tensorial.*

The following statement and the subsequent remark describe the relation between locally polytrivial and hypertrivial modules; this is employed in the proof of our next main result.

LEMMA 2.23. *Let G be a group and let A be a G -module. Then A is locally polytrivial if and only if A is hypertrivial of height at most ω .*

PROOF. Assume first that A is locally polytrivial and let \mathcal{E} be the collection of all finitely generated G -submodules of A . Each $E \in \mathcal{E}$ admits a finite series of G -submodules

$$\{0\} = E_0 < E_1 < \dots < E_{k(E)} = E$$

whose factors are trivial G -modules. For each nonnegative integer n , let A_n be the G -submodule of A generated by all F_n , where $F \in \mathcal{E}$ and $k(F) \geq n$. Then,

$$\{0\} = A_0 \leq A_1 \leq \dots \leq A_n \leq A_{n+1} \leq \dots \bigcup_{n \in \mathbb{N}} A_n = A$$

is an ascending series of G -submodules of A whose factors are trivial G -modules. Therefore, A is hypertrivial of height at most ω .

Conversely, let

$$\{0\} = A_0 \leq A_1 \leq \dots \leq A_n \leq A_{n+1} \leq \dots A_\omega = A$$

be an ascending series of G -submodules of A such that every A_{n+1}/A_n is a trivial G -module. If E is any finitely generated G -submodule of A , there is a nonnegative integer n such that A_n contains E . Then E is a polytrivial G -module and the statement is proved. □

Lemma 2.23 shows that the class of locally polytrivial G -modules is contained in that of hypertrivial G -modules, and this inclusion is strict, a property which is in contrast to the well-known fact that hypercentral groups form a proper subclass of $\mathcal{L}\mathfrak{N}$. This can be seen from the following example. Let p be a prime and let

$$A = \langle a \rangle \times \langle b_1 \rangle \times \langle b_2 \rangle \times \dots \times \langle b_n \rangle \times \dots$$

be the direct product of infinitely many groups of order p ; consider the automorphisms x and y_k (for each $k > 1$) of A defined by the positions

$$a^x = a, \quad b_1^x = b_1, \quad b_n^x = b_n b_{n-1} \quad \text{for all } n > 1$$

and

$$a^{y_k} = ab_k, \quad b_n^{y_k} = b_n \quad \text{for all } n \geq 1.$$

If $G = \langle x, y_k \mid k \geq 1 \rangle$, we have that $A = \langle a \rangle^G$ is a cyclic G -module, which is hypertrivial of height $\omega + 1$ but not polytrivial.

THEOREM 2.24. *The class $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$ of all (locally finite)-by-(locally nilpotent) groups is a Hall class.*

PROOF. Let N be a nilpotent normal subgroup of the group G such that G/N' belongs to $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$. To prove that $G \in (\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$, we may assume that G has no nontrivial locally finite normal subgroups, so in particular N is torsion-free. Consider any finitely generated subgroup E of G and let T/N be the subgroup of all elements of finite order of EN/N , so that EN/T is nilpotent. Since N/N' is \mathcal{T}_G -by-(locally polytrivial) as an E -module, it follows from Corollary 2.22 that N (and so also T) has an E -invariant series of finite length whose factors are either locally finite or locally polytrivial

E -modules. However, locally polytrivial E -modules are hypertrivial by Lemma 2.23 and hence an application of Lemma 2.10 yields that T is hypercentrally embedded in EN . Thus, EN is hypercentral and so G is locally nilpotent. \square

If \mathfrak{X} is any group class, $\mathbf{L}\mathfrak{X}$ denotes the class of all *locally \mathfrak{X} -groups*, that is, groups each of whose finite subsets lies in an \mathfrak{X} -subgroup; of course, if \mathfrak{X} is subgroup-closed, a group G is locally \mathfrak{X} if and only if all its finitely generated subgroups belong to \mathfrak{X} .

It is known that $\mathbf{L}(\mathfrak{N}\mathfrak{F})$ is a Hall class (see for instance [1, Proposition 3.1]) and it has been recently proved that also $\mathbf{L}(\mathfrak{F}\mathfrak{N})$ is a Hall class (see [7]). Since the intersection of Hall classes is likewise a Hall class, a combination of Theorem 2.24 and our next lemma provides a different proof of this latter fact.

LEMMA 2.25. $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N}) = \mathbf{L}((\mathbf{L}\mathfrak{F})\mathfrak{N})$ and $\mathbf{L}(\mathfrak{F}\mathfrak{N}) = ((\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})) \cap \mathbf{L}(\mathfrak{N}\mathfrak{F})$.

PROOF. It is obvious that $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N}) \leq \mathbf{L}((\mathbf{L}\mathfrak{F})\mathfrak{N})$. Conversely, if G is any group in $\mathbf{L}((\mathbf{L}\mathfrak{F})\mathfrak{N})$, the elements of finite order of G form a locally finite subgroup T and G/T is locally nilpotent. Thus, $\mathbf{L}((\mathbf{L}\mathfrak{F})\mathfrak{N}) \leq (\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$ and the first equality is proved.

To prove the second equality, note that

$$\mathbf{L}(\mathfrak{F}\mathfrak{N}) \leq \mathbf{L}((\mathbf{L}\mathfrak{F})\mathfrak{N}) = (\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$$

by the first part of the proof. Since every finite-by-nilpotent group is obviously nilpotent-by-finite, we have also $\mathbf{L}(\mathfrak{F}\mathfrak{N}) \leq \mathbf{L}(\mathfrak{N}\mathfrak{F})$ and hence

$$\mathbf{L}(\mathfrak{F}\mathfrak{N}) \leq ((\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})) \cap \mathbf{L}(\mathfrak{N}\mathfrak{F}).$$

Suppose now that G is a group in $((\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})) \cap \mathbf{L}(\mathfrak{N}\mathfrak{F})$ and let E be any finitely generated subgroup of G . Since E is nilpotent-by-finite, it satisfies the maximal condition; then E is finite-by-nilpotent because it belongs to $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$. The proof is complete. \square

We give now an example which shows in particular that (locally finite)-by-(finitely generated nilpotent) groups do not form a Hall class.

THEOREM 2.26. *There exists a torsion-free nilpotent linear group G over a field of characteristic 0 that is not finitely generated, but G/G' is periodic-by-(finitely generated abelian).*

PROOF. Let $\text{Tr}_1(3, \mathbb{Q})$ be the group of all unitriangular matrices of order 3 over the rationals, and let $G \leq \text{Tr}_1(3, \mathbb{Q})$ be the group generated by the matrices

$$x = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \quad \text{and} \quad z(a) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & 0 & 1 \end{pmatrix},$$

where a ranges over all rational numbers. Clearly $[y, x] = z(1)$, so

$$G' = \langle z(1) \rangle \quad \text{and} \quad G/G' \simeq \mathbb{Z} \times \mathbb{Z} \times \mathbb{Q}/\mathbb{Z}$$

is periodic-by-(finitely generated abelian). However, G is a torsion-free nilpotent group that is not finitely generated. \square

Although the above example is linear, it turns out that the situation is completely different in the case of a positive characteristic; this is essentially due to an intrinsic property of linear groups of positive characteristic. If G is a linear group, we denote by $u(G)$ the *unipotent radical* of G , that is, its unique maximal unipotent normal subgroup.

LEMMA 2.27. *Let G be a linear group over a field of positive characteristic and let N be a nilpotent normal subgroup of G . Then N' is locally finite.*

PROOF. Put $U = u(G)$. Since G has positive characteristic, U is locally finite and G/U is isomorphic to a completely reducible linear group (see for instance [6, Lemma 2.13]). Then also NU/U is completely reducible (see [20, Corollary 1.8]) and hence, being nilpotent, it is central-by-finite (see [20]). It follows from Schur's theorem that $N'U/U$ is finite and hence N' is locally finite. \square

The above lemma suggests that certain classes of groups with a locally finite obstruction at the bottom can behave as Hall classes in suitable universes, even if they are not Hall classes. This idea can be formalized as follows.

If \mathfrak{U} is a fixed class of groups, we say that a group class \mathfrak{X} is a \mathfrak{U} -Hall class (or also a *Hall class within the universe \mathfrak{U}*) if it contains every \mathfrak{U} -group G admitting a nilpotent normal subgroup N such that $G/N' \in \mathfrak{X}$. Of course, if \mathfrak{U} is the class of all groups, \mathfrak{U} -Hall classes are just the usual Hall classes.

THEOREM 2.28. *Let \mathfrak{X} be any class of groups. Then $(\mathbf{L}\mathfrak{F})\mathfrak{X}$ is a Hall class within the universe of linear groups of positive characteristic.*

Notice that the concept of a \mathfrak{U} -Hall class is further generalized in Section 5, by imposing a restriction also to the nilpotent normal subgroup N in the above definition.

3. Supersolubility conditions

The aim of this section is to prove that $(\mathbf{L}\mathfrak{F})\mathfrak{V}$ is a Hall class, when \mathfrak{V} is one of the most natural generalizations of the class of supersoluble groups, like those of paranilpotent groups, of hypercyclic groups and of locally supersoluble groups.

Note preliminarily that (locally finite)-by-supersoluble groups do not form a Hall class (see Theorem 2.26). This is essentially due to the fact that (locally finite)-by-supersoluble groups cannot be characterized in a way similar to that of Corollary 2.4, replacing centrality by the property of being cyclic, an obvious example being the additive group of rational numbers.

We first deal with the class of (locally finite)-by-paranilpotent groups. Recall that a normal subgroup N of a group G is said to be *paranilpotently embedded* in G if there exists a finite series

$$\{1\} = N_0 \leq N_1 \leq \cdots \leq N_t = N$$

of normal subgroups of G such that, for $i = 0, \dots, t - 1$, the factor group N_{i+1}/N_i is *paracentral* in G , that is, it is abelian and all its subgroups are normal in G/N_i . A group G is *paranilpotent* if it is paranilpotently embedded in itself. Paranilpotent groups were introduced in [19] under the name of *parasoluble groups*, while the term paranilpotent group has recently been employed (see for instance [4, 5]). Of course, every nilpotent group is paranilpotent, while paranilpotent groups are locally supersoluble.

As in the previous section, we obtain results corresponding to Corollaries 2.2 and 2.4 for the class of (locally finite)-by-paranilpotent groups.

If G is any group, a G -module A is called *paranilpotent* if it has a finite series of G -submodules

$$\{0\} = A_0 < A_1 < \dots < A_t = A$$

such that every subgroup of A_{i+1}/A_i is a G -submodule for all $i = 0, 1, \dots, t - 1$. Of course, any abelian normal subgroup of a paranilpotent group G is a paranilpotent G -module by conjugation.

The class of all paranilpotent G -modules is easily seen to be tensorial and quotient-closed (see for instance [10]), so that Lemma 2.1 has the following consequence.

COROLLARY 3.1. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by-paranilpotent is tensorial.*

The following result corresponds to Lemma 2.14, bearing in mind that the only nontrivial automorphism fixing all subgroups of a nonperiodic abelian group is the inversion.

LEMMA 3.2. *Let G be a group, A a G -module that is torsion-free as an abelian group and B a G -submodule of A such that $A/B \in \mathcal{T}_G$. If g is an element of G centralizing B (respectively inverting B), then g centralizes A (respectively inverts A).*

PROOF. Let a be any element of A and let k be a positive integer such that $ka \in B$. Then, $(ka)g = \varepsilon(ka)$, where $\varepsilon \in \{1, -1\}$, so that $k(ag - \varepsilon a) = 0$ and hence $ag = \varepsilon a$ since A is torsion-free as an abelian group. □

LEMMA 3.3. *Let G be a group and let N be a normal subgroup of G . If N has a G -invariant series of finite length whose factors are either locally finite or paracentral in G , then N contains a locally finite G -invariant subgroup T such that N/T is paranilpotently embedded in G .*

PROOF. Let

$$\{1\} = N_0 < N_1 < \dots < N_t = N$$

be a G -invariant series of finite length whose factors are either locally finite or paracentral in G . Without loss of generality, we may assume that N has no locally finite nontrivial normal subgroups, so in particular N_1 is torsion-free and paracentral in G . Moreover, we may suppose by induction on t that N/N_1 contains a locally finite G -invariant subgrp M/N_1 such that N/M is paranilpotently embedded in G . Put

$C = C_M(N_1)$, so that M/C has order at most 2 and N/C is paranilpotently embedded in G . Clearly, $N_1 \leq \zeta_1(C)$, so $C/\zeta_1(C)$ is locally finite and hence C is abelian by Schur's theorem. An application of Lemma 3.2 yields that C is paracentral in G and hence G is paranilpotent. \square

COROLLARY 3.4. *A group G is (locally finite)-by-paranilpotent if and only if it has a normal series of finite length whose factors are either locally finite or paranilpotently embedded in G .*

The first main result of this section is a consequence of Corollaries 3.1 and 3.4.

THEOREM 3.5. *The class of all (locally finite)-by-paranilpotent groups is a Hall class.*

We move now from paranilpotent groups to *hypercyclic groups*, that is, groups having an ascending normal series with cyclic factors. In general, a normal section X/Y of a group G is *hypercyclically embedded* in G if it has an ascending G -invariant series with cyclic factors. Notice that if N is a hypercyclically embedded normal subgroup of G , then $N \cap G'$ is contained in the hypercentre of G' ; in particular, the commutator subgroup of any hypercyclic group is hypercentral. Moreover, it is easy to see that paranilpotent groups are hypercyclic and that hypercyclic groups are locally supersoluble.

Let G be a group. We say that a G -module A is *hypercyclic* if it has an ascending series of G -submodules

$$\{0\} = A_0 < A_1 < \cdots < A_\alpha < A_{\alpha+1} < \cdots < A_\tau = A$$

each of whose factors is cyclic as an abelian group; the least ordinal τ for which such a series exists is called the *height* of A . It is known that the class of all hypercyclic G -modules is tensorial (see [16]), so that Lemma 2.1 has the following consequence.

COROLLARY 3.6. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by-hypercyclic is tensorial.*

LEMMA 3.7. *Let G be a group and let N be a normal subgroup of G . If N has a G -invariant series of finite length each of whose factors is either locally finite or hypercyclically embedded in G , then N contains a locally finite G -invariant subgroup T such that N/T is hypercyclically embedded in G .*

PROOF. As in the proofs of Lemmas 2.10 and 3.3, we may suppose that N has no nontrivial locally finite normal subgroups and contains a G -invariant subgroup M such that M is hypercyclically embedded in G and N/M is locally finite. Then $M \cap G' \leq \bar{\zeta}(N \cap G')$, so that $(N \cap G')/\bar{\zeta}(N \cap G')$ is locally finite and hence $N \cap G'$ is hypercentral and torsion-free by Lemma 2.10. Put $Z = \zeta_1(N \cap G')$. By hypothesis, Z contains a nontrivial cyclic G -invariant subgroup and we may consider a subgroup A of Z that is maximal with respect to the condition that all its subgroups are normal in G ; obviously A is hypercyclically embedded in G . If B/A is the subgroup of all elements of finite order of Z/A , it follows from Lemma 3.2 that all subgroups of B are normal in G .

Thus, $A = B$, so Z/A is torsion-free and hence either $Z = A$ or Z/A contains a nontrivial cyclic G -invariant subgroup. This procedure shows that Z is hypercyclically embedded in G . Since all factors of the upper central series of $N \cap G'$ are torsion-free, we may employ the same argument to prove that all such factors are hypercyclically embedded in G . Therefore, $N \cap G'$ is hypercyclically embedded in G . Since $[N, G] \leq N \cap G'$, it follows that also N is hypercyclically embedded in G . \square

COROLLARY 3.8. *A group G is (locally finite)-by-hypercyclic if and only if it has a normal series of finite length whose factors are either locally finite or hypercyclically embedded in G .*

As in the previous cases, the following can be deduced.

THEOREM 3.9. *The class of all (locally finite)-by-hypercyclic groups is a Hall class.*

Finally, we consider the class of (locally finite)-by-(locally supersoluble) groups; we need the following easy lemma.

LEMMA 3.10. *Let G be a group whose finitely generated subgroups are (locally finite)-by-supersoluble. Then G is (locally finite)-by-(locally supersoluble).*

PROOF. We may obviously assume that G has no nontrivial locally finite normal subgroups. Since the commutator subgroup of any finitely generated subgroup of G is (locally finite)-by-nilpotent, G' belongs to $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{N})$ by Lemma 2.25 and hence G' is torsion-free and locally nilpotent. Let E be any finitely generated subgroup of G and let T be the unique maximal locally finite normal subgroup of E . Then E/T is supersoluble and $T \leq \zeta_1(E)$, because $T \cap E' = \{1\}$. Therefore, E is supersoluble and G is locally supersoluble. \square

In the proof of Theorem 2.24, a major role is played by locally polytrivial modules and hypertrivial modules. Here, this role is taken by locally paranilpotent modules and their relations with hypercyclic modules.

The example just before Theorem 2.24 shows that hypercyclic modules need not be locally paranilpotent, and in contrast to Lemma 2.23, we prove that locally paranilpotent modules are not in general hypercyclic of height at most ω . To see this, consider an uncountable free abelian group

$$A = \bigoplus_{i \in I} \langle a_i \rangle$$

and let

$$G = \text{Dr}_{i \in I} \langle g_i \rangle$$

be an elementary abelian 2-group acting on A according to the rules

$$a_i g_i = -a_i \quad \text{and} \quad a_i g_j = a_i \quad \text{if } i \neq j.$$

Then A is a locally paranilpotent G -module which is hypercyclic of height strictly larger than ω . However, it is straightforward to show that any locally

paranilpotent module is at least hypercyclic, and this is enough to prove that (locally finite)-by-(locally supersoluble) groups form a Hall class.

THEOREM 3.11. *The class of all (locally finite)-by-(locally supersoluble) groups is a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N' is (locally finite)-by-(locally supersoluble). Consider any finitely generated subgroup E of G , so in particular EN/N is (locally finite)-by-supersoluble. Since N/N' is \mathcal{T}_G -by-(locally paranilpotent) as an E -module, it is also \mathcal{T}_G -by-hypercyclic. The nilpotency of N and Corollary 3.6 yield that N (and so also EN) has an EN -invariant series of finite length whose factors are either locally finite, or abelian and hypercyclic as EN -modules. It follows now from Lemma 3.7 that EN is (locally finite)-by-hypercyclic and so E is (locally finite)-by-supersoluble. Therefore, G is (locally finite)-by-(locally supersoluble) by Lemma 3.10. \square

A similar proof to that of Lemma 2.25 shows that the class of locally (finite-by-supersoluble) groups is the intersection of the class of (locally finite)-by-(locally supersoluble) groups with the class of locally (nilpotent-by-finite) groups. Thus, locally (finite-by-supersoluble) groups form a Hall class by Theorem 3.11; a different proof of this result is given in [7].

4. FC-nilpotency conditions

Let G be a group. The *FC-centre* of G is the characteristic subgroup $FC_1(G)$ consisting of all elements of G with only finitely many conjugates, or equivalently of those elements $g \in G$ for which the centralizer $C_G(g)$ has finite index in G . The *upper FC-central series* of G is the ascending characteristic series $\{FC_\alpha(G)\}_\alpha$ recursively defined by setting $FC_0(G) = \{1\}$,

$$FC_{\alpha+1}(G)/FC_\alpha(G) = FC_1(G/FC_\alpha(G))$$

for each ordinal α and

$$FC_\lambda(G) = \bigcup_{\beta < \lambda} FC_\beta(G)$$

if λ is a limit ordinal. The last term of the upper *FC-central series* of G is called the *FC-hypercentre* of G , and G is said to be *FC-hypercentral* if it coincides with its *FC-hypercentre*. Moreover, G is called *FC-nilpotent* if $FC_k(G) = G$ for some nonnegative integer k . Of course, nilpotent-by-finite groups are *FC-nilpotent* and it is known that any finitely generated *FC-nilpotent* group is nilpotent-by-finite (see [13]). It follows that every *FC-hypercentral* group is locally *FC-nilpotent* and that the class of locally *FC-nilpotent* groups coincides with that of locally (nilpotent-by-finite) groups.

In this section, we prove that $(\mathbf{L}\mathfrak{F})\mathfrak{H}$ is a Hall class, when \mathfrak{H} is chosen to be one of the classes of *FC-nilpotent*, *FC-hypercentral* or locally *FC-nilpotent* groups.

Let G be a group. If A is a G -module and a is any element of A , we denote by G_a the stabilizer of a in G , that is, the subgroup formed by all elements $g \in G$ such that $ag = a$. It is easy to see that the set $FC_1^G(A)$, consisting of all elements a of A such that the index $|G : G_a|$ is finite, is a G -submodule of A . We can now define recursively the upper FC -series $\{FC_\alpha^G(A)\}$ of A by

$$FC_0^G(A) = \{0\}, \quad FC_{\alpha+1}^G(A)/FC_\alpha^G(A) = FC_1^G(A)/FC_\alpha^G(A)$$

for each ordinal α and

$$FC_\lambda^G(A) = \bigcup_{\beta < \lambda} FC_\beta^G(A)$$

if λ is a limit ordinal. The last term of this series is called the FC^G -hypercentre of A , and A is called hyper- FC if coincides with its FC^G -hypercentre. Moreover, the G -module A is said to be poly- FC if $A = FC_k^G(A)$ for some nonnegative integer k .

Since G -modules that are poly- FC and G -modules that are hyper- FC form tensorial classes (see for instance [1]), an application of Lemma 2.1 yields the following result.

COROLLARY 4.1. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by-(poly- FC) and the class of all G -modules that are \mathcal{T}_G -by-(hyper- FC) are tensorial.*

To prove that (locally finite)-by-(FC -nilpotent) groups form a Hall class, it is not possible to emulate the proof of Theorem 2.5 since there is no analogue of Corollary 2.4 in this case. In fact, we now construct a group that is not (locally finite)-by-(FC -hypercentral), although it is locally finite over its FC -centre.

Let H and K be groups and let N be a normal subgroup of K . Then $H \wr_N K$ denotes the split extension of the direct product B of copies H_{Nk} of H , where Nk runs over K/N , and K permutes the copies H_{Nk} in the obvious way. In particular, N centralizes the base group of $H \wr_N K$, which is B .

Let $C_1, C_2, \dots, C_n, \dots$ be a countably infinite sequence of groups of order 2, and set by induction $W_{n+1} = C_{n+1} \wr W_n$ for each nonnegative integer n , where $W_0 = \{1\}$; moreover, let W be the direct limit of the groups W_n with the obvious inclusions. Consider now an infinite dihedral group $X = \langle x, y \rangle$, where x has infinite order and $y^2 = 1$, and set $G_1 = X$ and $A_1 = \langle x \rangle$. Suppose that we have constructed, for some $n \geq 1$, a group G_n containing a free abelian normal subgroup A_n of finite rank such that $G_n/A_n \simeq W_n$. Put $G_{n+1} = X \wr_{A_n} G_n$ and let $A_{n+1} = A_n \times B_{n+1}$, where B_{n+1} denotes the normal closure of $\langle x \rangle$ in G_{n+1} . Then A_{n+1} is a free abelian normal subgroup of finite rank of G_{n+1} and $G_{n+1}/A_{n+1} \simeq W_{n+1}$.

Let $G = \bigcup_n G_n$ be the direct limit of the G_n , so $A = \bigcup_n A_n$ is a free abelian normal subgroup of G and $G/A \simeq W$ is locally finite. For each positive integer k , the subgroup A_k centralizes the base group of G_h for all $h > k$, and so every element of A_k has at most $|W_k|$ conjugates in G_h . It follows that A is contained in the FC -centre of G . If g is any element of $G \setminus A$, there is a positive integer m such that $g \in G_m \setminus A_m$ and hence there exists a in A_{m+1} with $[a, g] \neq 1$. Thus, $C_G(A) = A$ and so G has no nontrivial periodic normal subgroups. Since $G/A \simeq W$ has no nontrivial finite normal subgroups

(see for instance [17, Part 2, Theorem 6.23]), A coincides with the FC -hypercentre of G . Consequently, G is locally finite over the FC -centre, but it is not (locally finite)-by- $(FC$ -hypercentral).

In contrast to the above example, we can at least prove that a result corresponding to Corollary 2.4 holds in the case of modules. To this aim, we need the following result that should be seen in relation with a well-known theorem of Mal'cev on the upper central series of torsion-free groups (see for instance [17, Part 1, Theorem 2.25]).

LEMMA 4.2. *Let G be a group and let A be a G -module that is torsion-free as an abelian group. Then $A/FC_\alpha^G(A)$ is torsion-free as an abelian group for each ordinal α .*

PROOF. It is clearly enough to prove the statement when $\alpha = 1$. Let a be any element of A such that $b = ka \in FC_1^G(A)$ for some positive integer k . If g belongs to the stabilizer G_b , we have $k(a(g-1)) = b(g-1) = 0$ and hence $a(g-1) = 0$ because A is torsion-free as an abelian group. Thus, $G_a = G_b$ has finite index in G , so $a \in FC_1^G(A)$ and hence $A/FC_1^G(A)$ is torsion-free as an abelian group. \square

COROLLARY 4.3. *Let G be a group and let A be a G -module that is torsion-free as an abelian group.*

- (a) *If A has a finite series of G -submodules each of whose factors either belongs to \mathcal{T}_G or is poly- \mathcal{FC} , then A is poly- \mathcal{FC} .*
- (b) *If A has a finite series of G -submodules each of whose factors either belongs to \mathcal{T}_G or is hyper- \mathcal{FC} , then A is hyper- \mathcal{FC} .*

PROOF. We only prove part (a) because the same argument works also for part (b). Let

$$\{0\} = A_0 < A_1 < \cdots < A_t = A$$

be a finite series of G -submodules of A such that, for each $i = 0, 1, \dots, t-1$, the G -module A_{i+1}/A_i either belongs to \mathcal{T}_G or is poly- \mathcal{FC} . Clearly, A_1 is poly- \mathcal{FC} and if T/A_1 is the subgroup consisting of all elements of finite order of A/A_1 , it follows from Lemma 4.2 that T is also poly- \mathcal{FC} . Moreover, the G -module A/T is torsion-free as an abelian group and so it is poly- \mathcal{FC} by induction on t . Therefore, A is poly- \mathcal{FC} and the statement is proved. \square

THEOREM 4.4. *The class of all (locally finite)-by- $(FC$ -nilpotent) groups is a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N is (locally finite)-by- $(FC$ -nilpotent). To prove that G itself is (locally finite)-by- $(FC$ -nilpotent), we may suppose that G has no nontrivial locally finite normal subgroups. Then it follows from Theorem 2.5 and Lemma 2.19 that G contains a nilpotent normal subgroup M such that G/M is (locally finite)-by- $(FC$ -nilpotent) and G/M is FC -nilpotent, and of course M is torsion-free.

We claim that M is contained in $FC_k(G)$ for some nonnegative integer k . Consider $A = \zeta_1(M)$ as a G -module by conjugation. Since M/A is torsion-free (see for instance

[17, Part 1, Theorem 2.25]), induction on the nilpotency class of M yields that M/A is contained in some term with finite ordinal type of the upper FC -central series of G/A . Moreover, M/M' is \mathcal{T}_G -by-(poly- \mathcal{FC}) as a G -module, and so it follows from Corollary 4.1 that A has a finite series of G -submodules each of whose factors is either in \mathcal{T}_G or poly- \mathcal{FC} . Thus, A is poly- \mathcal{FC} by Corollary 4.3(a), so that $A \leq FC_h(G)$ for some integer $h > 0$ and hence the claim is proved.

Since $M \leq FC_k(G)$ and G/M is FC -nilpotent, it follows that G is FC -nilpotent and the proof is complete. □

The same argument, essentially replacing poly- \mathcal{FC} by hyper- \mathcal{FC} and part (a) of Corollary 4.3 by part (b), allows us to prove the following.

THEOREM 4.5. *The class of all (locally finite)-by-(FC -hypercentral) groups is a Hall class.*

The final result of this section shows that $(\mathbf{L}\mathfrak{F})(\mathbf{L}(\mathfrak{R}\mathfrak{F}))$ is a Hall class. Notice that obviously, if G is a group and A is a G -module that is locally poly- \mathcal{FC} , then A is also hyper- \mathcal{FC} (although, as in the case of locally paranilpotent modules, it is not difficult to see that hyper- \mathcal{FC} modules are not in general locally poly- \mathcal{FC}).

THEOREM 4.6. *The class of all (locally finite)-by-(locally FC -nilpotent) groups is a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N' is (locally finite)-by-(locally FC -nilpotent). To prove that G itself is (locally finite)-by-(locally FC -nilpotent), we may obviously suppose that G has no nontrivial locally finite normal subgroups. Thus, Theorem 2.5 and Lemma 2.19 yield that G contains a nilpotent normal subgroup M such that G/M' is (locally finite)-by-(locally FC -nilpotent) and G/M is locally FC -nilpotent; of course, M is torsion-free.

Consider now any finitely generated subgroup E of G , so in particular the group EM/M is FC -nilpotent. Since M/M' is \mathcal{T}_E -by-(locally poly- \mathcal{FC}) as an E -module, it is also \mathcal{T}_E -by-(hyper- \mathcal{FC}). The nilpotency of M and Corollary 4.1 yield that M has an EM -invariant series of finite length whose factors are either locally finite, or torsion-free abelian and hyper- \mathcal{FC} as EM -modules.

It follows now from Corollary 4.3 that each factor of the upper central series of M is hyper- \mathcal{FC} as an EM -module. Thus, M is contained in the FC -hypercentre of EM and so EM is FC -hypercentral. In particular, E is nilpotent-by-finite and hence G is locally FC -nilpotent. □

5. Solubility and rank conditions

In this short section, we deal with group classes of the form $(\mathbf{L}\mathfrak{F})\mathfrak{Y}$, where \mathfrak{Y} is either a class of generalized soluble groups or a class of groups satisfying suitable rank conditions. To avoid the repetition of the same argument, we prove the following general lemma, which is of independent interest.

LEMMA 5.1. *Let \mathfrak{X} and \mathfrak{Y} be group classes such that $\mathfrak{X} = \mathbf{S}_n\mathfrak{X}$ and $\mathfrak{N}\mathfrak{Y} = \mathfrak{Y}$. If $\mathfrak{X}\mathfrak{N}$ is a Hall class, then $\mathfrak{X}\mathfrak{Y}$ is also a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N' belongs to $\mathfrak{X}\mathfrak{Y}$, so G has a normal subgroup M with $N' \leq M$, $M/N' \in \mathfrak{X}$ and $G/M \in \mathfrak{Y}$. Thus, $MN/N' \in \mathfrak{X}\mathfrak{N}$ and hence MN belongs to the Hall class $\mathfrak{X}\mathfrak{N}$. Let L be a normal \mathfrak{X} -subgroup of MN such that MN/L is nilpotent and let c be a positive integer with $\gamma_c(MN) \leq L$. Then $X = \gamma_c(MN) \cap M$ is a normal subgroup of G that belongs to $\mathfrak{X} = \mathbf{S}_n\mathfrak{X}$. Moreover, G/X belongs to $\mathfrak{N}\mathfrak{Y} = \mathfrak{Y}$ and hence G is in $\mathfrak{X}\mathfrak{Y}$. Therefore, $\mathfrak{X}\mathfrak{Y}$ is a Hall class. \square

By Theorem 2.5, the above lemma applies in particular to $\mathfrak{X} = \mathbf{L}\mathfrak{F}$ when either $\mathfrak{Y} = \mathfrak{S}$ or $\mathfrak{Y} = \mathbf{L}\mathfrak{S}$, but also if \mathfrak{Y} is the class $\mathbf{P}_n\mathfrak{A}$ consisting of all hyperabelian groups.

THEOREM 5.2. *The group classes $(\mathbf{L}\mathfrak{F})\mathfrak{S}$, $(\mathbf{L}\mathfrak{F})(\mathbf{P}_n\mathfrak{A})$ and $(\mathbf{L}\mathfrak{F})(\mathbf{L}\mathfrak{S})$ are Hall classes.*

The fact that $(\mathbf{L}\mathfrak{F})\mathfrak{S}$ is a Hall class can be stated in a quantitative way, by using Theorem 2.6 in the proof of Lemma 5.1.

As a further consequence of Theorem 2.5, we have the following easy result concerning the class \mathfrak{N}^k of soluble groups admitting a finite normal series with nilpotent factors of length at most k .

THEOREM 5.3. *The class $(\mathbf{L}\mathfrak{F})\mathfrak{N}^k$ is a Hall class for each positive integer k .*

PROOF. By Theorem 2.5, the statement is true if $k = 1$. Suppose $k > 1$ and let G be a group containing a nilpotent normal subgroup N such that $G/N' \in (\mathbf{L}\mathfrak{F})\mathfrak{N}^k$. Then G/N' has a normal subgroup $M/N' \in (\mathbf{L}\mathfrak{F})\mathfrak{N}^{k-1}$ such that G/M is nilpotent. Since $k > 1$, the group MN/N' belongs to the class $(\mathbf{L}\mathfrak{F})\mathfrak{N}^{k-1}$ by Fitting's theorem; whence MN lies in $(\mathbf{L}\mathfrak{F})\mathfrak{N}^{k-1}$ by induction on k . Therefore, $G \in (\mathbf{L}\mathfrak{F})\mathfrak{N}^k$ and the statement is proved. \square

In the second half of this section, we deal with group classes defined by the imposition of a rank condition. We start by recalling the necessary definitions.

Recall that a group G is said to have *finite (Prüfer) rank* if there is a positive integer r such that every finitely generated subgroup of G can be generated by r elements. Of course, the tensor product of two abelian groups of finite rank has finite rank (see for instance [22, Lemma 10]) and hence the class of groups of finite rank is easily seen to be a Hall class.

Let A be an abelian group. The *0-rank* $r_0(A)$ of A is defined as the rank of the torsion-free group A/T , where T is the periodic part of A ; moreover, for each prime p , the *p -rank* $r_p(A)$ of A is the rank of the p -component of A . Of course, A has finite rank if and only if

$$\sup\{r_0(A), r_p(A) \mid p \text{ a prime}\} < \infty.$$

We say that A has *finite total rank* if

$$r_0(A) + \sum_p r_p(A) < \infty.$$

A group G is said to have *finite abelian section rank* if every abelian section of G has finite 0-rank and finite p -rank for all primes p . Moreover, G has *finite abelian total rank* if it admits a series of finite length each of whose infinite factors is an abelian group of finite total rank; of course, every group of finite abelian total rank is soluble-by-finite and has finite rank, while any torsion-free nilpotent group of finite rank has finite abelian total rank, because all factors of its upper central series are torsion-free. Notice that the class of groups with finite abelian total rank is not quotient-closed, since \mathbb{Q}/\mathbb{Z} is the direct product of infinitely many Prüfer groups; however, if G is a group with finite abelian total rank and N is a periodic normal subgroup of G , it is easy to see that G/N also has finite abelian total rank.

If G is a group, in the following, we denote by \mathcal{R}_G the class of all G -modules that have finite rank as abelian groups. Our next result is an application of Lemma 2.1.

COROLLARY 5.4. *If G is a group, the class of all G -modules that are \mathcal{T}_G -by- \mathcal{R}_G is tensorial.*

The following result is well known (and even trivial in the abelian case), so for completeness, we only sketch its proof.

LEMMA 5.5. *Let G be a group admitting a normal series of finite length whose factors are either locally finite or abelian of finite rank. Then G contains a locally finite normal subgroup T such that G/T has finite rank.*

PROOF. By induction, it is enough to prove the statement when G contains a torsion-free abelian normal subgroup A of finite rank such that G/A is locally finite. Then $C = C_G(A)$ has finite index in G because periodic linear groups over the field of rational numbers are finite (see for instance [20, Theorem 9.33]). Moreover, C' is locally finite by Schur's theorem and the abelian group C/C' is periodic-by-(finite rank), which completes the proof. \square

THEOREM 5.6. *The class of all (locally finite)-by-(finite rank) groups is a Hall class.*

PROOF. The proof runs along the same lines as that of Theorem 2.17, just replacing Engel modules by modules whose underlying group has finite rank, and using Corollary 5.4 and Lemma 5.5. \square

We point out that since periodic subgroups of $\mathrm{GL}(n, \mathbb{Q})$ are finite and their order is bounded in terms of n , a slight modification of Lemma 5.5 allows us to obtain a quantitative version of Theorem 5.6.

THEOREM 5.7. *The class of all (locally finite)-by-(finite abelian total rank) groups is a Hall class.*

PROOF. Let G be a group and let N be a nilpotent normal subgroup of G such that G/N contains a locally finite normal subgroup L/N' with G/L of finite abelian total rank. We may obviously assume that G has no nontrivial locally finite normal subgroups, so in particular G has finite rank by Theorem 5.6. Let c be the nilpotency class

of the torsion-free nilpotent group N' . For each nonnegative integer $i < c$, we have that $\zeta_{i+1}(N')/\zeta_i(N')$ is torsion-free abelian of finite rank and hence $L/C_L(\zeta_{i+1}(N')/\zeta_i(N'))$ is finite (see [20, Theorem 9.33]). Thus, the normal subgroup

$$C = \bigcap_{i=0}^{c-1} C_L(\zeta_{i+1}(N')/\zeta_i(N'))$$

has finite index in L , so G/C has finite abelian total rank. Since $N' \leq \zeta_c(C)$, the group $C/\zeta_c(C)$ is locally finite and hence C is torsion-free nilpotent (see [17, Part 1, Corollary 2 to Theorem 4.21]). It follows that C has finite abelian total rank and hence G also has finite abelian total rank. The statement is proved. \square

THEOREM 5.8. *The class of all (locally finite)-by-(finite abelian section rank) groups is a Hall class.*

PROOF. Let G be a group containing a nilpotent normal subgroup N such that G/N' is (locally finite)-by-(finite abelian section rank). We may suppose by Lemma 2.19 that G/N has finite abelian section rank. Of course, it can also be assumed that N is torsion-free. Since N/N' is (locally finite)-by-(finite abelian total rank), it follows from Theorem 5.7 that N has finite abelian total rank. Therefore, G has finite abelian section rank and the statement is proved. \square

6. Obstructions of finite exponent

In the previous sections, we proved that $(\mathbf{L}\mathfrak{F})\mathfrak{X}$ is a Hall class for several natural choices of the group class \mathfrak{X} . It is therefore natural to ask what happens when $\mathbf{L}\mathfrak{F}$ is replaced by its subclass \mathfrak{B} consisting of all locally finite groups of finite exponent. In dealing with this problem, the role played by Schur's theorem in the case of locally finite obstructions is taken by the following result of Mann [12].

LEMMA 6.1. *Let G be a group such that $G/\zeta_1(G)$ is locally finite of finite exponent e . Then G' has finite exponent bounded by a function of e .*

If k is any positive integer and A is an abelian group of exponent k , the tensor product $A \otimes_{\mathbb{Z}} B$ has exponent dividing k for every abelian group B . Thus, the proof of Lemma 2.1 can be used to prove the following result, where $\mathcal{B}_G^{(k)}$ denotes the class of all G -modules whose underlying group has finite exponent dividing k .

LEMMA 6.2. *Let G be a group and let \mathcal{M} be a quotient-closed, tensorial class of G -modules. Then for each positive integer k , the class of all G -modules that are $\mathcal{B}_G^{(k)}$ -by- \mathcal{M} is tensorial.*

If G is a group and \mathcal{M} is a class of G -modules, we denote by \mathbf{PM} the class of all G -modules A admitting a finite series of G -submodules

$$\{0\} = A_0 < A_1 < \dots < A_t = A$$

such that A_{i+1}/A_i belongs to \mathcal{M} for each $i = 0, 1, \dots, t - 1$. The G -modules in the class \mathcal{PM} are called *poly-M*.

LEMMA 6.3. *Let G be a group and let \mathcal{M} be a class of G -modules that is closed with respect to forming G -submodules. If A is a G -module and B is a G -submodule of A such that $B \in \mathcal{M}$ and $A/B \in \mathcal{B}_G^{(k)}$ for some positive integer k , then there exists a G -submodule C of A with $C \in \mathcal{B}_G^{(k)}$ and $A/C \in \mathcal{M}$.*

PROOF. Clearly, $kA \leq B$ and so $kA \in \mathcal{M}$. Since the map $\varphi : a \mapsto ka$ is a G -homomorphism of A onto kA , it follows that the kernel C of φ is a G -submodule of A whose exponent as an abelian group divides k and the G -module A/C is in \mathcal{M} . □

The above lemma and a simple induction argument prove the following result.

COROLLARY 6.4. *Let G be a group and let \mathcal{M} be a class of G -modules that is closed with respect to forming G -submodules. If k is a positive integer and A is a G -module admitting a series of G -submodules of finite length t each of whose factors belongs to $\mathcal{B}_G^{(k)} \cup \mathcal{M}$, then there exists a G -submodule C of A such that $C \in \mathcal{B}_G^{(k)}$ and A/C is *poly-M*.*

A crucial point in our considerations is to prove that \mathfrak{BN} is a Hall class.

LEMMA 6.5. *Let G be a group and let N be a normal subgroup of G . If N has a finite G -invariant series each of whose factors is either in \mathfrak{B} or central in G , then there is a positive integer k such that $[N, {}_k G]$ belongs to \mathfrak{B} .*

PROOF. Let

$$\{1\} = N_0 < N_1 < \dots < N_t = N$$

be a finite G -invariant series of minimal length t such that for each nonnegative integer $i < t$, N_{i+1}/N_i either belongs to \mathfrak{B} or is central in G . By induction, there is a positive integer h such that $M/N_1 = [N, {}_h G]N_1/N_1 \in \mathfrak{B}$, and without loss of generality we may suppose that $N_1 \leq \zeta_1(G)$. Then the group $M/\zeta_1(M)$ is locally finite of finite exponent and so by Lemma 6.1, M' also lies in \mathfrak{B} . Application of Lemma 6.3 to the G -module M/M' when \mathcal{M} is the class of all trivial G -modules yields that M contains a G -invariant subgroup $L \geq M'$ of finite exponent such that $M/L \leq \zeta_1(G/L)$. Thus, $[N, {}_{h+1} G] \leq [M, G] \leq L$ has finite exponent and the statement is proved. □

COROLLARY 6.6. *A group G is \mathfrak{B} -by-nilpotent if and only if it has a normal series of finite length each of whose factors either belongs to \mathfrak{B} or is central in G .*

THEOREM 6.7. *The class \mathfrak{BN} is a Hall class.*

PROOF. It is enough to reproduce the proof of Theorem 2.5, replacing Lemma 2.1 by Lemma 6.2 and Corollary 2.4 by Corollary 6.6. □

COROLLARY 6.8. *The group classes $\mathfrak{B}\mathfrak{E}$, $\mathfrak{B}(\mathfrak{P}_n\mathfrak{N})$ and $\mathfrak{B}(\mathbf{L}\mathfrak{E})$ are Hall classes.*

PROOF. Since $\mathfrak{B}\mathfrak{N}$ is a Hall class by Theorem 6.7, the statement follows directly from Lemma 5.1 by choosing $\mathfrak{X} = \mathfrak{B}$ and \mathfrak{Y} as one of the classes $\mathfrak{S}, \mathfrak{P}_n\mathfrak{N}, \mathfrak{L}\mathfrak{S}$. \square

COROLLARY 6.9. *The class $\mathfrak{B}\mathfrak{N}^k$ is a Hall class for each positive integer k .*

PROOF. The case $k = 1$ is just Theorem 6.7. Suppose $k > 1$ and let G be a group containing a nilpotent normal subgroup N such that $G/N' \in \mathfrak{B}\mathfrak{N}^k$. Then there is a normal subgroup M/N' of G/N' such that $G/M \in \mathfrak{N}$ and $M/N' \in \mathfrak{B}\mathfrak{N}^{k-1}$. Since $k > 1$, Fitting's theorem yields that MN/N' belongs to $\mathfrak{B}\mathfrak{N}^{k-1}$, so MN lies in $\mathfrak{B}\mathfrak{N}^{k-1}$ by induction. It follows that $G \in \mathfrak{B}\mathfrak{N}^k$ and so the statement is proved. \square

Our next theorem shows that the class \mathfrak{B} behaves better than \mathfrak{F} and $\mathfrak{L}\mathfrak{F}$, at least in our context. If e is any positive integer, we denote by \mathfrak{B}_e the class of all groups of finite exponent dividing e , so $\mathfrak{B} = \bigcup_e \mathfrak{B}_e$.

THEOREM 6.10. *The class $\mathfrak{B}\mathfrak{Y}$ is a Hall class, where \mathfrak{Y} is any of the following classes of groups:*

- (a) *the class of finitely generated nilpotent groups;*
- (b) *the class of hypercentral groups;*
- (c) *the class of locally nilpotent groups;*
- (d) *the class of Engel groups;*
- (e) *the class of supersoluble groups;*
- (f) *the class of paranilpotent groups;*
- (g) *the class of hypercyclic groups;*
- (h) *the class of locally supersoluble groups;*
- (i) *the class of FC-nilpotent groups;*
- (j) *the class of FC-hypercentral groups;*
- (k) *the class of locally FC-nilpotent groups.*

PROOF. Let \mathfrak{Y} be any of the classes considered in the statement and let N be a nilpotent normal subgroup of a group G such that G/N' belongs to $\mathfrak{B}\mathfrak{Y}$. It follows from Theorem 6.7 and Lemma 2.19 that G contains a normal subgroup M such that G/M' is in $\mathfrak{B}\mathfrak{Y}$, G/M is in \mathfrak{Y} and $\gamma_k(M)$ has finite exponent for some positive integer k . To prove that G itself is in $\mathfrak{B}\mathfrak{Y}$, we may factor out $\gamma_k(M)$ and so assume that M is nilpotent.

Put $A = \zeta_1(M)$, so $G/A \in \mathfrak{B}\mathfrak{Y}$ by induction on the nilpotency class of M . If K/A is a normal \mathfrak{B} -subgroup of G/A such that $G/K \in \mathfrak{Y}$, we have that G/B is also a \mathfrak{Y} -group, where $B = M \cap K$. Since $A \leq \zeta_1(B)$, it follows from Lemma 6.1 that B' belongs to \mathfrak{B} , so the replacement of G by G/B' allows us to assume that B is abelian.

We complete the proof only for parts (b), (c) and (h) since all other statements can be proved in similar ways.

Let \mathfrak{Y} be the class of hypercentral groups. Since M/M' is \mathcal{B}_G -by-hypercentrivial as a G -module, it follows from Lemma 6.2 that M has a finite G -invariant series each of whose factors is either in \mathfrak{B} or hypercentrally embedded in G . Thus, the G -module B has a finite series of G -submodules each of whose factors either belongs to \mathcal{B}_G or is hypertrivial and hence it contains a G -submodule C such that $C \in \mathcal{B}_G$ and B/C is

hypertrivial by Corollary 6.4. Of course, C is a normal subgroup of finite exponent of G and G/C is hypercentral. This completes the proof of part (b).

Suppose now that $\mathfrak{Y} = \mathbf{L}\mathfrak{N}$ and let e be a positive integer such that G/M' belongs to $\mathfrak{B}_e(\mathbf{L}\mathfrak{N})$. Consider any finitely generated subgroup E of G . Since M/M' is $\mathcal{B}_E^{(e)}$ -by-(locally polytrivial) as an E -module, it follows from Lemma 6.2 that the lower central factor $\gamma_i(M)/\gamma_{i+1}(M)$ likewise is a $\mathcal{B}_E^{(e)}$ -by-(locally polytrivial) E -module for each positive integer $i \leq c$, where c is the nilpotency class of M . Thus, the E -module B has a series of E -submodules of length at most c each of whose factors is $\mathcal{B}_E^{(e)}$ -by-(locally polytrivial). Since locally polytrivial E -modules are hypertrivial by Lemma 2.23, an application of Corollary 6.4 yields that B contains an E -invariant subgroup C such that B/C is hypercentrally embedded in EB/C and C has finite exponent dividing $h = e^{2c}$. Of course, EB/C is locally nilpotent and we also have $C \leq B[h] = \{b \in B \mid b^h = 1\}$, so that $EB[h]/B[h]$ is nilpotent and $G/B[h]$ is locally nilpotent. Therefore, $G \in \mathfrak{B}(\mathbf{L}\mathfrak{N})$.

Finally, if \mathfrak{Y} is the class of locally supersoluble groups, the proof runs along the same lines as part (c), just replacing locally polytrivial G -modules and hypertrivial G -modules by locally paranilpotent G -modules and hypercyclic G -modules, respectively; note that a group is locally supersoluble if and only if it is locally hypercyclic. □

To complete our analysis of classes of the form $\mathfrak{B}\mathfrak{Y}$, where \mathfrak{Y} is a class of generalized nilpotent groups, we need some additional definitions.

Let G be a group and let A be a G -module. We say that A is a *Baer G -module* if A is a polytrivial $\langle g \rangle$ -module for each element g of G . Similarly, A is called a *Gruenberg G -module* if it is hypertrivial as a $\langle g \rangle$ -module for every $g \in G$. Finally, A is said to be a *Fitting G -module* if it is polytrivial as a $\langle g \rangle^G$ -module for each $g \in G$. Of course, the classes of Fitting, Baer and Gruenberg G -modules are closed with respect to forming submodules, quotient modules and extensions; our next lemma shows that these classes are also tensorial.

LEMMA 6.11. *If G is any group, the class of Fitting G -modules, the class of Baer G -modules and the class of Gruenberg G -modules are tensorial.*

PROOF. Let A and B be Fitting (respectively Baer or Gruenberg) G -modules, and consider an arbitrary element g of G . Since A and B are polytrivial $\langle g \rangle^G$ -modules (respectively polytrivial $\langle g \rangle$ -modules or hypertrivial $\langle g \rangle$ -modules), also the tensor product $A \otimes_{\mathbb{Z}} B$ is a polytrivial $\langle g \rangle^G$ -module (respectively polytrivial $\langle g \rangle$ -module or hypertrivial $\langle g \rangle$ -module). Therefore, $A \otimes_{\mathbb{Z}} B$ is a Fitting (respectively Baer or Gruenberg) G -module and the statement is proved. □

The following results add some further Hall classes to the list given in Theorem 6.10.

THEOREM 6.12. *The class $\mathfrak{B}\mathfrak{Y}$ is a Hall class, where \mathfrak{Y} is one of the following classes of groups:*

- (l) *the class of Fitting groups;*
- (m) *the class of Baer groups;*
- (n) *the class of Gruenberg groups.*

PROOF. We only prove the statement for Fitting groups, since the other cases can be proved in a similar fashion. Let G be a group containing a nilpotent normal subgroup N such that G/N' is \mathfrak{B} -by-Fitting. As in the first part of the proof of Theorem 6.10, we may reduce to the case in which N contains an abelian G -invariant subgroup B such that G/B is a Fitting group. Since N/N' is \mathcal{B}_G -by-Fitting as a G -module, it follows from Lemma 6.2 that N has a finite G -invariant series each of whose factors either is in \mathfrak{B} or is an abelian group that is a Fitting G -module by conjugation. Thus, the G -module B has a finite series of G -submodules each of whose factors either belongs to \mathcal{B}_G or is a Fitting G -module. It follows from Corollary 6.4 that B contains a G -submodule C such that $C \in \mathcal{B}_G$ and B/C is a Fitting G -module.

If g is any element of G , we have that B/C is a polytrivial $\langle g \rangle^G$ -module, while $\langle g \rangle^G B/B$ is nilpotent because G/B is a Fitting group; thus, $\langle g \rangle^G C/C$ is nilpotent. It follows that G/C is a Fitting group and hence G is \mathfrak{B} -by-Fitting. □

THEOREM 6.13. *The class $\mathfrak{B}\mathfrak{V}$ is a Hall class, where \mathfrak{V} is one of the following classes of groups:*

- (o) *the class of groups of finite rank;*
- (p) *the class of groups with finite abelian section rank.*

PROOF. The proof is similar to that of part (b) of Theorem 6.10. □

Our results seem to show that obstructions of finite exponent can be treated more easily than locally finite ones. However, this is not always true, as proved by the class of groups with finite abelian total rank. In fact, let $T = \text{Tr}(2, \mathbb{Q})$ be the group of all lower triangular matrices of degree 2 over the rationals and consider its central subgroup Z generated by the matrix

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Then $G = T/Z$ is a nilpotent group such that $G/G' \simeq \mathbb{Q} \times \mathbb{Q}$ and G' is the direct product of infinitely many Prüfer groups. It follows that \mathfrak{B} -by-(finite abelian total rank) groups do not form a Hall class. This anomaly depends essentially on the fact that the class of groups with finite abelian total rank is not quotient-closed.

The results of this section fail to be true if the class \mathfrak{B} is replaced by the class $\overline{\mathfrak{B}}$ consisting of all groups of finite exponent. Actually, it was proved by Endimioni and Traustason [9] that there exists a group G having a nilpotent normal subgroup N such that $G/N' \in \overline{\mathfrak{B}\mathfrak{A}}$, but G is not even in $\overline{\mathfrak{B}(\mathbb{L}\mathfrak{S})}$; in particular, $\overline{\mathfrak{B}\mathfrak{A}}$ is not a Hall class. Moreover, the replacement in their example of the Adjan group by the group constructed in [14, Theorem 31.4] shows that in Theorem 6.13, the class \mathfrak{B} cannot be replaced by the class $\overline{\mathfrak{B}}$.

As we did in the case of a locally finite obstruction at the bottom, one can slightly modify the proofs to obtain quantitative versions of Theorems 6.7, 6.13(o) and Corollary 6.8 (for the class $\mathfrak{B}\mathfrak{E}$). For instance, we have the following result.

THEOREM 6.14. *Let G be a group and let N be a normal subgroup of G . If N is nilpotent of class c and G/N' is (of finite exponent e)-by-(nilpotent of class d), then there exist positive integers h and k , depending only on c , d and e , such that G is (of exponent at most h)-by-(nilpotent of class at most k).*

It is reasonable to ask whether analogues of Corollary 6.6 can be proved for other group classes of the form $\mathfrak{B}\mathfrak{Y}$, since results of this type would provide alternative proofs for some of the theorems in this section. Our final result shows that, for instance, this is the case when \mathfrak{Y} is the class of paranilpotent groups.

LEMMA 6.15. *Let G be a group and let \mathcal{M} be a class of G -modules that is closed with respect to forming G -submodules and G -homomorphic images, contains all trivial G -modules and satisfies the condition $M(g-1) = \{0\}$ for each $M \in \mathcal{M}$ and $g \in G'$. If N is a normal subgroup of G admitting a finite G -invariant series whose factors are either in \mathfrak{B} or G -modules in \mathcal{M} , then there exists a G -invariant subgroup B of N of finite exponent such that N/B has a finite G -invariant series whose factors are G -modules in \mathcal{M} .*

PROOF. Let

$$\{1\} = N_0 < N_1 < \cdots < N_t = N$$

be a finite G -invariant series of minimal length t such that each N_{i+1}/N_i is either in \mathfrak{B} or a G -module in \mathcal{M} . By induction, there exists a G -invariant subgroup $M \geq N_1$ of N such that M/N_1 is in \mathfrak{B} and N/M has a finite G -invariant series whose factors are G -modules in \mathcal{M} . Thus, we may suppose that N_1 is a G -module in \mathcal{M} , and in particular that G' centralizes N_1 . Consider the normal subgroup $X = (M \cap G')N_1$ of G . Clearly, $N_1 \leq \zeta_1(X)$, so $X/\zeta_1(X) \in \mathfrak{B}$ and hence also X' belongs to \mathfrak{B} by Lemma 6.1. Application of Lemma 6.3 to the G -module X/X' and to the class \mathcal{M} yields that X/X' contains a \mathfrak{B} -subgroup B/X' that is normal in G and such that X/B belongs to \mathcal{M} . Since M/X is a central factor of G , it follows that N/B has a finite G -invariant series whose factors are either in \mathfrak{B} or G -modules in \mathcal{M} . Of course, $B \in \mathfrak{B}$ and so the proof is complete. \square

COROLLARY 6.16. *A group G is \mathfrak{B} -by-paranilpotent if and only if it has a normal series of finite length whose factors are either in \mathfrak{B} or paracentral in G .*

References

- [1] B. Amberg, S. Franciosi and F. de Giovanni, 'Groups with an FC-nilpotent triple factorization', *Ric. Mat.* **36** (1987), 103–114.
- [2] C. Casolo, *Groups with all Subgroups Subnormal* (AGTA Lost Monographs, Aduvare, Rome, 2022).

- [3] M. De Falco, F. de Giovanni, C. Musella and Y. P. Sysak, 'On the upper central series of infinite groups', *Proc. Amer. Math. Soc.* **139** (2011), 385–389.
- [4] F. de Giovanni and M. Trombetti, 'Infinite minimal non-hypercyclic groups', *J. Algebra Appl.* **14** (2015), 1550143, 15 pages.
- [5] F. de Giovanni, M. Trombetti and B. A. F. Wehrfritz, 'Linear groups whose proper subgroups are close to being nilpotent', *Comm. Algebra* **49** (2021), 3020–3033.
- [6] F. de Giovanni, M. Trombetti and B. A. F. Wehrfritz, 'Subnormality in linear groups', *J. Pure Appl. Algebra* **227** (2023), 107185, 16 pages.
- [7] F. de Giovanni, M. Trombetti and B. A. F. Wehrfritz, 'Hall classes of groups', to appear.
- [8] F. de Giovanni, M. Trombetti and B. A. F. Wehrfritz, 'Hall classes in linear groups', to appear.
- [9] G. Endimioni and G. Traustason, 'On torsion-by-nilpotent groups', *J. Algebra* **241** (2001), 669–676.
- [10] S. Franciosi and F. de Giovanni, 'Groups with many supersoluble subgroups', *Ric. Mat.* **40** (1991), 321–333.
- [11] P. Hall, 'Some sufficient conditions for a group to be nilpotent', *Illinois J. Math.* **2** (1958), 787–801.
- [12] A. Mann, 'The exponents of central factor and commutator groups', *J. Group Theory* **10** (2007), 435–436.
- [13] D. H. McLain, 'Remarks on the upper central series of a group', *Proc. Glasg. Math. Assoc.* **3** (1956), 38–44.
- [14] A. Y. Ol'shanskii, *Geometry of Defining Relations in Groups* (Kluwer, Dordrecht, 1989).
- [15] B. I. Plotkin, 'Some properties of automorphisms of nilpotent groups', *Soviet Math. Dokl.* **2** (1961), 471–474.
- [16] D. J. S. Robinson, 'A property of the lower central series of a group', *Math. Z.* **107** (1968), 225–231.
- [17] D. J. S. Robinson, *Finiteness Conditions and Generalized Soluble Groups* (Springer, Berlin, 1972).
- [18] P. Shumyatsky, 'On extensions of groups of finite exponent', *Glasg. Math. J.* **45** (2003), 535–538.
- [19] B. A. F. Wehrfritz, 'Supersoluble and locally supersoluble linear groups', *J. Algebra* **17** (1971), 41–58.
- [20] B. A. F. Wehrfritz, *Infinite Linear Groups* (Springer, Berlin, 1973).
- [21] B. A. F. Wehrfritz, *Group and Ring Theoretic Properties of Polycyclic Groups* (Springer, London, 2009).
- [22] B. A. F. Wehrfritz, 'Rank–Engel conditions on linear groups', *Rend. Circ. Mat. Palermo (2)* **70** (2021), 45–56.

F. DE GIOVANNI, Dipartimento di Matematica e Appl.,
Università di Napoli Federico II, Napoli, Italy
e-mail: degiovan@unina.it

M. TROMBETTI, Dipartimento di Matematica e Appl.,
Università di Napoli Federico II, Napoli, Italy
e-mail: marco.trombetti@unina.it

B. A. F. WEHRFRITZ, School of Mathematical Sciences,
Queen Mary University of London, London, UK
e-mail: b.a.f.wehrfritz@qmul.ac.uk