

Effectiveness of the predatory mite *Neoseiulus cucumeris* on two-spotted spider mite and western flower thrips: A quantitative assessment - Erratum

Erratum

Cite this article: Dalir S, Hajiqaanbar H, Fathipour Y, Khanamani M (2025) Effectiveness of the predatory mite *Neoseiulus cucumeris* on two-spotted spider mite and western flower thrips: A quantitative assessment - Erratum. *Bulletin of Entomological Research* **115**, 404. <https://doi.org/10.1017/S0007485325000239>

Sajjad Dalir¹, Hamidreza Hajiqaanbar¹, Yaghoob Fathipour¹ and Mostafa Khanamani²

¹Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran and ²Department of Plant Protection, Faculty of Agriculture, University of Jiroft, Kerman, Iran

<https://doi.org/10.1017/S0007485325000033>, Published online by Cambridge University Press: 7 February 2025

When this article was originally published in the Bulletin of Entomological Research it omitted to include Yaghoob Fathipour as a corresponding author. This article has since been updated to include Dr. Fathipour as a corresponding author as originally intended by the authors.

The publisher apologises for this error.

References

Dalir S, Hajiqaanbar H, Fathipour Y, Khanamani M (2025) Effectiveness of the predatory mite *Neoseiulus cucumeris* on two-spotted spider mite and western flower thrips: A quantitative assessment. *Bulletin of Entomological Research* **115**(2), 184–193. doi:[10.1017/S0007485325000033](https://doi.org/10.1017/S0007485325000033)