

Organoids in parasitology: a game-changer for studying host–nematode interactions

Matias Gaston Perez¹ , Diana Coman², Joana Neves² and Collette Britton¹ 

¹School of Biodiversity, One Health and Veterinary Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK and ²Centre for Host–Microbiome Interactions, Faculty of Dentistry, Oral & Craniofacial Sciences, King's College London, London, UK

Review Article

Cite this article: Perez MG, Coman D, Neves J, Britton C (2025) Organoids in parasitology: a game-changer for studying host–nematode interactions. *Parasitology*, 1–12. <https://doi.org/10.1017/S0031182025100620>

Received: 12 June 2025

Revised: 4 July 2025

Accepted: 7 July 2025

Keywords:

gastrointestinal nematodes; host–parasite interactions; immune modulation; intestinal epithelium; *in vitro* models; organoids; stem cells

Corresponding author: Matias Gaston Perez;
Email: matias.perez@glasgow.ac.uk

Abstract

Gastrointestinal (GI) nematode infections represent a significant health burden globally, affecting both humans and livestock. Traditional *in vitro* models to study host–parasite interactions, such as immortalized cell lines, have limitations that hinder the full understanding of these complex relationships. Organoid technology has emerged as a promising alternative, offering a physiologically relevant platform to study host–nematode interactions *in vitro*. Organoids are three-dimensional structures comprising differentiated cell types that recapitulate features of the corresponding organ. Technological advances for growing, maintaining and manipulating organoids have increased their applications to model infections, inflammation and cancer. This review discusses recent work using GI organoids to advance understanding of nematode–host interactions and modulation of GI epithelial cells. Additionally, we review studies that co-cultured GI organoids with innate lymphoid cells to study epithelial–immune cell cross-talk in the context of nematode infection. By bridging the gap between reductionist cell culture systems and whole-organism studies, organoids offer a powerful platform for investigating complex host–nematode interactions, and for developing and screening novel therapeutics.

Introduction

Gastrointestinal (GI) nematode infections pose a major health burden, affecting over 1.5 billion people, particularly in tropical and subtropical regions with inadequate sanitation (World Health Organization 2024). These macroparasites have evolved intricate life cycles that enable them to successfully colonize the GI tract of humans and other mammals. Prominent human species such as *Ascaris lumbricoides*, *Trichuris trichiura* and *Necator americanus* contribute to malnutrition, stunted growth and cognitive impairments, especially in children (Serván et al. 2024). Similarly, in livestock, GI nematodes cause significant economic losses due to reduced weight gain, decreased milk production and increased mortality (Strydom et al. 2023; Vande Velde et al. 2018). For example, *Haemonchus contortus* and *Ostertagia ostertagi*, which infect sheep and cattle, respectively, result in billions of dollars in losses annually (Charlier et al. 2020). Additionally, in horses, nematodes such as *Strongylus vulgaris* and Cyathostominae cause severe colic and life-threatening complications (Corning 2009).

GI nematodes are specially adapted to evade host defences, migrate through tissues and establish a niche, often leading to chronic infections (maizels And Gause 2023). Given the increasing resistance of GI nematodes to anthelmintic drugs, understanding how these parasites establish infection and evade host immunity is important for developing more sustainable control strategies in both human and veterinary medicine (Charlier et al. 2022; Hotez and Herricks 2015; Nielsen 2022; Vegvari et al. 2021). The intestinal epithelium plays a central role in host–parasite interactions by detecting pathogens and orchestrating protective mechanisms, leading to increased mucus secretion and enhanced peristalsis – collectively known as the ‘weep and sweep’ response (Bąska and Norbury 2022). Several studies in recent years identified epithelial tuft cells, a type of chemosensory secretory cell, as initial ‘sensors’ of parasite infection in the small intestine (SI) (Gerbe et al. 2016; Howitt et al. 2016; von Moltke et al. 2016). Further understanding of GI epithelial cell responses to nematode infection, and their interaction with immune cells is important in gaining insight into the initiation of host immune responses required for effective parasite control.

Mucosal tissues of the GI tract initiate innate and adaptive type 2 immune responses to nematode infection (Artis and Grencis 2008). Alarmins interleukin (IL)-25 from GI tuft cells, thymic stromal lymphopoietin (TSLP) from epithelial cells, and IL-33 from epithelial, endothelial and inflammatory immune cells, activate Group 2 innate lymphoid cells (ILC2s) (Hardman et al. 2013). Once activated, ILC2s play a crucial role as an early source of type-2 cytokines, in particular IL-4 and IL-13 which act in a feed forward loop to increase the numbers of tuft

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

and mucous-secreting cells in the GI epithelium (Fallon *et al.* 2006; Molofsky and Locksley 2023). Cytokines from ILC2 cells also act to recruit effector cells, including eosinophils, alternatively activated macrophages, and IgE- and IgG1-producing B cells, to resolve the infection, as well as promote tissue repair (Hartung and Julia 2022; Vivier *et al.* 2018). In the face of these protective mechanisms, nematodes have evolved multiple strategies to manipulate or suppress host immunity, enabling them to persist within their host for extended periods (Maizels *et al.* 2018). Much of our knowledge of nematode immunomodulation stems from studying worm excretory–secretory (ES) products and their effects on mammalian cells using immortalized cell lines or *in vivo* animal models (Maizels *et al.* 2018). Organoids ('mini-tissues') provide a new and powerful *in vitro* platform for studying host–pathogen interactions and immunomodulation. This review focuses on the application of organoids to advance understanding of host–parasite interactions and immunomodulation in the GI tract.

The need for more physiological model systems

Traditional cell culture models are widely used due to their ease of maintenance and scalability; however, they have significant limitations. Cell lines are often derived from cancerous or immortalized tissues, which compromise their ability to accurately mimic native epithelial cells (AGUILAR *et al.* 2021). For example, Caco-2 and HT-29 cell lines are derived from human colorectal adenocarcinoma and are used as *in vitro* models of gut epithelium (Haddad *et al.* 2023). However, they lack proper differentiation and fail to replicate the structural and functional complexity needed for studying host–parasite interactions. Even when grown in three-dimensional (3D) cultures, these cell lines do not capture the cellular diversity of the intestinal epithelium, limiting their utility in modelling *in vivo* conditions.

Primary cell cultures and tissue explants are physiologically relevant alternatives, as they maintain native tissue architecture and functionality (Schlaermann *et al.* 2016). However, their limited expansion and requirement to be derived from fresh tissue present significant challenges, limiting their application in long-term studies. Animal models, particularly rodents, have been instrumental in providing valuable insights into both local and systemic host responses to parasitic infections of humans and animals (Mukherjee *et al.* 2022). For example, the murine whipworm *Trichuris muris* is a useful model for the human parasite *Trichuris trichiura*, allowing researchers to study host immunity, parasite biology and host genetics in a controlled setting (Klementowicz *et al.* 2012). However, for many parasitic helminth species, no appropriate model system supports parasite development and maturation that mimics *in vivo* infection. Additionally, animal use comes with significant challenges, including the need for specialized housing, high maintenance costs, ethical concerns, and limited ability to track host–parasite interactions over the course of infection (de Graeff *et al.* 2019).

Organoid-based models have the potential to overcome these limitations by providing a controllable and scalable platform to investigate host–parasite interactions and, for some parasites, initial invasion. The development of organoids from tissue stem cells of human, murine, ruminant and other animal species presents a promising alternative, offering a physiologically relevant system for studying important helminth infections, while reducing reliance on murine models. There is increasing interest in organoid-based systems as alternative models, to bridge the gap between cell culture approaches and whole-organism studies.

Organoid types

Organoids are self-organising 3D structures developed from tissue stem cells or induced pluripotent stem cells. Using culture media containing defined growth factors and nutrients, the stem cells differentiate and form 3D structures with similar organization and function to the tissue of origin (Lancaster and Knoblich 2014; Sato and Clevers 2015; Sato *et al.* 2009). Organoids have been grown from a range of tissue stem cells including gut (abomasum in ruminants), SI, colon, lung, liver, kidney, brain, bile duct, retinal and pancreas (Kawasaki *et al.* 2022; Tang *et al.* 2022). They can be propagated and passaged as long as stem cells are present and can be cryopreserved, enabling the establishment of organoid biobanks (Perrone and Zilbauer 2021).

Importantly, although organoids typically grow as 3D structures embedded in matrices such as Matrigel, they can also be dissociated into single cells or small clusters and cultured as two-dimensional (2D) monolayers. This process involves plating organoid-derived cells onto coated culture plates, where they attach and form confluent monolayers. While these monolayers do not replicate the full 3D architecture of the tissue, they retain many aspects of cellular differentiation and polarization (Duque-Correa *et al.* 2022; White *et al.* 2024) and offer the advantage of more direct access to the apical (luminal) surface (Wang *et al.* 2017). For studying the GI tract, organoids have been developed using stem cells derived from gastric (abomasal), duodenal, ileum, jejunum, caecum and colon tissue (Barker *et al.* 2010; Jung *et al.* 2011; Smith *et al.* 2021; Williamson *et al.* 2013; Yui *et al.* 2012), with subsequent differentiation into secretory epithelial cell types (Paneth, tuft, goblet (mucous producing) and enteroendocrine cells), and absorptive enterocytes (Fukuda *et al.* 2014; Sato and Clevers 2015). The different cell types can be identified by immunohistochemistry on fixed organoids using antibodies to specific marker proteins, such as anti-POU2F3 or anti-DCLK-1 for tuft cells, anti-MUC2 for mucous cells and anti-LGR5 for stem cells (Gerbe *et al.* 2016; Perez *et al.* 2025). Additionally, reporter organoids expressing a fluorescence gene under the control of a specific cell marker gene promoter enable analysis and imaging in real-time. Examples of 3D GI organoids stained with antibodies or expressing a reporter gene (tuft cell *Dclk-1* promoter driving tdTomato) are shown in Figure 1, and detailed in Perez *et al.* (2025).

Considerations for studying nematode–host interactions using organoids

Developing organoid models to study parasites requires consideration of fundamental aspects of host–parasite interactions: what is the natural site of infection within the host, which host cell types are primarily targeted and/or modulated by the parasite, are there differences between hosts, which parasite life-stage(s) or molecular components are most relevant for investigation, how can parasites or parasite molecules be delivered into organoids and how can parasite-mediated effects be assayed? We will firstly consider host tissue and cell types and in the next section discuss organoid manipulation and monitoring of parasite-mediated effects on organoids.

Infection site tissue type

GI nematodes occupy various niches within the host, including lumen or submucosa depending on the parasite developmental stage and species, where they can induce substantial alterations




Figure 1. Differentiation of gastrointestinal organoids and cell type identification. (A) Confocal microscopy image showing MUC2⁺ cells in ovine small intestinal (SI) organoids cultured in organoid growth medium (OGM) for 4 days. (B) Representative images of *Dclk1*-tdTomato⁺ tuft cells (red tdTomato⁺ cells indicated by arrowhead) in murine SI organoids, either untreated or treated with IL-13 in OGM for 4 days. (C) Representative image of ovine abomasal (gastric) organoids cultured in OGM for 4 days, with or without IL-13 treatment.

in epithelial cells. While adult worms are confined to the intestinal lumen, infective third-stage larvae (L3) of some nematode species actively penetrate the GI epithelial barrier, as seen with *Heligmosomoides polygyrus*, *T. trichiura* and *Strongyloides stercoralis*, which invade the submucosa. Blood-feeding nematodes, such as *H. contortus* and human hookworms *Ancylostoma duodenale* and *N. americanus*, attach to the intestinal mucosa causing mechanical damage and blood loss. An intriguing aspect of some GI nematodes, such as *Ascaris* and hookworms, is the requirement for hepato-tracheal migration of L3 prior to maturation in the SI (Read and Skorping 1995). These diverse interactions highlight the range of host–parasite relationships that influence disease pathogenesis and immune responses, and the relevance of developing organoids from the appropriate host tissue.

When studying parasite effects on host organoids, it is important to recognize that nematodes exhibit specific tropisms for distinct regions of the GI tract, certain cell types, and may remain in the lumen or invade submucosal tissue. A recent study by White et al. (2024) established confluent 2D SI monolayers to investigate interactions between *Heligmosomoides bakeri* L4 or adult stages and host duodenal cells. Notably, when the basal side of the epithelial cells was exposed to L4 or adult *Heligmosomoides*

bakeri, the cellular response was greater than when the apical side was exposed, perhaps reflecting differences in expression of Pattern Recognition Receptors between the apical and basal surfaces (Price et al. 2018). Additionally, L4 parasites, which develop submucosally, induced greater upregulation of epithelial interferon-stimulated genes and repair genes compared to adult worms, which reside in the lumen. These findings underscore the importance of considering the tissue site of infection, orientation of epithelial cells and the relevant nematode developmental stages to dissect the dynamics of infection at both the host and parasite levels.

Parasite immunomodulation in organoids

While organoids mimic features of the tissue of origin, they lack the complexity of associated mesenchymal, stromal or immune cells. Parasite–epithelial–immune cell interactions can be examined using co-culture models (see below) or, alternatively, specific cytokines can be added to organoid cultures. For example, addition of type-2 cytokines IL-4 and IL-13 to murine SI organoids promotes organoid differentiation (budding), specifically expansion of secretory tuft and mucous cell types (see Figure 1B, C

and Drurey *et al.* 2021; Kato-Atar *et al.* 2022). This mimics the *in vivo* effects of these cytokines on epithelial cells during nematode infection and provides a useful platform to examine infection-associated changes and parasite-mediated immunomodulation. Using this approach, Drurey *et al.* (2021) and Kato-Atar *et al.* (2022) demonstrated that *H. polygyrus* adult ES products suppress IL-4/IL-13-induced tuft and goblet cell expansion, demonstrating the parasite's ability to dampen host innate responses by directly acting on epithelial cells. In contrast to IL-4/IL-13 treatment, GI organoids treated with IL-22, which is mainly produced by T helper 1 (Th1), Th17 and group 3 innate lymphoid cells (ILC3) (Keir *et al.* 2020; Perusina Lanfranca *et al.* 2016) results in reduced organoid budding, a dark appearance and increased expression of goblet cell marker Resistin-like beta molecule (RELM- β) (Lindholm *et al.* 2022). Interestingly, Lindholm *et al.* (2022) further demonstrated that BMP signalling within intestinal organoids acts as a feedback mechanism to limit IL-13-induced tuft cell hyperplasia, highlighting how organoids can effectively model cytokine-driven epithelial differentiation and regulatory pathways observed *in vivo*, providing a valid platform to study immunomodulation.

Organoids to study the establishment of infection

In vitro culture of parasitic helminths remains a challenge, making it difficult to examine helminth development and tissue invasion. Elegant studies from the Duque-Correa group (Duque-Correa *et al.* 2022) established 2D caecal epithelium grown in transwells to study nematode larval invasion and development. They showed that *T. muris* L1 larvae could degrade secreted mucus and invade intestinal epithelial cells within murine caecal organoids (caecaloids), replicating the early infection processes that occur *in vivo* and establishing the first *in vitro* system for whipworm infection. This was achieved using 2D epithelial cell cultures, derived from 3D organoids and later grown in transwells, where the apical intestinal surface was accessible to the larvae. The 2D system overcame the challenge of trying to deliver larvae to the lumen (apical side) of 3D organoids, which is prohibited by the relatively large size of most nematode larvae. Further studies using a transgel GI organoid system – grown on hydrogel to generate bilaterally accessible organoids – enabled live observation of epithelial cell infection by *T. muris* L1, accompanied by formation of syncytial tunnels as occurs *in vivo* and subsequent epithelial cell apoptosis (Hofer *et al.* 2024). This demonstrates the utility and power of organoids to examine in detail the processes of parasite infection and establishment *in vitro*.

Organoid technology can also be applied to other types of helminths; recently, spheroids, 3D structures comprising a single cell type rather than differentiated cell types, were developed to study growth of liver fluke *Fasciola hepatica* newly excysted juveniles (NEJ) (Vitkauskaitė *et al.* 2025). The spheroid culture was grown from HepG2 liver cells and was able to support growth of *F. hepatica*, with development of gut, muscle and tegumental structures, as well as secretion of digestive proteases, similar to fluke development *in vivo*. The NEJ grazed on the peripheral cells, rather than invading the spheroid structure.

Host differences

Organoids are generated from single stem cells from specific individuals and retain characteristics of the original tissue, such as

gene expression profiles and treatment responses (Bresnahan *et al.* 2020). To ensure robust and reproducible results, it is recommended that experiments include organoid lines derived from at least three biological replicate tissue samples. This practice helps to account for natural biological variability and enhance the reliability of the findings. On the other hand, the characteristics maintained in organoids provide a valuable platform for determining differences between individuals or animal breeds/strains, helping to determine why they respond differently to infections (Grencis 2015). This has application in determining mechanisms of resistance or susceptibility, relevant to animal selective breeding programmes, vaccine development and personalized medicine (Bose *et al.* 2021).

Delivery into organoids

While 3D organoids provide a more physiological system than cell culture or 2D monolayers, their 3D structure can make the delivery and uptake of parasites or their products to the apical (luminal) side more challenging. Different approaches have been tested to achieve this and, in some studies, where long-term culture is not a requirement, 2D monolayers are appropriate. In this section, we discuss some of the delivery approaches used, which are summarized in Table 1.

Microinjection

Microinjection was one of the pioneering techniques applied to organoids, allowing the introduction of molecules directly to the apical side of 3D organoids. The first studies to focus on host-parasite interactions used microinjection to deliver extracellular vesicles (EVs) purified from ES products of adult murine whipworm, *T. muris*, into murine colonic organoids (Eichenberger *et al.* 2018a). Following injection into the organoid lumen, labelled EVs were detected within the cytoplasm of organoid cells. Cellular uptake was observed at 37°C but not at 4°C, indicating this is an active process. Similar findings were observed using microinjection to deliver EVs from *N. brasiliensis* adult ES into murine SI organoids (Eichenberger *et al.* 2018b). While effects directly on organoids were not reported in these studies, *in vivo* administration of EVs from *N. brasiliensis* ES, but not *T. muris*, protected mice from chemically induced colitis, with reductions in pro-inflammatory cytokines IL-6 and IFN- γ (Eichenberger *et al.* 2018b). By performing microinjection of EVs from *T. muris* into murine caecaloids followed by RNA sequencing, Duque-Correa *et al.* (2020) showed that anti-inflammatory effects, including down-regulation of Type I interferon signalling, can be mediated directly on the caecal epithelium in the absence of immune cell types. The specific parasite-derived molecules responsible for these effects are not yet known but may involve proteins and/or small RNAs within EVs.

Microinjection allows precise targeting of specific regions within the organoid; however, it has several limitations. The variability in organoid size can lead to inconsistencies in injection efficiency, making the technique less reliable. Additionally, microinjection is labour-intensive and requires skilled researchers and specialized equipment. It is also impractical to deliver nematodes into the organoid lumen, as the limited luminal volume of organoids is insufficient to accommodate most nematode parasites (Duque-Correa *et al.* 2020).

Table 1. Delivery approaches used to study interaction of GI nematodes with epithelial cells

Delivery method	Parasite stage or component	Organoid type	References
Microinjection	<i>T. muris</i> (adult) EVs	Murine colonic 3D organoids	Eichenberger et al. 2018a; Duque-Correa et al. 2020
	<i>N. brasiliensis</i> (adult) EVs	Murine SI 3D organoids	Eichenberger et al. 2018b
Direct addition of nematodes	<i>T. circumcincta</i> (L3)	Ovine SI 3D organoids	Smith et al. 2021
	<i>O. ostertagi</i> (L3)	Bovine gastric 3D organoids	Faber et al. 2022
	<i>T. muris</i> (L1)	Murine SI 2D culture	Duque-Correa et al., 2022
	<i>T. spiralis</i> (L1)	Porcine SI 3D organoids	Liu et al. 2024
	<i>H. bakeri</i> (L4 and adult)	Murine SI 2D cultures	White et al. 2024
	<i>P. univalens</i> , Cyathostominae and <i>S. vulgaris</i> (L3)	Equine SI 2D cultures	Hellman et al. 2024
Direct addition of nematode-secreted products	<i>H. polygyrus</i> (adult) ES	Murine SI 3D organoids	Drurey et al. 2021; Karo-Atar et al. 2022
	<i>H. contortus</i> (adult) ES	Ovine gastric 3D organoids	Perez et al. 2025
		Ovine SI 3D organoids	
		Murine SI 3D organoids	
	<i>T. spiralis</i> (L1) ES	Porcine SI 3D organoids	Liu et al. 2024
	<i>Anisakis</i> (L3) EVs	Human 2D colonic organoids	Bellini et al. 2024
	MicroRNA – 5352 mimic	Ovine gastric 3D organoids	Perez et al. 2025
		Ovine SI 3D organoids	
		Murine SI 3D organoids	

EVs, extracellular vesicles; ES, excretory-secretory products.

Apical out organoids

An alternative for delivering parasites or ES products to the apical surface is to invert the polarity of 3D organoids using a method developed by Co et al. (2019). This involves removing the extracellular matrix (ECM) from the Matrigel (basement membrane extract), which is added to stem cells to support organoid formation (Co et al. 2021). Organoids are then incubated in low-binding culture plates, which induce a reversal of polarity (apical-out) while preserving barrier integrity and key functional characteristics. This technique was successfully applied to abomasal and SI organoids from different animal species (Smith et al. 2021), allowing access to the apical surface for experimental manipulation. However, the host-driven mechanisms that govern organoid invagination and polarity reversal remain poorly understood and apical-out organoids do not differentiate in the same way as apical-in organoids (Paužuolis et al. 2024). Smith et al. (2021) were able to demonstrate infection of apical-out organoid cells with the bacterial pathogen *Salmonella typhimurium*, but this method has not yet been explored to study nematode infectivity or development.

Direct addition of nematode larvae or ES products to 3D organoid culture medium

An alternative to microinjection is the direct addition of nematodes or their secreted products to organoid culture medium. Following incubation, larvae or ES can enter into the organoid lumen and/or cells. This strategy was used to introduce infective L3 of the important veterinary nematodes *Teladorsagia circumcincta* and *O. ostertagi* into ovine or bovine abomasal (gastric) organoids, respectively (Faber et al. 2022; Smith et al. 2021). Strikingly, the L3 larvae transversed the Matrigel and organoid membrane, from

the basal (outer) to apical (inner) side into the lumen (Faber et al. 2022; Smith et al. 2021). Larvae survived in the lumen for up to two weeks, were active and invaded organoid cells, but did not develop beyond L3 stage (Faber et al. 2022; Smith et al. 2021). This implies that the organoid model may not fully replicate the route of invasion and/or the *in vivo* conditions that support larval development. Future work could explore the feasibility of 2D monolayers, as used for *T. muris* (Duque-Correa et al. 2022), to promote larval development. Nonetheless, 3D organoids can provide valuable insights into the ability of larvae to interact with and cross epithelial barriers. Intriguingly, organoids exposed to *O. ostertagi* L3 or ES products showed a rapid expansion or ‘ballooning’ within 1 hour of treatment, which is thought to result from fluid influx into the lumen (Faber et al. 2022).

The ease of adding soluble substances to organoid culture medium and then monitoring effects has encouraged greater use of this approach. Several studies (Drurey et al. 2021; Karo-Atar et al. 2022; Perez et al. 2025) showed that ES from adult *H. polygyrus* or *H. contortus* can suppress the expansion of tuft and mucous cells driven by IL-4/IL-13, demonstrating a direct effect of ES on epithelial responses. Identifying the molecules and mechanisms involved has the potential to lead to new therapeutics to block nematode-mediated immunosuppression, while conversely, such parasite molecules could be developed as ‘helminth therapy’ to suppress GI inflammatory conditions, such as inflammatory bowel disease or allergies (Ryan et al. 2020). *T. spiralis* L1 larvae or ES were also shown to inhibit pro-inflammatory responses, reducing the pathology induced in SI organoids by porcine epidemic diarrhea virus (Liu et al. 2024).

These studies also revealed that ES from adult *H. polygyrus* or *H. contortus* alone in the absence of cytokine treatment had

a strong inhibitory effect on SI organoid budding and development. Large spheroid organoids devoid of differentiated cells were observed following 24 h of exposure to ES or L3 larvae, indicating that nematode-secreted molecules induce de-differentiation of organoids and lead to a fetal-like and/or repair state (Drurey et al. 2021; Karo-Atar et al. 2022; Perez et al. 2025). A similar phenotype had previously been observed following growth of organoids derived from SI stem cells from *H. polygyrus*-infected mice, or after IFN- γ treatment of SI organoids, recapitulating the fetal-like granuloma-associated cells occurring *in vivo* during *H. polygyrus* infection (Nusse et al. 2018). Our laboratory found that spheroid formation occurred only with *H. contortus* ES that had been concentrated (through a 3-kDa membrane), but not with unconcentrated ES, indicating a concentration-dependent effect (Perez et al. 2025), analogous to the localized effects *in vivo*. Spheroid formation in organoids was not observed with ES from adult *N. brasiliensis*, which is cleared rapidly from the murine SI (Karo-Atar et al. 2022), suggesting that epithelial de-differentiation may promote nematode survival in the GI tract. This effect is different to the organoid swelling observed with *O. ostertagi* ES or L3 (Faber et al. 2022), which happens rapidly (within 1 h). For both phenotypes, organoids provide an appropriate *in vitro* system to help understand changes induced in epithelial cells during *in vivo* infection.

Organoid transfection with parasite small RNAs

The above studies demonstrate the utility of organoids in dissecting biological activity of nematode ES products on the GI epithelium. Previous characterization of nematode ES identified proteins with immunomodulatory activity (Hewitson et al. 2009; Maizels et al. 2018); however, little is known about the biological effects of nematode-secreted small RNAs. Small regulatory microRNAs (miRNAs) modulate gene expression post-transcriptionally by binding to the 3'UTR of their target mRNAs, resulting in inhibition of protein translation and transcript degradation (Chekulaeva and Filipowicz 2009). The repertoires of miRNAs expressed and secreted by a variety of nematode species have been described (Britton et al. 2020; Buck et al. 2014). Our laboratory focussed on a single miRNA, miR-5352, which is conserved across GI parasitic nematodes, but not present in tissue-dwelling nematodes (Gu et al. 2017; Perez et al. 2025). This led our group to hypothesize that miR-5352 may be involved in regulating host gene expression in the GI tract, which we tested by transfecting SI organoids with a mimic of miR-5352.

Efficient delivery of the miRNA mimic into the cytoplasm of epithelial cells required transfection reagents with high efficiency, low toxicity and lipid formulations specifically designed for small RNA delivery. Organoids were exposed to miRNA mimic, labelled with Cy3 fluorescent dye, and transfection reagent (DharmaFECT TM) at 37°C for 1 h before being embedded in Matrigel and returned to culture (Perez et al. 2025). This approach significantly increased the number of transfected epithelial cells compared to using naked miRNA mimics, indicated by Cy3 label inside cells. Importantly, this single GI nematode miRNA suppressed differentiation of tuft and mucous cells in the presence of IL-13, similar to the effect of *H. polygyrus* or *H. contortus* complete ES. Kruppel-like transcription factor gene *Klf-4* was identified as a key target of nematode miR-5352, and *KLF-4* expression was significantly reduced in miR-5352 transfected organoids (Perez et al. 2025). *KLF-4* promotes cell differentiation by inhibiting Wnt signalling, and loss of *KLF-4* leads to stem cell maintenance (Zhang

et al. 2006), thus identifying mechanistically how nematode ES can modulate gut cell differentiation. Interestingly, nematode miR-5352 shares its seed sequence (key regulatory sequence of miRNAs) with mammalian miR-92a, which also suppresses *Klf-4*, suggesting that GI nematodes have hijacked a host regulatory pathway to suppress innate cell responses and promote nematode survival.

2D monolayers to study host–parasite interactions


Development of 2D monolayers, derived from 3D organoids, provides an alternative, accessible apical surface for studying host–parasite interactions and parasite invasion, as demonstrated by the studies of White et al. (2024) using *H. bakeri*, and Duque-Correa et al. (2022) with *T. muris* L1 larvae. Hellman et al. (2024) utilized equine SI 2D cultures (enteroids) stimulated with IL-4 and IL-13 and exposed the apical side to infective larvae of *Parascaris univalens*, Cyathostominae and *Strongylus vulgaris*. Changes in epithelial cell morphology and an increase in MUC2 expression were observed with *P. univalens* L3, indicating a direct effect of larvae on mucous cells. Using human 2D colonic organoids, Bellini et al. (2024) reported that *Anisakis* L3 EVs have tumorigenic and immunomodulatory capabilities. Following organoid treatment with EVs there was a reduction in the receptor for IL-33, potentially dampening type 2 immunity and inflammatory responses.

These findings underscore the utility of different organoid technologies and delivery routes for advancing understanding of GI nematode infections and modulation of epithelial cells. Developments in organoid technology, including co-culture, will continue to improve their relevance to infection biology and as platforms to screen and evaluate new therapeutics.

Immune cell co-culture

Studying the effects of GI nematode infection on epithelial organoids has advanced our understanding of host–pathogen interactions; however, it is important to recognize that no cellular compartment reacts in isolation *in vivo*. Thus, to better appreciate the implications of GI nematode infection, additional layers of complexity should be incorporated in organoid–parasite models. Mucosal tissue resident immune cells involved in the mediation of type 2 immune responses are a relevant compartment to introduce to the culture systems.

Helminth infection models have been instrumental in the discovery of ILC2s, the innate counterparts of adaptive T helper 2 (Th2) cells (Moro et al. 2010). ILC2s were initially discovered in the adipose tissue of the peritoneal cavity of mice. Upon infection with *N. brasiliensis*, ILC2s produced IL-13 which facilitated goblet cell hyperplasia in the SI and thus helminth expulsion (Moro et al. 2010). Subsequently, ILC2s were also described in the SI tissue where they mediated IL-13–dependent protection against *N. brasiliensis* (Neill et al. 2010; Price et al. 2010) and *T. muris* (Saenz et al. 2010) by inducing the ‘weep and sweep’ response. Since their discovery, the roles of intestinal ILC2s in helminth immunity have been further characterized. *N. brasiliensis* infection in the intestine promoted the activation of ILC2s through the tuft cell–ILC2 axis (Schneider et al. 2018), where tuft cell-derived IL-25 activated ILC2 which in turn led to the production of IL-13. Additionally, IL-33 and TSLP secreted by epithelial cells upon helminth infection further promoted ILC2 activation and induction of a type 2 immune response, necessary for helminth clearance (Hung et al. 2020). Besides the intestine, the protective roles of ILC2s against *N. brasiliensis* larvae extended to the lung (Turner et al. 2013).



B




Figure 2. (A) Schematic representation of ILC and organoid co-culture set-up. In brief, small intestinal organoids are generated prior to the co-culture establishment. Organoids are split approximately 48 h prior to placing them together with the ILCs. Pan-ILCs (live, CD45⁺, Lin⁻ (CD3, CD45R, CD11b, TER-119, Ly-G6, CD19, CD5, CD127⁺)) or specific ILC groups of interest are isolated from the lamina propria of RORyt – GFP mice, using fluorescence activated cell sorting (FACS). Once isolated, the ILCs are placed together with the organoids, spun down, resuspended in matrigel and placed in a well. Cultures are fed every 2 days until the experimental end point. (B) Confocal microscopy image of ILC precursors, stained for the immune marker CD45, co-cultured with small intestinal organoids, stained for the epithelial cell adhesion molecule (EpCAM) (JOWETT et al. 2022) (<https://creativecommons.org/licenses/by/4.0/#ref-appropriate-credit>).

In the lung, ILC2s use IL-9 in an autocrine manner to enhance their function and support tissue repair during the recovery phase (Turner et al. 2013). Interestingly, helminths were able to interfere with ILC2 activation even when not co-localized. Upon infection with *H. polygyrus*, activation of ILC2s in the lung was hindered due to the release of immunomodulatory ES products, such as *H. polygyrus* Alarmin Release Inhibitor (HpARI) (Osbourne et al. 2017) and *H. polygyrus* Binds Alarmin Receptor and Inhibits (HpBARI) (Vacca et al. 2020).

Answering mechanistic questions and dissecting specific immune-epithelial interactions has become streamlined in recent years with the establishment of co-culture models (Kromann et al. 2024). Murine ILC-organoid co-culture models have been generated by isolating ILCs from the lamina propria (Jowett et al. 2021, 2022; Lindemans et al. 2015; Read et al. 2022, 2024) or the mesenteric lymph nodes (Waddell et al. 2019) using fluorescence activated cell sorting techniques and co-culturing them with GI organoids. Similarly, human models have been established where intestinal

ILC2s were co-cultured with intestinal organoids (Möller et al. 2023). Additionally, organoid-maturated human ILC2s have been successfully co-cultured with lung-derived organoids, showcasing the versatility of this model (Jowett et al. 2022). To generate this co-culture, immune cells and epithelial organoids are resuspended together in basement membrane cellular extract to provide the ECM cues which enables the two compartments to interact in a manner similar to that *in vivo* (Figure 2). While this system advanced our understanding of tissue specific immune-epithelial interactions, only looking at the two compartments in a homeostatic setting provides limited applicability and can be enhanced by an additional compartment – parasites.

A tri-culture model involving ILC2s, organoids, and parasites would help elucidate how specific parasites or parasite-derived proteins, miRNAs, or EVs (Buck et al. 2014; Gu et al. 2017; Perez et al. 2025; Soichot et al. 2022) influence gene expression in epithelial and immune cells, and how these changes affect cellular functions, immune responses, and overall host–parasite interactions.

Moreover, it would allow us to answer questions related to the functionality and behaviour of ILC2s under helminth infections in a controlled environment. Studies have shown that *Proteobacteria*, a gut microbiome species (Pu *et al.* 2021), or *N. brasiliensis* (Huang *et al.* 2018) facilitated the migration of ILC2s from the intestine to the lung. In addition to pathogen-derived signals promoting this migration, tissue-specific imprinting has been implicated in determining the function and fate of post-migratory gut ILC2s in the lung (Jowett *et al.* 2022). Additionally, under specific microenvironmental conditions, ILC2s displayed a degree of plasticity as they transdifferentiated into ILC1s and ILC3s (Bielecki *et al.* 2021; Qin *et al.* 2024). Recent evidence also shows that ILC3s contribute to protective immunity against helminths by promoting tuft cell hyperplasia via RANKL signalling, thereby supporting the ILC2–tuft cell feedback loop (Xu *et al.* 2025). In future studies, using such a tri-culture model could enable assessment of the impact of parasite and parasite-derived products on epithelial and immune cell modulation and functionality, providing valuable insights into host–parasite dynamics.

Once established and characterized, this tri-culture model could be taken further and additional cellular compartments could be incorporated, such as neurons. Neuroimmune interactions have been previously described in GI nematode infections and ILC2s co-localize with neurons in neuroimmune cell units (Veiga-Fernandes and Artis 2018). Enteric neurons, via their ability to produce neuropeptides such as neuromedin U (Cardoso *et al.* 2017) and vasoactive intestinal peptide (Nussbaum *et al.* 2013), promoted the activation of ILC2s, and production of IL-5, IL-9 and IL-13, which in turn led to helminth expulsion. On the other hand, adrenergic neurons produced norepinephrine which inhibited the function of ILC2s, hindering helminth expulsion (Moriyama *et al.* 2018). ILC2s were reported to synthesise the neurotransmitter acetylcholine in the presence of IL-33 which was required for an appropriate type 2 immune response against *N. brasiliensis* (Chu *et al.* 2021; Roberts *et al.* 2021). Interestingly, *N. brasiliensis* and other mucosal-dwelling nematodes synthesize acetylcholinesterases, inhibitors of acetylcholine (Sanderson 1969; Selkirk *et al.* 2005).

Taken together, future development and application of tri-culture models incorporating epithelial cells, immune cells, and parasites will be essential to advance our understanding of host–parasite dynamics and immune interactions in helminth infections.

Conclusions and future directions

Organoids provide a powerful platform for studying the complex cellular and molecular responses of the host epithelium to nematode infections and, conversely, how nematodes modulate the host environment. Organoid technology is progressing, and new systems are being developed to increase their physiological relevance and longer-term survival, including the use of air–liquid interface (ALI) models and tissue engineering combined with microfluidics. In ALI models, cells are exposed to air on the apical side while the basolateral side is in contact with culture medium containing differentiation factors. This combination enhances cellular differentiation and 3D formation, as observed in human intestinal organoids which form a defined crypt-villus-like structure under ALI conditions (Baldassi *et al.* 2021; Ogawa *et al.* 2025). Tissue engineering strategies, including ECM-embedded scaffolds,

bioreactors and organoids-on-a-chip, have been developed as powerful tools for modelling both healthy and diseased organs (Wang *et al.* 2024) and can support organoid culture and better simulate the natural tissue architecture. Dual-access microfluidic systems facilitate maintenance of nutrients/removal of waste and dead cells, delivery of cytokines to the basal side, and co-culture with additional components such as immune cells, vasculature and microbiome, providing a more physiologically relevant platform (Hofer *et al.* 2024; Quintard *et al.* 2024). By maintaining organoids in conditions more akin to *in vivo* conditions, their physiological relevance can be enhanced and their lifespan extended, making them attractive for drug screening and potentially for maintaining parasite lifecycles, a current challenge for most nematode species.

The majority of organoid–nematode studies to date have focused on murine organoid models and model parasite species. Although some studies have begun to use ruminant-derived organoids (Faber *et al.* 2022; Perez *et al.* 2025; Smith *et al.* 2021), there is a clear need to further expand these approaches to include organoids and parasites from a broader range of animal species, as well as humans, to enable true translational impact for both veterinary and human clinical applications.

Nematodes can induce changes in organoid phenotype and cellular composition, which can be detected using antibodies to cell marker proteins (Drurey *et al.* 2021; Karo-Atar *et al.* 2022) or use of transgenic reporter organoids (Perez *et al.* 2025). While the latter require more initial effort to generate, they facilitate the study and imaging of organoid cells in real-time, without the need for fixation or antibodies, which may not be available/applicable for less widely studied host species. Reporter organoids can be generated from genetically modified mice (and potentially other animals) (Perez *et al.* 2025; Wadosky *et al.* 2022), or more rapidly using CRISPR–Cas9 gene knock-in technology. This introduces a fluorescent reporter tag to a specific gene of interest by electroporation, and transfected organoids are selected based on fluorescence. Different methodologies have been developed for CRISPR–Cas9 knock-in (Artegiani *et al.* 2020), with CRISPR–HOT (homology-independent organoid transgenesis) shown to be efficient, enabling real-time cell identification and lineaging. CRISPR–Cas9 or RNA interference mediated by short interfering or short hairpin RNA can also be used to knockdown genes of interest in organoids to examine gene function in a controlled environment. Technologies for organoid culture and co-culture, and CRISPR–Cas9 gene editing, are continuing to expand, providing increasingly sophisticated methodologies to study host cell responses and modulation by parasites and other infectious organisms or disease conditions (Huber *et al.* 2023; Sun *et al.* 2021).

Acknowledgements. The authors gratefully acknowledge Colin Loney, Imaging Manager (Medical Research Council–University of Glasgow Centre for Virus Research) for help with imaging and Dr David Smith, Dr Katie Hildersley and Prof Tom McNeilly (Moredun Research Institute, Edinburgh) for provision of ovine organoids for some of the work described in this review. We thank Prof Eileen Devaney for comments on the manuscript.

Author contributions. MGP and CB conceived the idea and scope of the review. MGP and DC performed the literature search and drafted the initial manuscript. CB, DC and JN contributed to the critical revision of the content and figures. All authors contributed to the writing, reviewed the final version of the manuscript, and approved it for submission.

Financial support. Some of the data discussed here were supported by the Royal Society through a Newton International Fellowship (NIF\R1\201159),

a McIntyre International Research Fellowship (University of Glasgow) and research training awards funded by The Company of Biologists and SULSA (Scottish Universities Life Sciences Alliance) to MGP, and by a Wellcome Collaborative Award (Ref 211814) to CB, D.C. and J.F.N. acknowledge funding support by The Leona M. and Harry B. Helmsley Charitable Trust. In addition, J.F.N. acknowledges funding from The Lister Institute of Preventive Medicine and from the BBSRC and the NC3Rs (NC/X002497/1).

Competing interests. J.F.N. is inventor on a patent application related to this work (WO/2024/256627). The remaining authors declare there are no competing interests.

Ethical standards. Not applicable.

References

- Aguilar C., Alves da Silva M, Saraiva M, Neyazi M, Olsson I A S and Bartfeld S (2021) Organoids as host models for infection biology - a review of methods. *Experimental Molecular Medicine* 53, 1471–1482. doi:10.1038/s12276-021-00629-4
- Artigiani B., Hendriks D, Beumer J, Kok R, Zheng X, Joore I, Chuva de Sousa L S, van Z J, Tans S and Clevers H (2020) Fast and efficient generation of knock-in human organoids using homology-independent CRISPR-Cas9 precision genome editing. *Nature Cell Biology* 22, 321–331. doi:10.1038/s41556-020-0472-5
- Artis D. and Grecis R K (2008) The intestinal epithelium: Sensors to effectors in nematode infection. *Mucosal Immunology* 1(4), 252–264. doi:10.1038/mi.2008.21
- Baldassi D., Gabold B and Merkel O (2021) Air-liquid interface cultures of the healthy and diseased human respiratory tract: Promises, challenges and future directions. *Advances Nanobiomedicine Research* 1, 2000111. doi:10.1002/anbr.202000111
- Barker N., Huch M, Kujala P, van de Wetering M, Snippert H J, van E J H, Sato T, Stange DE, Begthel H, van den Born M and Danenberg E. (2010) Lgr5(+ve) stem cells drive self-renewal in the stomach and build long-lived gastric units in vitro. *Cell Stem Cell* 6(1), 25–36. doi:10.1016/j.stem.2009.11.013.
- Bąska P. and Norbury L J (2022) The role of the intestinal epithelium in the “weep and sweep” response during gastro-intestinal helminth infections. *Animals* 12, 175. doi:10.3390/ani12020175
- Bellini I., Scribano D, Ambrosi C, Chiovoloni C, Rondón S, Pronio A, Palamara A T, Pietrantonio A, Kashkanova A, Sandoghdar V, D’amelio S and Cavallero S (2024) Anisakis extracellular vesicles elicit immunomodulatory and potentially tumorigenic outcomes on human intestinal organoids. *Parasites Vectors* 17(1), 393. doi:10.1186/s13071-024-06471-7
- Bielecki P., Riesenfeld S J, Hütter J C, Torlai Triglia E, Kowalczyk M S, Ricardo-Gonzalez R R, Lian M, Amezcua Vesely M C, Kroehling L, XU H, Slyper M, Muus C, Ludwig L S, Christian E, Tao L, Kedaigle A J, Steach H R, York A G, Skadow M H, Yaghoubi P, Dionne D, Jarret A, McGee H M, Porter C B M, Licon-Limón P, Bailis W, Jackson R, Gagliani N, Gasteiger G, Locksley R M, Regev A and Flavell R A (2021) Skin-resident innate lymphoid cells converge on a pathogenic effector state. *Nature* 592(7852), 128–132. doi:10.1038/s41586-021-03188-w
- Bose S., Clevers H and Shen X (2021) Promises and challenges of organoid-guided precision medicine. *Medicine* 2, 1011–1026. doi:10.1016/j.medj.2021.08.005
- Bresnahan E., Ramadori P, Heikenwalder M, Zender L and Lujambio A (2020) Novel patient-derived preclinical models of liver cancer. *Journal Hepatology* 72, 239–249. doi:10.1016/j.jhep.2019.09.028
- Britton C., Laing R and Devaney E (2020) Small RNAs in parasitic nematodes – Forms and functions. *Parasitology* 147, 855–864.
- Buck A. H., Coakley G, Simbari F, McSorley H J, Quintana J F, le Bihan T, Kumar S, Abreu-Goodger C, Lear M, Harcus Y and Ceroni A (2014) Exosomes secreted by nematode parasites transfer small RNAs to mammalian cells and modulate innate immunity. *Nature Communications* 5, 5488. doi:10.1038/ncomms5488.
- Cardoso V., Chesné J, Ribeiro H, García-Cassani B, Carvalho T, Bouchery T, Shah K, Barbosa-Morais N L, Harris N and Veiga-Fernandes H (2017) Neuronal regulation of type 2 innate lymphoid cells via neuromedin U. *Nature* 549(7671), 277–281. doi:10.1038/nature23469
- Charlier J., Bartley D J, Sotiraki S, Martinez-Valladares M, Claerebout E, von Samson-himmelstjerna G, Thamsborg S M, Hoste H, Morgan E R and Rinaldi L (2022) Anthelmintic resistance in ruminants: Challenges and solutions. *Advances Parasitology* 115, 171–227. doi:10.1016/bs.apar.2021.12.002
- Charlier J., Rinaldi L, Musella V, Ploeger H W, Chartier C, Vineer H R, Hinney B, von Samson-himmelstjerna G, Bacescu B, Mickiewicz M and Mateus T L (2020) Initial assessment of the economic burden of major parasitic helminth infections to the ruminant livestock industry in Europe. *Preventative Veterinary Medicine* 182, 105103. doi:10.1016/j.pvetmed.2020.105103
- Chekulaeva M. and Filipowicz W (2009) Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells. *Current Opinion Cell Biology* 21, 452–460. doi:10.1016/j.ccb.2009.04.004
- Chu C., Parkhurst C N, Zhang W, Zhou L, Yano H, Arifuzzaman M and Artis D (2021) The ChAT-acetylcholine pathway promotes group 2 innate lymphoid cell responses and anti-helminth immunity. *Science Immunology* 6(57), eabe3218. doi:10.1126/sciimmunol.abe3218
- CO J. Y., Margalef-Català M, Li X, Mah A T, Kuo C J, Monack D M and Amieva M R (2019) Controlling Epithelial Polarity: A Human Enteroid Model for Host-Pathogen Interactions. *Cell Reports* 26(9), 2509–2520.e4. doi:10.1016/j.celrep.2019.01.108
- CO J. Y., Margalef-Català M, Monack D M and Amieva M R (2021) Controlling the polarity of human gastrointestinal organoids to investigate epithelial biology and infectious diseases. *Nature Protocols* 16, 5171–5192. doi:10.1038/s41596-021-00607-0
- Corning S. (2009) Equine cyathostomins: A review of biology, clinical significance and therapy. *Parasites and Vectors* 2(Suppl 2), S1. doi:10.1186/1756-3305-2-S2-S1
- de Graeff N., Jongsma K R, Johnston J, Hartley S and Bredenoord A L (2019) The ethics of genome editing in non-human animals: A systematic review of reasons reported in the academic literature. *Philosophical Transactions Royal Society London B Biology Science* 374, 20180106. doi:10.1098/rstb.2018.0106
- Drurey C., Lindholm H T, Coakley G, Poveda M C, Löser S, Doolan R, Gerbe F, Jay P, Harris N, Oudhoff M J and Maizels RM. (2021) Intestinal epithelial tuft cell induction is negated by a murine helminth and its secreted products. *Journal Experimental Medicine* 219, e20211140. doi:10.1084/jem.20211140.
- Duque-Correa M. A., Goulding D, Rodgers F H, Gillis J A, Cormie C, Rawlinson K A, Bancroft A J, Bennett H M, Lotkowska M E, Reid A J, Speak A O, Scott P, Redshaw N, Tolley C, McCarthy C, Brandt C, Sharpe C, Ridley C, Moya J G, Carneiro C M, Starborg T, Hayes K S, Holroyd N, Sanders M, Thornton D J, Grecis R K and Berriman M (2022) Defining the early stages of intestinal colonisation by whipworms. *Nature Communications* 13, 1725. doi:10.1038/s41467-022-29334-0
- Duque-Correa M. A., Schreiber F, Rodgers F H and Berriman M (2020) Advances and perspectives in the use of in vitro models to study host-parasite interactions. *Frontiers in Cellular and Infection Microbiology* 10, 339. doi:10.3389/fcimb.2020.00339
- Eichenberger R. M., Ryan S, Jones L, Buitrago G, Polster R, Montes de Oca M, Zuvelek J, Giacomini P R, Dent L A, Engwerda C R, Field M A, Sotillo J and Loukas A (2018b) Hookworm secreted extracellular vesicles interact with host cells and prevent inducible colitis in mice. *Frontiers Immunology* 9, 850. doi:10.3389/fimmu.2018.0085.
- Eichenberger R. M., Talukder M H, Field M A, Wangchuk P, Giacomini P, Loukas A and Sotillo J (2018a) Characterization of *Trichuris muris* secreted proteins and extracellular vesicles provides new insights into host-parasite communication. *Journal Extracellular Vesicles* 7, 1428004. doi:10.1080/20013078.2018.1428004
- Faber M. N., Smith D, Price D R G, Steele P, Hildersley K A, Morrison L J, Mabbott N A, Nisbet A J and McNeilly T N (2022) Development of bovine gastric organoids as a novel in vitro model to study host-parasite interactions in gastrointestinal nematode infections. *Frontiers Cellular Infection Microbiology* 12, 904606. doi:10.3389/fcimb.2022.904606

- Fallon P. G., Ballantyne S J, Mangan N E, Barlow J L, Dasvarma A, Hewett D R, McIlgorm A, Jolin H E and McKenzie A N J (2006) Identification of an interleukin (IL)-25 dependent cell population that provides IL-4, IL-5, and IL-13 at the onset of helminth expulsion. *Journal Experimental Medicine* **203**, 1105–1116. doi:10.1084/jem.20051615
- Fukuda M., Mizutani T, Mochizuki W, Matsumoto T, Nozaki K, Sakamaki Y, Ichinose S, Okada Y, Tanaka T, Watanabe M and Nakamura T (2014) Small intestinal stem cell identity is maintained with functional Paneth cells in heterotopically grafted epithelium onto the colon. *Genes and Development* **28**(16), 1752–1757. doi:10.1101/gad.245233.114
- Gerbe F., Sidot E, Smyth D J, Ohmoto M, Matsumoto I, Dardalhon V, Cesses P, Garnier L, Pouzolles M, Brulin B and Bruschi M. (2016) Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature* **529**, 226–230. doi:10.1038/nature16527.
- Grencis R. K., (2015) Immunity to helminths: Resistance, regulation, and susceptibility to gastrointestinal nematodes. *Annual Review of Immunology* **33**, 201–225.
- GU H. Y., Marks N D, Winter A D, Weir W, Tzelos T, McNeilly T N, Britton C and Devaney E (2017) Conservation of a microRNA cluster in parasitic nematodes and profiling of miRNAs in excretory-secretory products and microvesicles of *Haemonchus contortus*. *PLoS Neglected Tropical Diseases* **11**(11), e0006056. doi:10.1371/journal.pntd.0006056
- Haddad M. J., Sztupecki W, Delayre-Orthez C, Rhazi L, Barbezier N, Depeint F and Anton P M (2023) Complexification of In Vitro Models of Intestinal Barriers, A True Challenge for a More Accurate Alternative Approach. *International Journal of Molecular Sciences* **24**(4), 3595. doi:10.3390/ijms24043595
- Hardman C. S., Panova V and McKenzie A N J (2013) IL-33 citrine reporter mice reveal the temporal and spatial expression of IL-33 during allergic lung inflammation. *European Journal of Immunology* **43**, 488–498.
- Hartung F. and Julia E-V B (2022) Trained immunity in type 2 immune responses. *Mucosal Immunology* **15**, 158–1169. doi:10.1038/s41385-022-00557-0
- Hewitson J. P., Grainger J R and Maizels R M (2009) Helminth immunoregulation: The role of parasite secreted proteins in modulating host immunity. *Molecular and Biochemical Parasitology* **167**, 1–11. doi:10.1016/j.molbiopara.2009.04.008
- Hofer M., Duque-Correa M A and Lutolf M P (2024) Patterned gastrointestinal organoid monolayers with bilateral access as observable models of parasite gut infection. *Nature Biomedical Engineering* **9**(7), 1075–1085. doi:10.1038/s41551-024-01313-4
- Hotez P. J. and Herricks J R (2015) Helminth elimination in the pursuit of sustainable development goals: a “worm index” for human development. *PLoS Neglected Tropical Diseases* **9**(4), e0003618. doi:10.1371/journal.pntd.0003618
- Howitt M. R., Lavoie S, Michaud M, Blum A M, Tran S V, Weinstock J V and Artis D (2016) Tuft Cells, Taste-Chemosensory Cells, Orchestrate Parasite Type 2 Immunity in the Gut. *Science* **351**, 1329–1333. doi:10.1126/science.aaf1648.
- Huang Y., Mao K, Chen X, Sun M A, Kawabe T, Li W, Usher N, Zhu J, JF U, Jr, Paul W E and Germain R N (2018) S1P-dependent interorgan trafficking of group 2 innate lymphoid cells supports host defense. *Science* **359**(6371), 114–119. doi:10.1126/science.aam5809
- Huber A., Dijkstra C and Ernst M E M F (2023) Generation of gene-of-interest knockouts in murine organoids using CRISPR-Cas-9. *Star Protocols* **4**, 102076. doi:10.1016/j.xpro.2023.102076
- Hung L. Y., Tanaka Y, Herbine K, Pastore C, Singh B, Ferguson A, Vora N, Douglas B, Zullo K, Behrens E M, Li Hui Tan T, Kohanski M A, Bryce P, Lin C, Kambayashi T, Reed D R, Brown B L, Cohen N A and Herbert D R (2020) Cellular context of IL-33 expression dictates impact on anti-helminth immunity. *Science Immunology* **5**(53), eabc6259. doi:10.1126/sciimmunol.abc6259
- Hellman S., Martin F, Tydén E, Sellin M E, Norman A, Hjertner B, Svedberg P and Fossum C (2024) Equine enteroid-derived monolayers recapitulate key features of parasitic intestinal nematode infection. *Veterinary Research* **55**(1), 25. doi:10.1186/s13567-024-01266-1.
- Jowett G. M., Norman M D A, YU T T L, Rosell Arévalo P, Hoogland D, Lust S T, Read E, Hamrud E, Walters N J, Niazi U, Chung M W H, Marciano D, Omer O S, Zabinski T, Danovi D, Lord G M, Hilborn J, Evans N D, Dreiss C A, Bozec L, Oommen O P, Lorenz C D, da Silva R M P, Neves J F and Gentleman E (2021) ILC1 drive intestinal epithelial and matrix remodelling. *Nature Materials* **20**(2), 250–259. doi:10.1038/s41563-020-0783-8
- Jowett G. M., Read E, Roberts L B, Coman D, Vilà González M, Zabinski T, Niazi U, Reis R, Trieu T J, Danovi D, Gentleman E, Vallier L, Curtis M A, Lord G M and Neves J F (2022) Organoids capture tissue-specific innate lymphoid cell development in mice and humans. *Cell Reports* **40**, 111281. doi:10.1016/j.celrep.2022.111281
- Jung P., Sato T, Merlos-Suárez A, Barriga F M, Iglesias M, Rossell D, Auer H, Gallardo M, Blasco M A, Sancho E, Clevers H and Batlle E (2011) Isolation and in vitro expansion of human colonic stem cells. *Nature Medicine* **17**(10), 1225–1227. doi:10.1038/nm.2470
- Karo-Atar D., Ouladan S, Javkar T, Joumier L, Matheson M K, Merritt S, Westfall S, Rochette A, Gentile M E, Fontes G, Fonseca G J, Parisien M, Diatchenko L, von Moltke J, Malleshaiah M, Gregorieff A and King I L (2022) Helminth-induced reprogramming of the stem cell compartment inhibits type 2 immunity. *Journal of Experimental Medicine* **219**, e20212311. doi:10.1084/jem.20212311
- Kawasaki M., Goyama T, Tachibana Y, Nagao I and Ambrosini Y M (2022) Farm and Companion Animal Organoid Models in Translational Research: A Powerful Tool to Bridge the Gap Between Mice and Humans. *Frontiers in Medical Technology* **12**(4), 895379. (PMID: 35647577; PMCID: PMC9133531). doi:10.3389/fmedt.2022.895379
- Keir M., YI Y, LU T and Ghilardi N (2020) The role of IL-22 in intestinal health and disease. *Journal of Experimental Medicine* **217**(3), e20192195. (PMID: 32997932; PMCID: PMC7062536). doi:10.1084/jem.20192195
- Klementowicz J. E., Travis M A and Grecis R K (2012) *Trichuris muris*: A model of gastrointestinal parasite infection. *Seminars Immunopathology* **34**, 815–828. doi:10.1007/s00281-012-0348-2
- Kromann E. H., Cearra A P and Neves J F (2024) Organoids as a tool to study homeostatic and pathological immune-epithelial interactions in the gut. *Clinical and Experimental Immunology* **218**(1), 28–39. doi:10.1093/cei/uxad118
- Lancaster M. A. and Knoblich J A (2014) Organogenesis in a dish: Modeling development and disease using organoid technologies. *Science* **345**, 1247125. doi:10.1126/science.1247125
- Lindemans C. A., Calafiore M, Mertelsmann A M, O’connor M H, Dudakov J A, Jenq R R, Velardi E, Young I F, Smith O M, Lawrence G, Ivanov J A, Fu Y Y, Takashima S, Hua G, Martin M L, O’rourke K P, Lo Y H, Mokry M, Romera-Hernandez M, Cupedo T, Dow L, Nieuwenhuis E E, Shroyer N F, Liu C, Kolesnick R, van den Brink M R M and Hanash A M (2015) Interleukin-22 promotes intestinal-stem-cell-mediated epithelial regeneration. *Nature* **528**(7583), 560–564. doi:10.1038/nature16460
- Lindholm H. T., Parmar N, Drurey C, Campillo Poveda M, Vornwald P M, Ostrop J, Díez-Sánchez A, Maizels R M and Oudhoff M J (2022) BMP signaling in the intestinal epithelium drives a critical feedback loop to restrain IL-13-driven tuft cell hyperplasia. *Science Immunology* **7**, eabl6543. doi:10.1126/sciimmunol.abl6543
- Liu Y., Tan J, Zhang N, QU Z, LI W, WU Y, Yin H, Liu G and FU B (2024) *Trichinella spiralis* excretory/secretory antigens ameliorate porcine epidemic diarrhea virus-induced mucosal damage in porcine intestinal organoids by alleviating inflammation and promoting tight junction. *International Journal Parasitology* **55**, 183–195. doi:10.1016/j.ijpara.2024.12.002
- Maizels R. M. and Gause W C (2023) Targeting helminths: The expanding world of type 2 immune effector mechanisms. *Journal of Experimental Medicine* **220**, e20221381. doi:10.1084/jem.20221381
- Maizels R. M., Smits H H and McSorley H J (2018) Modulation of Host Immunity by Helminths: The Expanding Repertoire of Parasite Effector Molecules. *Immunity* **49**, 801–818. doi:10.1016/j.immuni.2018.1.0.016
- Möller K. J., Wegner L H M, Malsy J, Baumdick M E, Borggrewe M, Jordan-Paiz A, Jung J M, Martrus G, Kretschmer P, Sagebiel A F, Schreurs R R C E, Hagen S H, Burmester G, Clauditz T S, Pals S T, Boettcher M, Mellinger N, Sauter G, Tomuschat C, Königs I, Schumacher U, Altfeld M, Bernink J H, Perez D, Reinshagen K and Bunders M J (2023) Expanded ILC2s in

- human infant intestines promote tissue growth. *Mucosal Immunology* 16(4), 408–421. doi:10.1016/j.mucimm.2023.04.004
- Molofsky A. B. and Locksley R M (2023) The ins and outs of innate and adaptive type 2 immunity. *Immunity* 56, 704–722. doi:10.1016/j.immuni.2023.03.014
- Moriyama S., Brestoff J R, Flamar A L, Moeller J B, Klose C S N, Rankin L C, Yudanin N A, Monticelli L A, Putzel G G, Rodewald H R and Artis D (2018) β 2-adrenergic receptor-mediated negative regulation of group 2 innate lymphoid cell responses. *Science* 359(6379), 1056–1061. doi:10.1126/science.aan4829
- Moro K., Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, Furusawa J, Ohtani M, Fujii H and Koyasu S (2010) Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature* 463(7280), 540–544. doi:10.1038/nature08636
- Mukherjee P., Roy S, Ghosh D and Nandi S K (2022) Role of animal models in biomedical research: A review. *Laboratory Animal Research* 38, 18. doi:10.1186/s42826-022-00128-1
- Neill D. R., Wong S H, Bellosi A, Flynn R J, Daly M, Langford T K, Bucks C, Kane C M, Fallon P G, Pannell R, Jolin H E and McKenzie A N (2010) Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature* 464(7293), 1367–1370. doi:10.1038/nature08900
- Nielsen M. K., (2022) Anthelmintic resistance in equine nematodes: Current status and emerging trends. *International Journal Parasitology Drugs and Drug Resistance* 20, 76–88. doi:10.1016/j.ijpddr.2022.10.005
- Nussbaum J. C., van Dyken S J, von Moltke J, Cheng L E, Mohapatra A, Molofsky A B, Thornton E E, Krummel M F, Chawla A, Liang H E and Locksley R M (2013) Type 2 innate lymphoid cells control eosinophil homeostasis. *Nature* 502(7470), 245–248. doi:10.1038/nature12526
- Nusse Y. M., Savage A K, Marangoni P, Rosendahl-Huber A K M, Landman T A, de Sauvage F J, Locksley R M and Klein O D (2018) Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche. *Nature* 559(7712), 109–113. doi:10.1038/s41586-018-0257-1
- Ogawa I., Nakai T, Iwao T and Matsunaga T (2025) Air-liquid interface culture combined with differentiation factors reproducing intestinal cell structure formation in vitro. *Biology Open* 14, bio061612. doi:10.1242/bio.061612
- Osborn M., Soares D C, Vacca F, Cohen E S, Scott I C, Gregory W F, Smyth D J, Toivakka M, Kemter A M, le Bihan T, Wear M, Hoving D, Filbey K J, Hewitson J P, Henderson H, González-Ciscar A, Errington C, Vermeren S, Astier A L, Wallace W A, Schwarze J, Ivens A C, Maizels R M and McSorley H J (2017) HpARI Protein Secreted by a Helminth Parasite Suppresses Interleukin-33. *Immunity* 47(4), 739–751.e5. doi:10.1016/j.immuni.2017.09.015
- PauZuolis M., Samperio Ventayol P, Neyazi M and Bartfeld S (2024) Organoids as a tool to study the impact of heterogeneity in gastrointestinal epithelium on host-pathogen interactions. *Clinical and Experimental Immunology* 218(1), 16–27. doi:10.1093/cei/uxae002
- Perez M. G., Gillan V, Anderson W A, Gerbe F, Herbert F, McNeilly T N, Maizels R M, Jay P, Devaney E and Britton C (2025) A secreted helminth microRNA suppresses gastrointestinal cell differentiation required for innate immunity. *Frontiers in Immunology* 16, 1558132. doi:10.3389/fimmu.2025.1558132
- Perrone F. and Zilbauer M (2021) Biobanking of human gut organoids for translational research. *Experimental and Molecular Medicine* 53(10), 1451–1458. doi:10.1038/s12276-021-00606-x
- Perusina Lanfranca M., Lin Y, Fang J, Zou W and Frankel T (2016) Biological and pathological activities of interleukin-22. *Journal of Molecular Medicine* 94, 523–534. doi:10.1007/s00109-016-1391-6
- Price A. E., Liang H E, Sullivan B M, Reinhardt R L, Easley C J, Erle D J and Locksley R M (2010) Systemically dispersed innate IL-13-expressing cells in type 2 immunity. *Proceedings of National Academy of Sciences Usa* 107(25), 11489–11494. doi:10.1073/pnas.1003988107
- Price A. E., Shamardani K, Lugo K A, Deguine J, Roberts A W, Lee B L and Barton G M (2018) A Map of Toll-like Receptor Expression in the Intestinal Epithelium Reveals Distinct Spatial, Cell Type-Specific, and Temporal Patterns. *Immunity* 49, 560–575.e6. doi:10.1016/j.immuni.2018.07.016
- PU Q., Lin P, Gao P, Wang Z, Guo K, Qin S, Zhou C, Wang B, WU E, Khan N, Xia Z, Wei X and WU M (2021) Gut Microbiota Regulate Gut-Lung Axis Inflammatory Responses by Mediating ILC2 Compartmental Migration. *Journal of Immunology* 207(1), 257–267. doi:10.4049/jimmunol.2001304
- Qin M., Fang Y, Zheng Q, Peng M, Wang L, Sang X and Cao G (2024) Tissue microenvironment induces tissue specificity of ILC2. *Cell Death Discovery* 10, 324. doi:10.1038/s41420-024-02096-y
- Quintard C., Tubbs E, Jonsson G, Jiao J, Wang J, Werschler N, Laporte C, Pitaval A, Bah T S, Pomeranz G, Bissardon C, Kaal J, Leopoldi A, Long D A, Blandin P, Achard J L, Battail C, Hagelkruys A, Navarro F, Fouillet Y, Penninger J M and Gidrol X (2024) A microfluidic platform integrating functional vascularized organoids-on-chip. *Nature Communications* 15(1), 1452. doi:10.1038/s41467-024-45710-4
- Read A. F. and Skorping A (1995) The evolution of tissue migration by parasitic nematode larvae. *Parasitology* 111(3), 359–371. doi:10.1017/s0031182000081919
- Read E., Jowett G M, Coman D and Neves J F (2022) Co-Culture of Murine Small Intestine Epithelial Organoids with Innate Lymphoid Cells. *Journal of Visualized Experiments* Mar 23 2022(181), e63554. doi:10.3791/63554
- Read E., Peña-Cearra A, Coman D, Jowett G M, Chung M W H, Coales I, Syntaka S, Finlay R E, Tachó-Piñot R, van der Post S, Naizi U, Roberts L B, Hepworth M R, Curtis M A and Neves J F (2024) Bi-directional signaling between the intestinal epithelium and type-3 innate lymphoid cells regulates secretory dynamics and interleukin-22. *Mucosal Immunology* 17(1), 1–12. doi:10.1016/j.mucimm.2023.11.002
- Roberts L. B., Schnoeller C, Berkachy R, Darby M, Pillaye J, Oudhoff M J, Parmar N, Mackowiak C, Sedda D, Quesniaux V, Ryffel B, Vaux R, Gounaris K, Berrard S, Withers D R, Horsnell W G C and Selkirk M E (2021) Acetylcholine production by group 2 innate lymphoid cells promotes mucosal immunity to helminths. *Science Immunology* 6(57), eabd0359. doi:10.1126/sciimmunol.abd0359
- Ryan S. M., Eichenberger R M, Ruscher R, Giacomini P R and Loukas A (2020) Harnessing helminth-driven immunoregulation in the search for novel therapeutic modalities. *PLoS Pathogens* 16(5), e1008508. doi:10.1371/journal.ppat.1008508
- Saenz S. A., Siracusa M C, Perrigou J G, Spencer S P, JF U, Jr, Tocker J E, Budelsky A L, Kleinschek M A, Kastelein R A, Kambayashi T, Bhandoola A and Artis D (2010) IL25 elicits a multipotent progenitor cell population that promotes T(H)2 cytokine responses. *Nature* 464(7293), 1362–1366. doi:10.1038/nature08901
- Sanderson B. E., (1969) Acetylcholinesterase activity in *Nippostrongylus brasiliensis* (Nematoda). *Comparative Biochemical Physiology* 29(3), 1207–1213. doi:10.1016/0010-406x(69)91024-x
- Sato T. and Clevers H (2015) SnapShot: Growing Organoids from Stem Cells. *Cell* 161(7), 1700–1700.e1. doi:10.1016/j.cell.2015.06.028
- Sato T., Vries R G, Snippert H J, van de Wetering M, Barker N, Stange D E and Clevers H (2009) Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* 459, 262–265. doi:10.1038/nature07935
- Schlaermann P., Toelle B, Berger H, Schmidt S C, Glanemann M, Ordemann J, Bartfeld S, Mollenkopf H J and Meyer T F (2016) A novel human gastric primary cell culture system for modelling *Helicobacter pylori* infection in vitro. *Gut* 65, 202–213. doi:10.1136/gutjnl-2014-307949
- Schneider C., O'leary C E, von Moltke J, Liang H E, Ang Q Y, Turnbaugh P J, Radhakrishnan S, Pellizzon M, MA A, & Locksley R M (2018) A Metabolite-Triggered Tuft Cell-ILc2 Circuit Drives Small Intestinal Remodeling. *Cell* 174(2), 271–284.e14. doi:10.1016/j.cell.2018.05.014
- Selkirk M. E., Lazari O, Hussein A S and Matthews J B (2005) Nematode acetylcholinesterases are encoded by multiple genes and perform non-overlapping functions. *Chemico-Biological Interactions* 157–158, 263–268. doi:10.1016/j.cbi.2005.10.039
- Serván A., Garimano N and Santini M S (2024) Systematic review and meta-analysis of soil-transmitted helminth infections in South America (2000–2024). *Acta Tropica* 260, 107400. doi:10.1016/j.actatropica.2024.107400
- Smith D., Price D R G, Burrells A, Faber M N, Hildersley K A, Chintoon-Uta C, Chapuis A F, Stevens M, Stevenson K, Burgess S T G (2021) The

- development of ovine gastric and intestinal organoids for studying ruminant host-pathogen interactions. *Frontiers Cellular Infection Microbiology* 11, 733811. doi:10.3389/fcimb.2021.733811.
- Soichot J., Guttman N, Rehrauer H, Joller N and Tritten L (2022) Nematode micrornas can Individually Regulate Interferon Regulatory Factor 4 and mtor in Differentiating T Helper 2 Lymphocytes and Modulate Cytokine Production in Macrophages. *Frontiers Molecular Bioscience* 9, 909312. doi:10.3389/fmolb.2022.909312
- Strydom T., Lavan R P, Torres S and Heaney K (2023) The Economic Impact of Parasitism from Nematodes, Trematodes and Ticks on Beef Cattle Production. *Animals (Basel)* 13, 1599. doi:10.3390/ani13101599
- Sun D., Evans L, Perrone F, Sokleva V, Lim K, Rezakhani S, Lutolf M, Zilbauer M and Rawlins EL (2021) A functional genetic toolbox for human tissue-derived organoids. *eLife* 10, e67886 doi:10.7554/eLife.67886.
- Tang X. Y., WU S, Wang D, Chu C, Hong Y, Tao M, HU H, XU M, Guo X and Liu Y (2022) Human organoids in basic research and clinical applications. *Signal Transduction and Targeted Therapy* 7(1), 168. (PMID: 35610212; PMCID: PMC9127490). doi:10.1038/s41392-022-01024-9
- Turner J. E., Morrison P J, Wilhelm C, Wilson M, Ahlfors H, Renaud J C, Panzer U, Helmby H and Stockinger B (2013) IL-9-mediated survival of type 2 innate lymphoid cells promotes damage control in helminth-induced lung inflammation. *Journal of Experimental Medicine* 210(13), 2951–2965. doi:10.1084/jem.20130071
- Vacca F, Chauché C, Jamwal A, Hinchy E C, Heieis G, Webster H, Ogunkanbi A, Sekne Z, Gregory W F, Wear M, Perona-Wright G, Higgins M K, Nys J A, Cohen E S and McSorley H J (2020) A helminth-derived suppressor of ST2 blocks allergic responses. *Elife* 9(e54017). doi:10.7554/eLife.54017
- Vande Velde F., Charlier J and Claerebout E (2018) Farmer Behavior and Gastrointestinal Nematodes in Ruminant Livestock - Uptake of Sustainable Control Approaches. *Frontiers in Veterinary Science* 5, 255. doi:10.3389/fvets.2018.00255
- Vegvari C., Giardina F, Malizia V, de Vlas S J, Coffeng L E and Anderson R M (2021) Impact of Key Assumptions About the Population Biology of Soil-Transmitted Helminths on the Sustainable Control of Morbidity. *Clinical Infectious Diseases* 72, S188–S194. doi:10.1093/cid/ciab195
- Veiga-Fernandes H. and Artis D (2018) Neuronal-immune system cross-talk in homeostasis. *Science* 359(6383), 1465–1466. doi:10.1126/science.aap9598
- Vitkauskaitė A., McDermott E, Lalor R, de Marco Verissimo C, Dehkordi M H, Thompson K, Owens P, Fearnhead H O, Dalton J P and Calvani N E D (2025) In vitro co-culture of *Fasciola hepatica* newly excysted juveniles (NEJs) with 3D HepG2 spheroids permits novel investigation of host-parasite interactions. *Virulence* 16(1). doi:10.1080/21505594.2025.2482159
- Vivier E., Artis D, Colonna M, Diefenbach A, Di Santo J P, Eberl G., Koyasu S., Locksley R.M., McKenzie A.N., Mebius R.E. and Powrie F. (2018) Innate lymphoid cells: 10 years on. *Cell* 174, 1054–1066.
- von Moltke J., Ji M, Liang H E and Locksley R M (2016) Tuft-Cell-Derived IL-25 Regulates an Intestinal ILC2-Epithelial Response Circuit. *Nature* 529, 221–225. doi:10.1038/nature16161
- Waddell A., Vallance J E, Hummel A, Alenghat T and Rosen M J (2019) IL-33 Induces Murine Intestinal Goblet Cell Differentiation Indirectly via Innate Lymphoid Cell IL-13 Secretion. *Journal of Immunology* 202(2), 598–607. doi:10.4049/jimmunol.1800292
- Wadosky K. M., Wang Y, Zhang X and Goodrich D W (2022) Generation of tumour organoids from genetically engineered mouse models of prostate cancer. *Journal of Visualized Experiments* 148, 10.3791/59710. doi:10.3791/59710
- Wang Y., DiSalvo M, Gunasekara D B, Dutton J, Proctor A, Lebhar M S, Williamson I A, Speer J, Howard R L, Smiddy N M, Bultman S J, Sims C E, Magness S T and Allbritton N L (2017) Self-renewing Monolayer of Primary Colonic or Rectal Epithelial Cells. *Cellular and Molecular Gastroenterology and Hepatology* 4(1), 165–182.e7. (PMID: 29204504; PMCID: PMC5710741). doi:10.1016/j.jcmgh.2017.02.011
- White R., Bermúdez-Barrientos J R, Robertson E, Duque Correa M A and Buck A H (2024) 2-D organoids demonstrate specificity in the interactions of parasitic nematodes and their secreted products at the basal or apical intestinal epithelium. *bioRxiv* 12(16), 628471. doi:10.1101/2024.12.16.628471
- Williamson G. L. M., Piedrahita I, JA B A T and Magness S T (2013) Cell lineage identification and stem cell culture in a porcine model for the study of intestinal epithelial regeneration. *PLoS One* 8(6), e66465. doi:10.1371/journal.pone.0066465
- World Health Organization (2024) Soil-transmitted helminth infections. Available at: <https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections> (Accessed December 2024).
- Wang Z. (2024) Unveiling the potential: Patient-derived organoids in cancer research and treatment. *Oral Oncology Reports* 9, 100240. doi:10.1016/j.oor.2024.100240.
- XU H., Wang Y, Wang W, FU Y X, Qiu J, Shi Y, Yuan L, Dong C, HU X, Chen Y G and Guo X (2025) ILC3s promote intestinal tuft cell hyperplasia and anthelmintic immunity through RANK signaling. *Science Immunology* 10(107), eadn1491. (Epub 2025 May 16. PMID: 40378237). doi:10.1126/sciimmunol.adn1491
- Yui S., Nakamura T, Sato T, Nemoto Y, Mizutani T, Zheng X, Ichinose S, Nagaishi T, Okamoto R, Tsuchiya K, Clevers H and Watanabe M (2012) Functional engraftment of colon epithelium expanded in vitro from a single adult Lgr5(+) stem cell. *Nature Medicine* 18(4), 618–623. doi:10.1038/nm.2695
- Zhang W., Chen X, Kato Y, Evans P M, Yuan S, Yang J, Rychahou P G, Yang V W, He X, Evers B M and Liu C (2006) Novel cross talk of Kruppel-like factor 4 and beta-catenin regulates normal intestinal homeostasis and tumor repression. *Molecular and Cellular Biology* 26, 2055–2064. doi:10.1128/MCB.26.6.2055-2064.2006