

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. *Tables* Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLES

Performance anxiety in professional musicians: a systematic review on prevalence, risk factors and clinical treatment effects

Fernholz I, Mumm JLM, Plag J, Noeres K, Rotter G, Willich SN, Ströhle A, Berghöfer A & Schmidt A 2287

An update on the efficacy of anti-inflammatory agents for patients with schizophrenia: a meta-analysis

Çakici N, van Beveren NJM, Judge-Hundal G, Koola MM & Sommer IEC 2307

ORIGINAL ARTICLES

Principal component analysis and brain-based predictors of emotion regulation in anxiety and depression

Klumpp H, Kinney KL, Bhaumik R & Fitzgerald JM 2320

A machine learning ensemble to predict treatment outcomes following an Internet intervention for depression

Pearson R, Pisner D, Meyer B, Shumake J & Beevers CG 2330

Auditory and visual hallucination prevalence in Parkinson's disease and dementia with Lewy bodies: a systematic review and meta-analysis

Eversfield CL & Orton LD 2342

Risk of schizophrenia, schizoaffective, and bipolar disorders by migrant status, region of origin, and age-at-migration: a national cohort study of 1.8 million people

Dykxhoorn J, Hollander A-C, Lewis G, Magnusson C, Dalman C & Kirkbride JB 2354

HOMA-IR increase after antidepressant treatment in depressed patients with the Met allele of the Val66Met BDNF genetic polymorphism

Martin S, Colle R, El Asmar K, Rigal A, Vievard A, Feve B, Becquemont L, Verstuyft C & Corruble E 2364

Pre-loss personal factors and prolonged grief disorder in bereaved mothers

Goldstein RD, Petty CR, Morris SE, Human M, Odendaal H, Elliott A, Tobacco D, Angal J, Brink L, Kinney HC, Prigerson HG & for the PASS Network 2370

Multidimensional understanding of religiosity/spirituality: relationship to major depression and familial risk

McClintock CH, Anderson M, Svob C, Wickramaratne P, Neugebauer R, Miller L & Weissman MM 2379

Identifying key targets for interventions to improve psychological wellbeing: replicable results from four UK cohorts

Stochl J, Sonesson E, Wagner AP, Khandaker GM, Goodyer I & Jones PB 2389

Risk and coaggregation of major psychiatric disorders among first-degree relatives of patients with bipolar disorder: a nationwide population-based study

Chen M-H, Hsu J-W, Huang K-L, Su T-P, Li C-T, Lin W-C, Tsai S-J, Cheng C-M, Chang W-H, Pan T-L, Chen T-J & Bai Y-M 2397

Prediction of remission in pharmacotherapy of untreated major depression: development and validation of multivariable prediction models

Furukawa TA, Kato T, Shinagawa Y, Miki K, Fujita H, Tsujino N, Kondo M, Inagaki M & Yamada M 2405

Optimizing patient expectancy in the pharmacologic treatment of major depressive disorder

Zilcha-Mano S, Brown PJ, Roose SP, Cappetta K & Rutherford BR 2414

Childhood determinants of suicidality: comparing males in military and civilian employed populations

Syed Sheriff R, Van Hooff M, Malhi G, Grace B & McFarlane A 2421

Bias-contingent attention bias modification and attention control training in treatment of PTSD: a randomized control trial

Lazarov A, Suarez-Jimenez B, Abend R, Naim R, Shvil E, Helpman L, Zhu X, Papini S, Duroski A, Rom R, Schneier FR, Pine DS, Bar-Haim Y & Neria Y 2432

From laboratory to life: associating brain reward processing with real-life motivated behaviour and symptoms of depression in non-help-seeking young adults

Bakker JM, Goossens L, Kumar P, Lange IMJ, Michielse S, Schruers K, Bastiaansen JA, Lieveer R, Marcellis M, van Amelsvoort T, van Os J, Myin-Germeys I, Pizzagalli DA & Wichers M 2441

Hippocampal subfields and visuospatial associative memory across stages of schizophrenia-spectrum disorder

Wannan CMJ, Cropley VL, Chakravarty MM, Van Rheenen TE, Mancuso S, Bousman C, Everall I, McGorry PD, Pantelis C & Bartholomeusz CF 2452