

## IMPLICATIONS OF ILLITE/SMECTITE STABILITY DIAGRAMS: A DISCUSSION

**Key Words**—Chemical activity, Illite, Mixed layer, Phase equilibria, Smectite, Stability.

### INTRODUCTION

Although all interstratified illite/smectites contain some Mg and/or Fe, changes in the site occupancies of these cations are not required in the conversion of interlayers from smectite to illite (Garrels, 1984). Thus, to a first approximation, illite/smectite phase relationships can be represented in the simplified system  $K_2O-Al_2O_3-SiO_2-H_2O$ . On this premise, Garrels (1984) recently constructed chemical activity diagrams of the system  $K_2O-Al_2O_3-SiO_2-H_2O$  to compare phase relationships expected if illite/smectite (I/S) behaves as a solid solution (single phase) or as a mixture of two phases.

Analyses of natural aqueous solutions in contact with clays (Aagaard and Helgeson, 1983) were found to lie close to the phase boundary representing I/S (two phase) equilibrium, assuming arbitrary but reasonable illite and smectite compositions, whereas these analyses cross the field of I/S solid solution (one phase). It is difficult, however, to determine whether these aqueous solutions are actually in equilibrium with illite, because no compositions are available for natural solutions known to coexist with illite (Aagaard and Helgeson, 1983). In the absence of conclusive data, Garrels (1984) reasoned that the available data are more nearly consistent with the interpretation that I/S is a two-phase assemblage rather than a single-phase solid solution. Inasmuch as the evidence available to Garrels was not definitive, he suggested criteria for distinguishing between these two hypotheses and proposed long-term experiments to resolve the problem.

If I/S behaves as two phases, equilibration with kaolinite should result in isothermal, isobaric invariance, thus fixing solution composition at the kaolinite-illite-smectite triple point (Garrels, 1984, Figure 4). On the other hand, if I/S behaves as a solid solution, equilibration with kaolinite would not result in isothermal, isobaric invariance. A range of solution compositions ( $a_{K^+}/a_{H^+}$ ,  $a_{SiO_2}$ ) could, therefore, coexist at equilibrium with kaolinite and I/S solid solution (Garrels, 1984, Figure 3). Garrels' analysis is particularly timely because the critical data necessary to test these hypotheses concerning the nature of illite have recently become available.

### EXPERIMENTAL DATA AND THEIR IMPLICATIONS

Kittrick (1984) equilibrated Goose Lake illite with kaolinite in long-term solubility experiments. Equilib-

rium was indicated by constant solution compositions attained from undersaturation and supersaturation in experiments of long duration. Solubility measurements were made at intervals terminating with solution samples taken after more than 300 and more than 500 days. The final analyses for each experiment are shown in Figure 1 (circles). Sass (1984) equilibrated Goose Lake illite with 0.2 M and 2.0 M KCl/HCl solutions in the presence of kaolinite and quartz. Equilibrium was approached from two directions by starting with solutions having high and low  $a_{K^+}/a_{H^+}$  ratios. Each data point (triangles) in Figure 1 thus represents a reversal pair.

Data for both sets of experiments appear to lie close to a single straight line on a  $\log a_{K^+}/a_{H^+}$  vs.  $\log a_{SiO_2}$  diagram (Figure 1). Linear regression analysis was used to determine the correlation coefficient ( $-0.95$ ) and the equation of the line; its slope,  $-6.09$ , is directly related to the composition of the phase controlling the solubility. Calculations following the method of Mattigod and Kittrick (1979) indicate that the composition of the solubility controlling phase is approximately  $K_{0.25}Al_{2.00}(Si_{3.75}Al_{0.25})O_{10}(OH)_2$ . The close accord of solution compositions obtained in experiments with 0.2 M and 2 M solutions (Sass, 1984) suggests that  $K^+$  activity is solubility controlled. Thus,  $K^+$  behaves as a common ion and an integral part of the structure of this phase.

The mineralogical term applied to a phase of this composition is a matter of definition. The boundary between illite and smectite is customarily placed at  $-0.4$  charge units per half cell (Hower and Mowatt, 1966). A recent report of The Clay Minerals Society Nomenclature Committee (Bailey *et al.*, 1984) and S. W. Bailey (University of Wisconsin, Madison, Wisconsin, personal communication, 1985), however, suggest that the boundary should be defined by expandability, illite being a nonexpanding clay mineral. The minimum interlayer charge for illite by the latter definition is about  $-0.6$ , whereas the maximum interlayer charge for smectite is less than  $-0.4$ . Goose Lake illite has the formula  $K_{0.59}(Al_{1.58}Fe_{0.24}Mg_{0.15})(Si_{3.64}Al_{0.35})O_{10}(OH)_2$  (Kittrick, 1984) and is 25–30% expandable (Routson, 1970, p. 79). Thus, Goose Lake illite is actually an illite/smectite, while the solubility-controlling phase is a smectite.

### DISCUSSION

If illite is considered to be a single-phase solid solution, the kaolinite-illite boundary (Garrels, 1984,




Figure 1. Compositions of solutions equilibrated with Goose Lake illite + kaolinite; triangles, Sass (1984); circles, Kittrick (1984). Smectite (I/S), smectite component of I/S.

Figure 3) should be a curve corresponding to progressively lower dissolved-silica activities with increasing K-content of illite. Garrels (1984) proposed that this range of silica values indicates that illite is a solid solution, whereas fixed values of silica and  $a_{K^+}/a_{H^+}$  at the triple point (Garrels, 1984, Figure 4) indicates that illite is a two-phase assemblage.

The data points in Figure 1 define a univariant solubility line representing a single "illite" composition which appears to control solubility over a large range of  $a_{K^+}/a_{H^+}$  values. Inasmuch as the solubility-controlling phase has a composition in the range associated with smectites and the bulk chemical composition of Goose Lake illite cannot be represented by this composition alone, this phase may represent the smectite component of I/S. Thus, assuming congruent solubility, Goose Lake illite appears to behave as a two-phase assemblage.

Although equilibrium with respect to the illite component of I/S has not been attained, kaolinite-smectite equilibrium may be used to deduce illite phase relationships qualitatively. The presence of an illite-smectite boundary and a triple point can be inferred on this basis, even though direct experimental evidence is lacking. Consequently, the portion of the solubility line (Figure 1) at values of  $a_{K^+}/a_{H^+}$  greater than the triple point represent metastable kaolinite-smectite equilibrium.

Because similar experimental results have been obtained for illites from other localities (Kittrick, 1984; Sass, 1984), inferences for Goose Lake illite may be generally applicable to I/S random. Thus, recent evidence strongly suggests that I/S random behaves as a two-phase assemblage (Kittrick, 1984; Sass and Rosenberg, 1984; Nadeau *et al.*, 1984) as inferred by Garrels (1984).

#### ACKNOWLEDGMENTS

We thank R. M. Garrels for making his calculations available to us and for review of the manuscript. Computer analysis and graphics were provided by Mrs. Monica Frye. Acknowledgment is also made to the donors of the Petroleum Research Fund, administered by the American Chemical Society for support of this research.

Department of Geology P. E. ROSENBERG  
Department of Agronomy and Soils J. A. KITTRICK  
Washington State University  
Pullman, Washington 99164

Geoscience Research and Engineering B. M. SASS  
Department  
Battelle, Pacific Northwest Laboratories  
Richland, Washington 99352

#### REFERENCES

- Aagaard, P. and Helgeson, H. C. (1983) Activity/composition relations among silicates and aqueous solutions: II. Chemical and thermodynamic consequences of ideal mixing of atoms on homologous sites in montmorillonites, illites and mixed-layer clays: *Clays & Clay Minerals* **31**, 207-217.
- Bailey, S. W., Brindley, G. W., Fanning, D. S., Kodama, H., and Martin, R. T. (1984) Report of the Clay Mineral Society Nomenclature Committee for 1982 and 1983: *Clays & Clay Minerals* **32**, 239-240.
- Garrels, R. M. (1984) Montmorillonite/illite stability diagrams: *Clays & Clay Minerals* **32**, 161-166.
- Hower, J. and Mowatt, T. C. (1966) The mineralogy of illites and mixed-layer illite/montmorillonites: *Amer. Mineral.* **51**, 825-854.
- Kittrick, J. A. (1984) Solubility measurements of phases in three illites: *Clays & Clay Minerals* **32**, 115-124.
- Mattigod, S. V. and Kittrick, J. A. (1979) Aqueous solubility studies of muscovites: apparent nonstoichiometric solute activities at equilibrium: *Soil Sci. Soc. Amer. J.* **43**, 180-187.
- Nadeau, P. H., Wilson, M. J., McHardy, W. J., and Tait, J. M. (1984) Interstratified clays as fundamental particles: *Science* **225**, 923-925.
- Routson, R. C. (1970) Illite solubility: Ph.D. thesis, Washington State University, Pullman, Washington, 123 pp.
- Sass, B. M. (1984) Stability relationships of illite in solutions between 25° and 250°C: M.S. thesis, Washington State University, Pullman, Washington, 131 pp.
- Sass, B. M. and Rosenberg, P. E. (1984) Multi-phase solubility in natural illites between 25° and 250°C: *Program and Abstracts*, Annual Meeting, Clay Minerals Society, Baton Rouge, Louisiana, p. 101.

(Received 15 January 1985; accepted 24 June 1985; Ms. 1443)