



PAPER

Non-linear biphasic mixture model: Existence and uniqueness results

Meraj Alam¹, Adrian Muntean² and Raja Sekhar G P³

¹Department of Mathematics, Ecole Centrale School of Engineering, Mahindra University, Hyderabad, 500043, Telangana, India

²Department of Mathematics and Computer Science, Karlstad University, Karlstad, 651 88, Sweden

³Department of Mathematics, Indian Institute of Technology Kharagpur, Kharagpur, West Bengal, 721302, India

Corresponding author: Raja Sekhar G P; Email: rajas@iitkgp.ac.in

Received: 10 February 2022; **Revised:** 09 August 2023; **Accepted:** 07 May 2024; **First published online:** 18 September 2024

Keywords: in-vitro tumour; biphasic mixture theory; hydraulic resistivity; weak solutions; fixed-point theory

2020 Mathematics Subject Classification: 76Txx, 76Zxx (Primary); 35Q74, 35D30 (Secondary)

Abstract

This paper is concerned with the development and analysis of a mathematical model that is motivated by interstitial hydrodynamics and tissue deformation mechanics (poro-elasto-hydrodynamics) within an in-vitro solid tumour. The classical mixture theory is adopted for mass and momentum balance equations for a two-phase system. A main contribution of this study is we treat the physiological transport parameter (i.e., hydraulic resistivity) as anisotropic and heterogeneous, thus the governing system is strongly coupled and non-linear. We derived a weak formulation and then formulated the equivalent fixed-point problem. This enabled us to use the Galerkin method, and the classical results on monotone operators combined with the well-known Schauder and Banach fixed-point theorems to prove the existence and uniqueness of results.

1. Introduction

The study of fluid flow through porous media has gained the attention of many researchers over the years. Examples of natural porous materials are living tissues, rocks, soils, etc. On the other hand, there are manmade porous materials, e.g., concrete, foam, ceramic, etc. Because of their wide presence and hard-to-estimate effective properties, flow through porous material is studied by engineers and scientists. This leads to a coupled phenomenon where fluid flow and solid deformation in porous materials interplay. This is a classical problem in geomechanics and biomechanics. Recently, one of the most studied topics in the field of fluid mechanics is flow through biological tissues such as tumours, glycocalyx layers and articular cartilage. This paper deals with the mathematical modelling and analysis of the coupled phenomena of fluid flow and solid deformation (in short *poro-elasto-hydrodynamics*) within an in-vitro tumour model. Typically, a tumour is assumed to be a deformable porous medium that consists of multiple phases, e.g., one fluid phase and many solid ones. A tumour may exist in isolation (in-vitro) or may be present in normal tissue (in-vivo) [1]. Although the internal geometrical structure of tumours is complicated, developing mathematical models for approximate situations is very useful. Theoretical predictions generated from such approximate models may help to reduce the number of animal experiments that need to be carried out and also suggest new experimental programmes that identify optimal tumour therapy schedules [2].

Mathematical models of tumours in general can be divided into three categories: discrete, continuum and hybrid. Here, we focus on continuum models that treat cells as averaged populations and are based on the continuum mechanics approach to porous media combined with mixture theory [3]. The early

mathematical models on the avascular tumour growth [4] assumed that tumours are made of single type of cells having a constant density. However, various experimental, and theoretical evidence have shown that such a description is not sufficient to study the tumour dynamics [4, 5]. Hence, multiphase models came into play. In this case, one can consider density variations within mixture components to evaluate the evolution of partial stresses. Biot's theory of poroelasticity and the theory of mixture are commonly adapted models to explore *poro-elasto-hydrodynamics* [1]. Alike continuum-level approaches involve the development of a set of equations to represent the mechanical behaviour of a soft tissue such as a tumour (which is assumed as a deformable porous material) at the macro scale, using a porous media approach.

The first multiphase model for tumours is proposed by Please et al. [6]. The authors proposed a diffusion equation for cell concentration and generalised Darcy equation for cells and water motion inside the tumour. Further, such multiphase models have been studied analytically and numerically by several authors, one can refer [5, 7, 8]. Sumets et al. [9] described a new boundary-integral representation for biphasic mixture theory, where they solved elasto-hydrodynamic-mobility problems using boundary element methods. Dey and Raja Sekhar [1] used a biphasic mixture model for poro-elasto-hydrodynamics and nutrient transport inside an in-vitro solid tumour. The authors assumed the presence of an unknown fluid source/sink in the model and biphasic mixture equations have been solved explicitly in the case of one-dimensional spherical geometry. Slvia and Wheeler [10] presented a coupled geomechanics and unsteady reservoir flow model using the theory of poroelasticity. They established the existence and uniqueness of a weak solution and computed *a priori* error estimates for the numerical solution with stress-dependent permeability. In [11], a non-linear model for a poroelastic medium (described by quasi-static Biot-equations) coupled to transport equations of substances was considered. They have modelled time and space-dependent processes in deformable cellular tissues by the method of homogenisation, starting from a reactive flow system coupling mechanics at the pore scale. The model was analysed and the global-in-time existence and uniqueness of the solution were shown. Cao et al. [12] have considered a non-linear steady flow model in a deformable biological domain based on the theory of poroelasticity (non-linearity is due to the assumption of dilation-dependent interstitial permeability of the solid matrix). They established the existence and uniqueness of a weak solution. Looking through mentioned literature, we observe that there is a gap in dealing with the existence and uniqueness of corresponding biphasic mixture equations that describe the coupled phenomena of fluid flow and solid deformation within biomaterials such as tumours. Attempting to fill such a gap, Alam et al. [13, 14] developed a well-posedness theory and certain regularity results in $2d$ and $3d$ for poro-elasto-hydrodynamics model inside an in-vitro solid tumour. Further, in the case of an in-vivo solid tumour, Alam et al. [15] developed existence and uniqueness resulting in a weak sense for poro-elasto-hydrodynamics while assuming the hydraulic resistivity is heterogeneous and deformation dependent.

We note that, in general, poro-elasto-hydrodynamics models within a tumour may not lead directly to linear biphasic mixture equations. In practice, due to the non-uniform blood vessel distribution, the supply of fluids and macromolecules within a tumour is heterogeneous. As a consequence, physiological transport parameters (e.g., hydraulic resistivity or permeability) depend on space and deformation. For instance, in the case of soft permeable tissue and gel, Barry and Aldis [16] and Holmes and Mow [17] considered permeability depending exponentially on the strain. Also, some of the biological tissues and cells display anisotropic permeability [18]. In particular, articular cartilage typically exhibits anisotropic behaviour [19]. Further, in the case of multicellular tumour tissue, Givero and Preziosi [20] followed Holmes and Mow [17] and considered that the permeability depends on the volumetric deformation (or strain). Dey and Sekhar [1] assumed that the hydraulic conductivity of soft tumour depends on the radial distance. These non-linear effects inserted in the physiological parameter yield non-linear biphasic mixture equations. As far as we know, for non-linear biphasic mixture models, there is a lack of literature regarding the existence, uniqueness and regularity of the solution. In this paper, we present a non-linear biphasic mixture model that represents poro-elasto-hydrodynamics, which do not account for any new growth of tumour cells. The physiological transport parameter (hydraulic resistivity) is assumed to be

deformation dependent, which yields the non-linearity in the model. We develop a local weak solvability theory.

1.1 Biphasic mixture theory

In this subsection, we introduce the generic governing equations. We use biphasic mixture theory to represent the fluid and solid phases of the tumour. Following [1, 8, 21], we apply the conservation of mass and momentum to the fluid and solid phases, viewing the fluid as viscous Newtonian and the solid as deformable, and accounting for momentum exchange between the two phases. Let \mathbf{V}_f and \mathbf{V}_s denote the velocities of the fluid and solid phases, respectively. The apparent densities of the fluid and solid phases are denoted by $\tilde{\rho}_f$ and $\tilde{\rho}_s$, respectively, and their corresponding volume fractions by ϕ_f and ϕ_s . The true densities of the fluid and solid phases are then $\rho_f = \phi_f \tilde{\rho}_f$ and $\rho_s = \phi_s \tilde{\rho}_s$. Accordingly, in Ω , the mass and linear momentum balance equations for the fluid phase are given by

$$\frac{\partial(\tilde{\rho}_f \phi_f)}{\partial t} + \nabla \cdot [(\tilde{\rho}_f \phi_f) \mathbf{V}_f] = \tilde{\rho}_f S_f, \tag{1.1}$$

$$\rho_f \left(\frac{\partial \mathbf{V}_f}{\partial t} + (\mathbf{V}_f \cdot \nabla) \mathbf{V}_f \right) = \nabla \cdot \mathbf{T}_f + \mathbf{\Pi}_f + \mathbf{b}_f, \tag{1.2}$$

where \mathbf{T}_f denotes the stress tensor for the fluid phase

$$\mathbf{T}_f = -[\phi_f P - \lambda_f \nabla \cdot \mathbf{V}_f] \mathbf{I} + \mu_f [\nabla \mathbf{V}_f + (\nabla \mathbf{V}_f)^T]. \tag{1.3}$$

The corresponding mass and linear momentum equations for the solid phase are

$$\frac{\partial(\tilde{\rho}_s \phi_s)}{\partial t} + \nabla \cdot [(\tilde{\rho}_s \phi_s) \mathbf{V}_s] = \tilde{\rho}_s S_s, \tag{1.4}$$

$$\rho_s \left(\frac{\partial \mathbf{V}_s}{\partial t} + (\mathbf{V}_s \cdot \nabla) \mathbf{V}_s \right) = \nabla \cdot \mathbf{T}_s + \mathbf{\Pi}_s + \mathbf{b}_s, \tag{1.5}$$

where \mathbf{T}_s denotes the stress tensor for the solid phase

$$\mathbf{T}_s = -[\phi_s P - \chi_s (\nabla \cdot \mathbf{U}_s)] \mathbf{I} + \mu_s [\nabla \mathbf{U}_s + (\nabla \mathbf{U}_s)^T]. \tag{1.6}$$

In Equations (1.1) and (1.4), S_f and S_s are fluid and solid source terms, respectively. \mathbf{U}_s denotes the displacement of the solid phase. Hence, $\mathbf{V}_s = \frac{\partial \mathbf{U}_s}{\partial t}$. The average interstitial fluid pressure is P and \mathbf{b}_j $j = \{1, 2\}$ denotes the body force. Further, the volume fractions ϕ_f and ϕ_s are assumed to satisfy the following saturation assumption

$$\phi_f + \phi_s = 1. \tag{1.7}$$

We suppose that the two phases interact together via the drag forces $\mathbf{\Pi}_s$ and $\mathbf{\Pi}_f$, which by Newton’s third law, are equal and opposite. Following [7, 8], we define

$$-\mathbf{\Pi}_s = \mathbf{\Pi}_f = K(\mathbf{V}_s - \mathbf{V}_f) - (\nabla \phi_s) P, \tag{1.8}$$

where $K = \frac{\mu_f}{k}$ is the hydraulic resistivity or drag coefficient, where k is the permeability of the porous matrix (the precise nature of K will be defined in the next section). Furthermore, μ_f (μ_s) is the dynamic viscosity of the fluid phase (solid phase), while λ_f and χ_s denote the Lamé coefficient and shear modulus of the fluid and solid phases. The elastic modulli χ_s and μ_s are related to the Young’s modulus (\mathcal{Y}) and Poisson’s ratio (ν_p) via the relationship

$$\chi_s = \frac{\nu_p \mathcal{Y}}{(1 + \nu_p)(1 - 2\nu_p)} \text{ and } \mu_s = \frac{\mathcal{Y}}{2(1 + \nu_p)}.$$

1.2 Main modelling assumptions

Having presented a more generic set of mixture theory equations in the absence of any assumptions or boundary conditions, here we list some biologically suitable assumptions restricted to a specific tumour model. Our choice of model is motivated by the study of an isolated (in-vitro) tumour that behaves as a heterogeneous deformable porous medium. Usually, tumour tissues are considered incompressible fluids with no voids present. It is assumed that each phase has an equal constant density. Suppose $\Omega \subset \mathbb{R}^d$, $d = \{2, 3\}$ is a bounded Lipschitz domain that is filled by the tumour. Let $\partial\Omega$ be its boundary (Figure 1). One may note that the solid tumour is essentially a multicellular spheroid. When nutrients perfuse the interstitial space, a large number of cells receive adequate food for survival and proliferation. As a consequence, the tumour grows in size. For a growing tumour, the permeability and the effective mechanics parameters (χ_s and μ_s) may depend on the volume fraction of the cell population [8]. Moreover, the volume fractions depend on space. Hence, it is extremely difficult to analyse mathematically the growth model and fluid transport model simultaneously. To simplify this, we assume that the tumour tissue is not growing and all elastic parameters (χ_s and μ_s , etc.) are independent of volume fractions. Further, we make the following modelling assumptions (A1)–(A4):

- (A1) Nutrient perfusion and transport occur on much shorter timescales than the timescale for tumour cell growth. Accordingly, we view the tumour as a static, perfused biological domain. On the short timescale associated with nutrient transport to (and within) the tumour, cell death and proliferation are assumed to be negligible. Therefore, we fix $S_s = 0$ in the tumour and normal tissue regions. Further, on the timescale of interest, the solid volume fraction ϕ_s remains constant, and for simplicity, we assume it to be independent of spatial position and time as (see [1]).
- (A2) A fluid source is attached to the mixture, hence $S_f \neq 0$, [1, 22].
- (A3) Motion of the cells and interstitial fluid flow is so slow that the inertial terms can be neglected in both phases, see e.g. [8, 23].
- (A4) *Structure of the hydraulic resistivity:* Various experimental and theoretical investigations indicate clearly that for the deformable porous medium (or soft biological tissue such as articular cartilage, arterial tissue and tumour), the permeability also called in this context hydraulic resistivity depends on stress, dilatation, volume fractions (porosity), etc. There are several analytical expressions for permeability that are used in literature such as $k(x) = \exp[mU'(x)]$, where m is a constant and $U' = \frac{dU}{dx}$ with U as displacement, while $k(x) = k_0/[1 - mU'(x)]$ or $k(x) = k_0[1 + mU'(x)]$ for small $mU'(x)$, or $k(x) = k_0(\phi/\phi_0)^n$ [16, 17]. Here, k_0 is the permeability at reference porosity ϕ_0 and n is a variable that may be determined by fitting experimental data. In this framework, we propose two different cases. We assume that the hydraulic resistivity K depends explicitly on (a) the solid phase displacement \mathbf{U}_s and (b) the strain/dilatation i.e. on $\nabla \cdot \mathbf{U}_s$. In both of these cases, we admit anisotropic effects, i.e. \mathbf{K} is a square matrix of order $d = \{2, 3\}$. Note that in case (a), we do not use any specific, explicit expression of $\mathbf{K}(\mathbf{U}_s)$ in our analysis. One may think through the following choice

$$\mathbf{K}(\mathbf{U}_s) = (\alpha_1 |\mathbf{U}_s| + \alpha_3) \mathbf{I} + (\alpha_1 - \alpha_2) \frac{\mathbf{U}_s \mathbf{U}_s^T}{|\mathbf{U}_s|},$$

where α_i ($i = 1, 2, 3$) are real constants such that $\alpha_1 \geq 0$, $\alpha_2 \geq 0$ and $\alpha_3 > 0$. One can easily show that \mathbf{K} is Lipschitz and uniformly positive [24]. For case (b), one can think of a form $\mathbf{K}(\nabla \cdot \mathbf{U}_s) = (\gamma_1 + \gamma_2 |\nabla \cdot \mathbf{U}_s|) \mathbf{I}$, where $\gamma_i \geq 0$ are real constants, and \mathbf{I} is the identity matrix. One can observe that \mathbf{K} is Lipschitz and uniformly positive. We will state further assumptions on \mathbf{K} in the next Section 3, [16, 17].




Figure 1. Geometry of the problem.

The assumptions (A1)–(A4) are taken into account in Equations (1.1)–(1.8). This helps us to define a coupled non-linear system of steady-state mass and momentum balance equations in unknowns $(\mathbf{V}_f, \mathbf{U}_s, P)$ as follows:

$$-\nabla \cdot (2\mu_f D(\mathbf{V}_f) + \lambda_f (\nabla \cdot \mathbf{V}_f) \mathbf{I} - \phi_f P \mathbf{I}) + \mathbf{K}(\boldsymbol{\zeta}) \mathbf{V}_f = \mathbf{b}_f \text{ in } \Omega, \tag{1.9}$$

$$-\nabla \cdot (2\mu_s D(\mathbf{U}_s) + \lambda_s (\nabla \cdot \mathbf{U}_s) \mathbf{I} - \phi_s P \mathbf{I}) - \mathbf{K}(\boldsymbol{\zeta}) \mathbf{V}_f = \mathbf{b}_s \text{ in } \Omega, \tag{1.10}$$

$$\nabla \cdot (\phi_f \mathbf{V}_f) = S_f \text{ in } \Omega, \tag{1.11}$$

where $\boldsymbol{\zeta}$ is either $\nabla \cdot \mathbf{U}_s$ or \mathbf{U}_s . Further, $D(\cdot)$ denotes the deviatoric matrix, which is defined as $D(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})')$, where $(\nabla \mathbf{u})'$ denotes the transpose of the matrix $\nabla \mathbf{u}$. We have made the approximation $\mathbf{V}_s = \frac{\partial \mathbf{U}_s}{\partial t} \approx 0$ which is consistent with the infinitesimal strain theory that is used in e.g. [1, 25]. The mass balance equations for the fluid phase include a source term S_f , which models fluid exchange with vasculature and lymph vessels. For a closed mixture (for instance the case of avascular tumours), we consider $S_f = 0$ so that $\nabla \cdot (\phi_f \mathbf{V}_f) = 0$. This is the counterpart of the incompressibility constraint. Note that in Equation (1.11), even though the density of each phase is constant, the vector \mathbf{V}_f is not solenoidal. On the other hand, when the external sources/sinks are attached to the mixture, we have $S_f \neq 0$ [5, 22]. Typically, the fluid source S_f is assumed to be driven by the average transmural pressure and (trusting [1, 22]) takes the form

$$S_f = -L_p \left(\frac{A}{V} \right) \{1 + L_r A_r\} (P - P_F), \tag{1.12}$$

where L_p is the average hydraulic conductivity coefficient of capillary. In (1.12), A/V denotes the capillary surface area per unit tissue volume in the tumour tissue and $L_r A_r$ denotes the ratio of the strength of distributed solute source through the vasculature and solute sink through the lymph vessels and P_F is the weighted vascular pressure.

1.3 Boundary conditions

The model that we have considered here is supposed to mimic an in-vitro tumour. Accordingly, we prescribe

$$\mathbf{T}_f \cdot \mathbf{n} = \mathbf{T}_\infty \text{ and } \mathbf{U}_s = 0 \text{ on } \partial\Omega, \tag{1.13}$$

where \mathbf{n} is the outward normal unit vector to the boundary $\partial\Omega$.

2. Non-dimensional equations

Using the transformations $\hat{\mathbf{x}} = \frac{\mathbf{x}}{R}$, $\nabla' = R\nabla$, $\hat{P} = \frac{P}{P_F}$, $\hat{\mathbf{V}}_f = \frac{\mathbf{V}_f}{\frac{R P_F}{\mu_f}}$, $\hat{\mathbf{U}}_s = \frac{\mathbf{U}_s}{\frac{R^3 P_F}{\mu_f \nu}}$, $\mathbf{K} = \hat{\mathbf{K}} K_d$, where K_d is the hydraulic resistivity (drag coefficient) of the tumour tissue in the absence of deformation, R is the length

of the edge of the d -cube in which Ω is contained. The following dimensionless form of the governing Equations (1.9)–(1.11) (‘hat’ is dropped for convenience) is available in Ω

$$-\nabla \cdot \left(2D(\mathbf{V}_f) + \frac{\lambda_f}{\mu_f} (\nabla \cdot \mathbf{V}_f) \mathbf{I} - \phi_f P \mathbf{I} \right) + \frac{1}{Da} \mathbf{K}(\boldsymbol{\zeta}) \mathbf{V}_f = \mathbf{b}_f, \tag{2.1}$$

$$-\nabla \cdot \left(\frac{\varrho_t}{(1 + \nu_p)} D(\mathbf{U}_s) + \frac{\nu_p \varrho_t}{(1 + \nu_p)(1 - 2\nu_p)} (\nabla \cdot \mathbf{U}_s) \mathbf{I} - \phi_s P \mathbf{I} \right) - \frac{1}{Da} \mathbf{K}(\boldsymbol{\zeta}) \mathbf{V}_f = \mathbf{b}_s, \tag{2.2}$$

$$\nabla \cdot (\phi_f \mathbf{V}_f) = -\alpha_t^2 (1 + L_r A_r) (P - 1). \tag{2.3}$$

In (2.1) and (2.2), \mathbf{b}_f and \mathbf{b}_s are modified non-dimensional body forces, and $\varrho_t = \mathcal{Y} R^2 \rho_f / \mu_f^2$ is the dimensionless Young’s modulus Y associated with the solid phase. It contains the response of the solid phase (cellular phase + extracellular matrix) towards viscous drag due to interstitial fluid movement. $\alpha_t^2 = L_p(A/V)\mu_f$ is the strength of solute source, and $Da = \frac{K_d \mu_f}{R^2}$ is the Darcy number (permeability parameter). The corresponding boundary conditions are

$$\left(2D(\mathbf{V}_f) + \frac{\lambda_f}{\mu_f} (\nabla \cdot \mathbf{V}_f) \mathbf{I} - \phi_f P \mathbf{I} \right) \cdot \mathbf{n} = \mathbf{T}_\infty \text{ and } \mathbf{U}_s = 0 \text{ on } \partial\Omega. \tag{2.4}$$

For the sake of writing convenience, set $\lambda = \frac{\lambda_f}{\mu_f}$, $\alpha_1 = \frac{\varrho_t}{2(1+\nu_p)}$, $\alpha_2 = \frac{\nu_p \varrho_t}{(1+\nu_p)(1-2\nu_p)}$, and $a_0 = \alpha_t^2 (1 + L_r A_r)$. Observe that the system of Equations (2.1)–(2.3) is non-linear and fully coupled whenever $\boldsymbol{\zeta}$ is equals to either \mathbf{U}_s or $\nabla \cdot \mathbf{U}_s$, which is our primary interest for now. The main aim is to study the well-posedness of the non-linear system (2.1)–(2.3) subject to the data (2.4).

3. Well-posedness of the auxiliary sub-problems

$\mathbf{V}_f, P, \mathbf{U}_s$ are the unknown functions in the system of Equations (2.1)–(2.3). We assume the following:

- (A) The parameters $\phi_f > 0, \phi_s > 0, \lambda \geq 0, \alpha_1 > 0, \alpha_2 > 0, a_0 > 0, Da > 0$ are known real constants, and the functions $\mathbf{b}_j \in L^2(\Omega)^d$ where $j = f, s, \mathbf{T}_\infty \in L^2(\partial\Omega)^d$ are also known. $c_k > 0, c_p > 0, c_t > 0, c_s > 0$ are some real constants that appear in Korn’s, Poincare’s, trace and Sobolev’s inequalities, respectively.¹
- (B) Let $\mathbf{K} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a symmetric, uniformly bounded and positive definite matrix. This ensures that there exist positive constants k_1 and k_2 such that for all $\boldsymbol{\xi}, \mathbf{x} \in \mathbb{R}^d$, we have:

$$(i) k_1 \boldsymbol{\xi} \cdot \boldsymbol{\xi} \leq \mathbf{K}(\mathbf{x}) \boldsymbol{\xi} \cdot \boldsymbol{\xi} \text{ and } (ii) \|\mathbf{K}(\mathbf{x})\| \leq k_2, \tag{3.1}$$

$\|\cdot\|$ denotes the Euclidean norm.

- (C) We assume that the hydraulic resistivity \mathbf{K} is Lipschitz continuous. To this extent, let us assume that there exists a constant $k_L > 0$ such that

$$\|\mathbf{K}(\mathbf{x}) - \mathbf{K}(\mathbf{y})\| \leq k_L \|\mathbf{x} - \mathbf{y}\| \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^d. \tag{3.2}$$

3.1 Concept of weak formulation

Choose the triplet of test functions $(\mathbf{W}, \mathbf{Z}, q) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$. Taking the scalar product of (2.1) with \mathbf{W} , (2.2) with \mathbf{Z} and (2.3) with q , and using the boundary conditions (2.4), we get the following non-linear weak formulation:

¹We refer to the Appendix section for details on function spaces and other preliminary results.

Find the triplet $(\mathbf{V}_f, \mathbf{U}_s, P) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ such that

$$2(D(\mathbf{V}_f) : D(\mathbf{W}))_\Omega + \lambda(\nabla \cdot \mathbf{V}_f, \nabla \cdot \mathbf{W})_\Omega - \phi_f(P, \nabla \cdot \mathbf{W})_\Omega \tag{3.3}$$

$$+ \frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})\mathbf{V}_f, \mathbf{W})_\Omega = (\mathbf{b}_f, \mathbf{W})_\Omega + (\mathbf{T}_\infty, \mathbf{W})_{\partial\Omega}$$

$$\phi_f(\nabla \cdot \mathbf{V}_f, q)_\Omega + a_0(P, q)_\Omega = (a_0, q)_\Omega \tag{3.4}$$

$$2\alpha_1(D(\mathbf{U}_s) : D(\mathbf{Z}))_\Omega + \alpha_2(\nabla \cdot \mathbf{U}_s, \nabla \cdot \mathbf{Z})_\Omega - \phi_s(P, \nabla \cdot \mathbf{Z})_\Omega$$

$$- \frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})\mathbf{V}_f, \mathbf{Z})_\Omega = (\mathbf{b}_s, \mathbf{Z})_\Omega \tag{3.5}$$

holds for all $(\mathbf{W}, \mathbf{Z}, q) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$. Here in $\boldsymbol{\zeta}$ is \mathbf{U}_s and $\nabla \cdot \mathbf{U}_s$.

Lemma 1. (Equivalence of weak formulations) *Suppose that parameters and data satisfy assumptions (A) and (B). Then any solution (in the sense of distributions) $(\mathbf{V}_f, \mathbf{U}_s, P) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ of the coupled problem (2.1)–(2.4) is also a solution to the variational problem (3.3)–(3.5). Conversely, any solution to the weak problem (3.3)–(3.5) satisfies (2.1)–(2.4) in the sense of distributions.*

The proof follows using standard arguments. We omit to show it.

3.2 Decoupled problem corresponding to (3.3)–(3.5), Case (a): $\mathbf{K}(\boldsymbol{\zeta}) = \mathbf{K}(\mathbf{U}_s)$

We note that the weak formulation (3.3)–(3.5) can be decoupled concerning the unknowns (\mathbf{V}_f, P) and \mathbf{U}_s . In case (a), we are dealing with $\mathbf{K}(\boldsymbol{\zeta}) = \mathbf{K}(\mathbf{U}_s)$, which is a non-linear function of \mathbf{U}_s and satisfies assumptions (3.1) and (3.2). In this case, we can solve (3.3)–(3.5) sequentially, that is, given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$ find $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ such that

$$(Q_{w_1}(\boldsymbol{\zeta})) \begin{cases} 2(D(\mathbf{V}_f) : D(\mathbf{W}))_\Omega + \lambda(\nabla \cdot \mathbf{V}_f, \nabla \cdot \mathbf{W})_\Omega - \phi_f(P, \nabla \cdot \mathbf{W})_\Omega \\ + \frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})\mathbf{V}_f, \mathbf{W})_\Omega + \phi_f(\nabla \cdot \mathbf{V}_f, q)_\Omega + a_0(P, q)_\Omega \\ = (\mathbf{b}_f, \mathbf{W})_\Omega + (\mathbf{T}_\infty, \mathbf{W})_{\partial\Omega} + (a_0, q)_\Omega \end{cases}$$

holds for all $(\mathbf{W}, q) \in H^1(\Omega)^d \times L^2(\Omega)$, and then for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, find $\mathbf{U}_s \in H_0^1(\Omega)^d$ such that

$$(Q_{w_2}(\mathbf{U}_s)) \begin{cases} 2\alpha_1(D(\mathbf{U}_s) : D(\mathbf{Z}))_\Omega + \alpha_2(\nabla \cdot \mathbf{U}_s, \nabla \cdot \mathbf{Z})_\Omega = \phi_s(P, \nabla \cdot \mathbf{Z})_\Omega \\ + \frac{1}{Da}(\mathbf{K}(\mathbf{U}_s)\mathbf{V}_f, \mathbf{Z})_\Omega + (\mathbf{b}_s, \mathbf{Z})_\Omega \end{cases}$$

holds for all $\mathbf{Z} \in H_0^1(\Omega)^d$. For notational convenience, we denote combined problem as $(Q_{w_1}(\boldsymbol{\zeta})) - (Q_{w_2}(\mathbf{U}_s))$.

3.3 Decoupled problem corresponding to (3.3)–(3.5), Case (b): $\mathbf{K}(\nabla \cdot \boldsymbol{\zeta}) = \mathbf{K}(\nabla \cdot \mathbf{U}_s)$

We can solve (3.3)–(3.5) sequentially, that is, for a given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$ find $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ such that

$$(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta})) \begin{cases} 2(D(\mathbf{V}_f) : D(\mathbf{W}))_\Omega + \lambda(\nabla \cdot \mathbf{V}_f, \nabla \cdot \mathbf{W})_\Omega - \phi_f(P, \nabla \cdot \mathbf{W})_\Omega \\ + \frac{1}{Da}(\mathbf{K}(\nabla \cdot \boldsymbol{\zeta})\mathbf{V}_f, \mathbf{W})_\Omega + \phi_f(\nabla \cdot \mathbf{V}_f, q)_\Omega + a_0(P, q)_\Omega \\ = (\mathbf{b}_f, \mathbf{W})_\Omega + (\mathbf{T}_\infty, \mathbf{W})_{\partial\Omega} + (a_0, q)_\Omega \end{cases}$$

holds for all $(\mathbf{W}, q) \in H^1(\Omega)^d \times L^2(\Omega)$, and then for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, find $\mathbf{U}_s \in H_0^1(\Omega)^d$ such that

$$(Q_{w_2}(\nabla \cdot \mathbf{U}_s)) \begin{cases} 2\alpha_1(D(\mathbf{U}_s) : D(\mathbf{Z}))_\Omega + \alpha_2(\nabla \cdot \mathbf{U}_s, \nabla \cdot \mathbf{Z})_\Omega = \phi_s(P, \nabla \cdot \mathbf{Z})_\Omega \\ + \frac{1}{Da}(\mathbf{K}(\nabla \cdot \mathbf{U}_s)\mathbf{V}_f, \mathbf{Z})_\Omega + (\mathbf{b}_s, \mathbf{Z})_\Omega \end{cases}$$

holds for all $\mathbf{Z} \in H_0^1(\Omega)^d$. For notational convenience, we denote combined problem as $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta})) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$.

3.4 Case (a): existence and uniqueness results for $(Q_{w_1}(\boldsymbol{\zeta}))$

In order to solve weak formulation $(Q_{w_1}(\boldsymbol{\zeta}))$, we use the following method. We rephrase the weak formulation $(Q_{w_1}(\boldsymbol{\zeta}))$ into an abstract setting. Set $\mathbb{Y} = H^1(\Omega)^d \times L^2(\Omega)$. To do so, define a mapping \mathcal{H}_ζ from \mathbb{Y} to \mathbb{Y} by

$$\begin{aligned} \langle \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle &= 2(D(\mathbf{V}_f) : D(\mathbf{W}))_\Omega + \lambda(\nabla \cdot \mathbf{V}_f, \nabla \cdot \mathbf{W})_\Omega - \phi_f(P, \nabla \cdot \mathbf{W})_\Omega \\ &+ \frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})\mathbf{V}_f, \mathbf{W})_\Omega + \phi_f(\nabla \cdot \mathbf{V}_f, q)_\Omega + a_0(P, q)_\Omega - [(\mathbf{b}_f, \mathbf{W})_\Omega + (\mathbf{T}_\infty, \mathbf{W})_{\partial\Omega} + (a_0, q)_\Omega]. \end{aligned} \tag{3.6}$$

Using the mapping \mathcal{H}_ζ , the variational formulation $(Q_{w_1}(\boldsymbol{\zeta}))$ can equivalently be written as: for a given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$, find $(\mathbf{V}_f, P) \in \mathbb{Y}$ such that

$$\langle \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle = 0 \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}. \tag{3.7}$$

Conversely, if (3.7) holds, then (2.1) and (2.3) with the first boundary condition in the equation (2.4) satisfy in the sense of distributions. Hence, our immediate task is to find a pair $(\mathbf{V}_f, P) \in \mathbb{Y}$ that satisfies (3.7). In order to do so, we proceed as follows. The mapping \mathcal{H}_ζ satisfies the following lemma:

Lemma 3.1. *Suppose that parameters and data satisfy assumptions (A) and (B). If \mathcal{H}_ζ is a mapping from \mathbb{Y} into itself defined by (3.6), then the following statements hold:*

- (i) \mathcal{H}_ζ is continuous.
- (ii) There exists a real number $r > 0$ such that

$$\langle \mathcal{H}_\zeta(\mathbb{V}), \mathbb{V} \rangle > 0, \text{ for all } \mathbb{V} \in \mathbb{Y} \text{ with } \|\mathbb{V}\|_{\mathbb{Y}} = r,$$

i.e., \mathcal{H}_ζ is coercive on a ball of radius r in \mathbb{Y} . Here, for any $\mathbb{V} = (\mathbf{V}_f, P) \in \mathbb{Y} = H^1(\Omega)^d \times L^2(\Omega)$, $\|\cdot\|_{\mathbb{Y}}$ is defined as

$$\|\mathbb{V}\|_{\mathbb{Y}}^2 = \|(\mathbf{V}_f, P)\|_{\mathbb{Y}}^2 = \|\mathbf{V}_f\|_{1,\Omega}^2 + \|P\|_{0,\Omega}^2.$$

Proof. (i) The continuity of the mapping \mathcal{H}_ζ can be shown using the continuity of scalar product. Indeed, let $\{\mathbb{V}^m\}_{m \geq 1} = \{(\mathbf{V}_f^m, P^m)\}_{m \geq 1}$ be any sequence in \mathbb{Y} that converges strongly to $\mathbb{V} = (\mathbf{V}_f, P) \in \mathbb{Y}$ as $m \rightarrow \infty$, i.e.

$$\|\mathbf{V}_f^m - \mathbf{V}_f\|_{1,\Omega} \rightarrow 0, \quad \|P^m - P\|_{0,\Omega} \rightarrow 0 \text{ as } m \rightarrow \infty. \tag{3.8}$$

Relying on the definition of \mathcal{H}_ζ and on Cauchy-Schwarz inequality, we get

$$\begin{aligned} |\langle \mathcal{H}_\zeta(\mathbb{V}^m) - \mathcal{H}_\zeta(\mathbb{V}), (\mathbf{W}, q) \rangle| &\leq 2\|\mathbb{D}^f(\mathbf{V}_f^m - \mathbf{V}_f)\|_{0,\Omega}\|\mathbb{D}^f(\mathbf{W})\|_{0,\Omega} \\ &+ \lambda\|\nabla \cdot (\mathbf{V}_f^m - \mathbf{V}_f)\|_{0,\Omega}\|\nabla \cdot \mathbf{W}\|_{0,\Omega} + \phi_f\|P^m - P\|_{0,\Omega}\|\nabla \cdot \mathbf{W}\|_{0,\Omega} \\ &+ \frac{k_2}{Da}\|\mathbf{V}_f^m - \mathbf{V}_f\|_{0,\Omega}\|\mathbf{W}\|_{0,\Omega} + \phi_f\|\nabla \cdot (\mathbf{V}_f^m - \mathbf{V}_f)\|_{0,\Omega}\|q\|_{0,\Omega} \\ &+ a_0\|P^m - P\|_{0,\Omega}\|q\|_{0,\Omega}. \end{aligned}$$

Using (3.8), we obtain

$$|\langle \mathcal{H}_\zeta(\mathbf{V}_f^m, P^m) - \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle| \rightarrow 0 \quad \forall (\mathbf{W}, q) \text{ as } m \rightarrow \infty.$$

This argument establishes the continuity of \mathcal{H}_ζ .

- (ii) For any $\mathbb{V} = (\mathbf{V}_f, P) \in \mathbb{Y}$, we have

$$\begin{aligned} \langle \mathcal{H}_\zeta(\mathbb{V}), \mathbb{V} \rangle &= 2\|D(\mathbf{V}_f)\|_{0,\Omega}^2 + \lambda\|\nabla \cdot \mathbf{V}_f\|_{0,\Omega}^2 - \phi_f(P, \nabla \cdot \mathbf{V}_f)_\Omega + \frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})\mathbf{V}_f, \mathbf{V}_f)_\Omega \\ &+ \phi_f(\nabla \cdot \mathbf{V}_f, P)_\Omega + a_0\|P\|_{0,\Omega}^2 - [(\mathbf{b}_f, \mathbf{V}_f)_\Omega + (\mathbf{T}_\infty, \mathbf{V}_f)_{\partial\Omega} + (a_0, P)_\Omega]. \end{aligned} \tag{3.9}$$

Using Cauchy-Schwarz, Korn's and trace inequalities, we obtain

$$\begin{aligned} \langle \mathcal{H}_\zeta(\mathbb{V}), \mathbb{V} \rangle &\geq \alpha \|\mathbf{V}_f\|_{1,\Omega}^2 + \lambda \|\nabla \cdot \mathbf{V}_f\|_{0,\Omega}^2 + a_0 \|P\|_{0,\Omega}^2 \\ &\quad - (\|\mathbf{b}_f\|_{0,\Omega} + \sqrt{c_i} \|\mathbf{T}_\infty\|_{0,\partial\Omega}) \|\mathbb{V}\|_{1,\Omega} - \|a_0\|_{0,\Omega} \|P\|_{0,\Omega}, \end{aligned} \tag{3.10}$$

where $\alpha = \frac{1}{c_k} \min\{2, \frac{k_1}{Da}\}$. Further, (3.10) can be rewritten as

$$\langle \mathcal{H}_\zeta(\mathbb{V}), \mathbb{V} \rangle \geq \alpha_3 (\|\mathbf{V}_f\|_{1,\Omega}^2 + \|P\|_{0,\Omega}^2) - \alpha_4 \|\mathbb{V}\|_{1,0,\Omega}, \tag{3.11}$$

where

$$\alpha_3 = \min \{ \alpha, a_0 \}, \tag{3.12}$$

$$\alpha_4 = [(\|\mathbf{b}_f\|_{0,\Omega} + \sqrt{c_i} \|\mathbf{T}_\infty\|_{0,\partial\Omega})^2 + \|a_0\|_{0,\Omega}^2]^{1/2}. \tag{3.13}$$

If $\|\mathbb{V}\|_{\mathbb{Y}} = r_0$ for some $r_0 > 0$, then we have

$$\langle \mathcal{H}_\zeta(\mathbb{V}), \mathbb{V} \rangle > 0 \quad \forall \mathbb{V} \in \mathbb{Y}, \quad \text{when } r_0 > \frac{\alpha_4}{\alpha_3}. \tag{3.14}$$

This completes the proof of Lemma 3.1. □

Remark 3.2. Note that $\lambda = \frac{\lambda_f}{\mu_f}$ (which is the ratio of the two viscosity coefficients) plays a significant role in the coercivity proof. The literature [26, 27] suggests that researchers have debated on the sign (or value) of λ . According to the well-known Stokes-hypothesis, $\lambda = -2/3$ [27]. On the other hand, the existing literature also suggests that this ratio can be non-negative [26]. Thus, we consider both of these possibilities. If $\lambda \geq 0$, then the coercivity of \mathcal{H}_ζ , as shown by us in (3.14), holds. However, when $\lambda < 0$ (i.e. a typical Stokes hypothesis), the coercivity of \mathcal{H}_ζ holds with relevant restrictions on the constants. For instance, when $\lambda < 0$, the mapping \mathcal{H}_ζ is coercive if $\alpha = \frac{1}{c_k} \min\{2, \frac{k_1}{Da}\} > 2/3$. For convenience from here onward, we assume λ to be a non-negative constant.

Based on Lemma A.1 (see Appendix A), we now present the following existence results.

Theorem 3.3. *Suppose that (A) and (B) hold. Then for a given $\zeta \in H_0^1(\Omega)^d$, the problem (3.7) has at least one solution $(\mathbf{V}_f, P) \in \mathbb{Y} = H^1(\Omega)^d \times L^2(\Omega)$ satisfying the problems (2.1) and (2.3) with the first boundary condition in the equation (2.4) in the sense of distributions. Moreover, the solution (\mathbf{V}_f, P) satisfies the following a priori bound*

$$\|\mathbf{V}_f\|_{1,\Omega}^2 + \|P\|_{0,\Omega}^2 \leq \left(\frac{\alpha_4}{\alpha_3}\right)^2. \tag{3.15}$$

Proof. To prove this result, we use the Galerkin method. The spaces $H^1(\Omega)^d, L^2(\Omega)$ are separable Hilbert spaces. Hence, there exist corresponding bases $\{\mathbf{W}_i\}_{i=1}^\infty$ and $\{q_i\}_{i=1}^\infty$ of smooth functions. Let \mathbb{Y}_m be the space spanned by $\{(\mathbf{W}_i, q_i)\}_{i=1}^m$. The scalar product on \mathbb{Y}_m is induced by the scalar product of \mathbb{Y} . We define the approximate solution (\mathbf{V}_f^m, P^m) as follows:

$$\mathbf{V}_f^m = \sum_{i=1}^m a_i \mathbf{W}_i, \quad P^m = \sum_{i=1}^m c_i q_i, \tag{3.16}$$

with

$$(Q_m(\zeta)) \begin{cases} 2(D(\mathbf{V}_f^m) : D(\mathbf{W}))_\Omega + \lambda(\nabla \cdot \mathbf{V}_f^m, \nabla \cdot \mathbf{W})_\Omega - \phi_f(P^m, \nabla \cdot \mathbf{W})_\Omega \\ + \frac{1}{Da}(\mathbf{K}(\zeta)\mathbf{V}_f^m, \mathbf{W})_\Omega + \phi_f(\nabla \cdot \mathbf{V}_f^m, q)_\Omega + a_0(P^m, q)_\Omega \\ = (\mathbf{b}_f, \mathbf{W})_\Omega + (\mathbf{T}_\infty, \mathbf{W})_{\partial\Omega} + (a_0, q)_\Omega, \end{cases}$$

holding for all $(\mathbf{W}, q) \in \mathbb{Y}_m$ with $a_i, b_i, c_i \in \mathbb{R}$, for $i = 1, 2, \dots, m$. The task now is to ensure the existence of solutions to $(Q_m(\zeta))$ and show that $(Q_m(\zeta))$ recovers $(Q_{w_1}(\zeta))$ as $m \rightarrow \infty$. The linear structure of $(Q_m(\zeta))$ suggests that weak convergence is enough to pass the limit. Hence, in order to do so, we define a mapping \mathcal{H}_m inspired by the structure of mapping \mathcal{H} as

$$\langle \mathcal{H}_\zeta^m(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle = \langle \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle, \quad \text{for all } (\mathbf{W}, q) \in \mathbb{Y}_m \tag{3.17}$$

where \mathcal{H}_ζ is defined in (3.6). From Lemma 3.1, we deduce that the mapping \mathcal{H}_ζ^m satisfies the conditions needed for Lemma A.1 (see Appendix A) and hence, there exists a solution $(\mathbf{V}_f^m, P^m) \in \mathbb{Y}_m$ for each m such that

$$\langle \mathcal{H}_\zeta^m(\mathbf{V}_f^m, P^m), (\mathbf{W}, q) \rangle = 0, \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}_m. \tag{3.18}$$

It follows that (\mathbf{V}_f^m, P^m) satisfy $(Q_m(\zeta))$ and a_i, c_i can be determined.

Energy Estimates: Let $\mathbf{W} = \mathbf{V}_f^m$, and $q = P^m$ in $(Q_m(\zeta))$, then by performing calculations similar to those leading to (3.11), we obtain

$$\alpha_3(\|\mathbf{V}_f^m\|_{1,\Omega}^2 + \|P^m\|_{0,\Omega}^2) - \alpha_4\|(\mathbf{V}_f^m, P^m)\|_{1,0,\Omega} \leq 0, \tag{3.19}$$

where α_3 and α_4 are defined in (3.12)–(3.13). Consequently,

$$\|\mathbf{V}_f^m\|_{1,\Omega}^2 + \|P^m\|_{0,\Omega}^2 \leq \left(\frac{\alpha_4}{\alpha_3}\right)^2. \tag{3.20}$$

Inequality (3.20) implies that the sequence $\{(\mathbf{V}_f^m, P^m)\}_{m \geq 1}$ is uniformly bounded in \mathbb{Y} . Hence, it has a sub-sequence $\{(\mathbf{V}_f^m, P^m)\}_{m \geq 1}$ (for convenience, we denote it by the same symbol) and a pair $(\mathbf{V}_f, P) \in \mathbb{Y}$ such that

$$(\mathbf{V}_f^m, P^m) \rightharpoonup (\mathbf{V}_f, P) \text{ as } m \rightarrow \infty \text{ weakly in } \mathbb{Y}. \tag{3.21}$$

By taking the limit in (3.18) and using the weak convergence (3.21), we get

$$\langle \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle = 0, \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}_m. \tag{3.22}$$

A continuity argument shows that (3.22) holds for any $(\mathbf{W}, q) \in \mathbb{Y}$. Hence, (\mathbf{V}_f, P) is a solution of (3.7) and equivalently, of the weak formulation $(Q_{w_1}(\zeta))$. Using the lower semi-continuity property of norm in (3.20), we can achieve the following a priori bound on solution (\mathbf{V}_f, P) given by

$$\|\mathbf{V}_f\|_{1,\Omega}^2 + \|P\|_{0,\Omega}^2 \leq \left(\frac{\alpha_4}{\alpha_3}\right)^2. \tag{3.23}$$

□

Proposition 3.4. *Suppose the hypotheses of Theorem 3.3 hold. Then, the weak formulation $(Q_{w_1}(\zeta))$ has a unique solution that depends continuously on the given data.*

Proof. *Uniqueness:* Let (\mathbf{V}_f^1, P^1) and (\mathbf{V}_f^2, P^2) be two solutions that satisfy Equation (3.7) or equivalently the weak formulation $(Q_{w_1}(\zeta))$. Define $(\mathbf{V}_f, P) = (\mathbf{V}_f^1 - \mathbf{V}_f^2, P^1 - P^2)$. Then, from (3.7), we have

$$\langle \mathcal{H}_\zeta(\mathbf{V}_f^1, P^1) - \mathcal{H}_\zeta(\mathbf{V}_f^2, P^2), (\mathbf{W}, q) \rangle = 0 \tag{3.24}$$

for all $(\mathbf{W}, q) \in \mathbb{Y}$. Replace (\mathbf{W}, q) by (\mathbf{V}_f, P) in (3.24) and using the definition of \mathcal{H} , we find

$$\alpha\|\mathbf{V}_f\|_{1,\Omega}^2 + \lambda\|\nabla \cdot \mathbf{V}_f\|_{0,\Omega}^2 + a_0\|P\|_{0,\Omega}^2 \leq 0.$$

The above implies $\mathbf{V}_f = 0, \mathbf{U}_s = 0$ and $P = 0$ almost everywhere in Ω . Hence, the weak formulation $(Q_{w_1}(\zeta))$ has a unique solution. □

Continuous dependence: Let (\mathbf{V}_f^1, P^1) and (\mathbf{V}_f^2, P^2) be two solutions of $(Q_{w_1}(\zeta))$ corresponding to the two sets of data $(\mathbf{T}_{\infty,1}, \mathbf{b}_{f,1}, a_{0,1})$ and $(\mathbf{T}_{\infty,2}, \mathbf{b}_{f,2}, a_{0,2})$, then the difference $(\mathbf{V}_f, P) = (\mathbf{V}_f^1 - \mathbf{V}_f^2, P^1 - P^2)$ satisfies

$$\alpha\|\mathbf{V}_f\|_{1,\Omega}^2 + \lambda\|\nabla \cdot \mathbf{V}_f\|_{0,\Omega}^2 + a_0\|P\|_{0,\Omega}^2 - (\|\mathbf{b}_f\|_{0,\Omega} + \sqrt{c_t}\|\mathbf{T}_\infty\|_{0,\partial\Omega})\|\mathbf{V}\|_{1,\Omega} - \|a_0\|_{0,\Omega}\|P\|_{0,\Omega} \leq 0,$$

or,

$$\begin{aligned} & \| \mathbf{V}_f^1 - \mathbf{V}_f^2 \|_{1,\Omega}^2 + \| P^1 - P^2 \|_{0,\Omega}^2 \\ & \leq \frac{1}{\alpha_3^2} [(\| \mathbf{b}_{f,1} - \mathbf{b}_{f,2} \|_{0,\Omega} + \sqrt{c_f} \| \mathbf{T}_{\infty,1} - \mathbf{T}_{\infty,2} \|_{0,\partial\Omega})^2 + \| a_{0,1} - a_{0,2} \|_{0,\Omega}^2]. \end{aligned} \tag{3.25}$$

Thus, if $(\mathbf{T}_{\infty,1}, \mathbf{b}_{f,1}, a_{0,1})$ is close to $(\mathbf{T}_{\infty,2}, \mathbf{b}_{f,2}, a_{0,2})$, then the left-hand side of (3.25) (the difference of solutions) must be small. This establishes the well-posedness of the auxiliary linear problem $(Q_{w_1}(\boldsymbol{\zeta}))$. Next, we would like to consider the sub-problem, $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$.

3.5 Case (a): existence and uniqueness results for $(Q_{w_2}(\mathbf{U}_s))$

We note that $\mathbf{K}(\boldsymbol{\zeta}) = \mathbf{K}(\mathbf{U}_s)$ is a non-linear function of \mathbf{U}_s (see assumption (A4) in Subsection 1.2) which makes $(Q_{w_2}(\mathbf{U}_s))$ a semilinear problem. By introducing a semilinear form $B(\cdot, \cdot) : H_0^1(\Omega)^d \times H_0^1(\Omega)^d \rightarrow \mathbb{R}$ that is given by

$$B(\mathbf{U}_s, \mathbf{Z}) = 2\alpha_1(D(\mathbf{U}_s) : D(\mathbf{Z}))_{\Omega} + \alpha_2(\nabla \cdot \mathbf{U}_s, \nabla \cdot \mathbf{Z})_{\Omega} - \frac{1}{Da}(\mathbf{K}(\mathbf{U}_s)\mathbf{V}_f, \mathbf{Z})_{\Omega} \tag{3.26}$$

and a linear form $L : H_0^1(\Omega)^d \rightarrow \mathbb{R}$ defined as

$$L(\mathbf{Z}) = \phi_s(P, \nabla \cdot \mathbf{Z})_{\Omega} + (\mathbf{b}_s, \mathbf{Z})_{\Omega}, \tag{3.27}$$

weak problem $(Q_{w_2}(\mathbf{U}_s))$ can be rewritten as an abstract formulation. For a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, find $\mathbf{U}_s \in H_0^1(\Omega)^d$ such that

$$B(\mathbf{U}_s, \mathbf{Z}) = L(\mathbf{Z}) \text{ for all } \mathbf{Z} \in H_0^1(\Omega)^d. \tag{3.28}$$

In order to show existence and uniqueness results for problem (3.28), we will use the Browder-Minty theorem (see Theorem A.3, Appendix A), which is based on the monotone operator approach. To justify the hypotheses of the Browder-Minty theorem, we prove the following results in the form of lemmas.

Lemma 2. *The correspondence $\mathbf{Z} \mapsto B(\mathbf{U}_s, \mathbf{Z})$ is a bounded linear operator and $L \in (H_0^1(\Omega)^d)^*$.*

Proof. Clearly, the mapping $\mathbf{Z} \mapsto B(\mathbf{U}_s, \mathbf{Z})$ is linear (obvious) and bounded. Indeed, using Cauchy-Schwarz, Hölder’s and Sobolev inequalities, we find

$$\begin{aligned} |B(\mathbf{U}_s, \mathbf{Z})| & \leq 2\alpha_1 \|D(\mathbf{U}_s)\|_{0,\Omega} \|D(\mathbf{Z})\|_{0,\Omega} + \alpha_2 \|\nabla \cdot \mathbf{U}_s\|_{0,\Omega} \|\nabla \cdot \mathbf{Z}\|_{0,\Omega} \\ & + \frac{1}{Da} \|\mathbf{K}(\mathbf{U}_s)\|_{L^\infty(\Omega)} \|\mathbf{V}_f\|_{0,\Omega} \|\mathbf{Z}\|_{0,\Omega} \leq \left((2\alpha_1 + \alpha_2) \|\nabla \mathbf{U}_s\|_{0,\Omega} \right. \\ & \left. + \frac{k_2 \sqrt{c_p}}{Da} \|\mathbf{V}_f\|_{0,\Omega} \right) \|\nabla \mathbf{Z}\|_{0,\Omega}. \end{aligned} \tag{3.29}$$

Now, L is linear (obvious) and bounded. Indeed, we have

$$|L(\mathbf{Z})| \leq (\phi_s \|P\|_{0,\Omega} + \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega}) \|\nabla \mathbf{Z}\|_{0,\Omega}. \tag{3.30}$$

This implies $L \in (H_0^1(\Omega)^d)^*$.

The Lemma 2 implies that there exists an operator (non-linear) $\mathcal{A} : H_0^1(\Omega)^d \rightarrow (H_0^1(\Omega)^d)^* = H^{-1}(\Omega)^d$ with

$$(\mathcal{A}\mathbf{U}_s, \mathbf{Z}) = B(\mathbf{U}_s, \mathbf{Z}). \tag{3.31}$$

Thus, the variational problem (3.28) equivalently reduces to the operator equation: find $\mathbf{U}_s \in H_0^1(\Omega)^d$ such that

$$\mathcal{A}\mathbf{U}_s = L \tag{3.32}$$

in the sense

$$(\mathcal{A}\mathbf{U}_s, \mathbf{Z}) = L(\mathbf{Z}), \text{ for all } \mathbf{Z} \in H_0^1(\Omega)^d. \tag{3.33}$$

Further, estimate (3.29) implies the non-linear operator \mathcal{A} is bounded. Indeed,

$$\|\mathcal{A}\mathbf{U}_s\|_{H^{-1}(\Omega)^d} \leq \left[(2\alpha_1 + \alpha_2)\|\nabla\mathbf{U}_s\|_{0,\Omega} + \frac{k_2\sqrt{c_p}}{Da}\|\mathbf{V}_f\|_{0,\Omega} \right]. \tag{3.34}$$

□

Lemma 3. If $\frac{2\alpha_1}{c_k} > \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da}$, then the semilinear form $B(\cdot, \cdot)$ is elliptic that is there exists a constant $c > 0$ such that

$$B(\mathbf{U}_s^1, \mathbf{U}_s^1 - \mathbf{U}_s^2) - B(\mathbf{U}_s^2, \mathbf{U}_s^1 - \mathbf{U}_s^2) \geq c\|\nabla\mathbf{U}_s^1 - \nabla\mathbf{U}_s^2\|_{0,\Omega}^2 \text{ for all } \mathbf{U}_s^1, \mathbf{U}_s^2 \in H_0^1(\Omega)^d. \tag{3.35}$$

Note: This lemma implies that \mathcal{A} is strongly monotone.

Proof: Indeed, consider

$$\begin{aligned} (\mathcal{A}\mathbf{U}_s^1 - \mathcal{A}\mathbf{U}_s^2, \mathbf{U}_s^1 - \mathbf{U}_s^2) &= B(\mathbf{U}_s^1, \mathbf{U}_s^1 - \mathbf{U}_s^2) - B(\mathbf{U}_s^2, \mathbf{U}_s^1 - \mathbf{U}_s^2) \\ &= 2\alpha_1\|D(\mathbf{U}_s^1) - D(\mathbf{U}_s^2)\|_{0,\Omega}^2 + \alpha_2\|\nabla \cdot \mathbf{U}_s^1 - \nabla \cdot \mathbf{U}_s^2\|_{0,\Omega}^2 \\ &\quad - \frac{1}{Da}((\mathbf{K}(\mathbf{U}_s^1) - \mathbf{K}(\mathbf{U}_s^2))\mathbf{V}_f, \mathbf{U}_s^1 - \mathbf{U}_s^2)_\Omega \geq 2\alpha_1\|D(\mathbf{U}_s^1) - D(\mathbf{U}_s^2)\|_{0,\Omega}^2 \\ &\quad + \alpha_2\|\nabla \cdot \mathbf{U}_s^1 - \nabla \cdot \mathbf{U}_s^2\|_{0,\Omega}^2 - \frac{1}{Da}\|\mathbf{K}(\mathbf{U}_s^1) - \mathbf{K}(\mathbf{U}_s^2)\|_{0,\Omega}\|\mathbf{V}_f\|_{L^4(\Omega)}\|\mathbf{U}_s^1 - \mathbf{U}_s^2\|_{L^4(\Omega)} \\ &\geq \frac{2\alpha_1}{c_k}\|\nabla\mathbf{U}_s^1 - \nabla\mathbf{U}_s^2\|_{0,\Omega}^2 - \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da}\|\nabla\mathbf{U}_s^1 - \nabla\mathbf{U}_s^2\|_{0,\Omega}^2 \\ &= \left(\frac{2\alpha_1}{c_k} - \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da} \right) \|\nabla\mathbf{U}_s^1 - \nabla\mathbf{U}_s^2\|_{0,\Omega}^2 \end{aligned} \tag{3.36}$$

To reach (3.36), we have applied Hölder’s, Poincaré’s and Sobolev’s inequalities and Lipschitz continuous property of \mathbf{K} . Thus, if $\frac{2\alpha_1}{c_k} > \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da}$, then B is elliptic with $c = \left(\frac{2\alpha_1}{c_k} - \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da} \right)$.

Lemma 4. The non-linear operator \mathcal{A} as in (3.31) is continuous from $H_0^1(\Omega)^d$ to $H^{-1}(\Omega)^d$.

Proof. Let $\mathbf{U}_s^n \rightarrow \mathbf{U}_s$ in $H_0^1(\Omega)^d$ as $n \rightarrow \infty$. Consider

$$\begin{aligned} |(\mathcal{A}\mathbf{U}_s^n - \mathcal{A}\mathbf{U}_s, \mathbf{Z})| &\leq 2\alpha_1\|D(\mathbf{U}_s^n) - D(\mathbf{U}_s)\|_{0,\Omega}\|D(\mathbf{Z})\|_{0,\Omega} \\ &\quad + \alpha_2\|\nabla \cdot \mathbf{U}_s^n - \nabla \cdot \mathbf{U}_s\|_{0,\Omega}\|\nabla \cdot \mathbf{Z}\|_{0,\Omega} + \frac{1}{Da}\|\mathbf{K}(\mathbf{U}_s^n) - \mathbf{K}(\mathbf{U}_s)\|_{0,\Omega}\|\mathbf{V}_f\|_{L^4(\Omega)}\|\mathbf{Z}\|_{L^4(\Omega)} \end{aligned}$$

On applying Lipschitz continuity of \mathbf{K} and Sobolev inequality, we find

$$\begin{aligned} \|\mathcal{A}\mathbf{U}_s^n - \mathcal{A}\mathbf{U}_s\|_{H^{-1}(\Omega)^d} &\leq 2\alpha_1\|D(\mathbf{U}_s^n) - D(\mathbf{U}_s)\|_{0,\Omega} + \alpha_2\|\nabla \cdot \mathbf{U}_s^n - \nabla \cdot \mathbf{U}_s\|_{0,\Omega} \\ &\quad + \frac{k_L c_s}{Da}\|\mathbf{U}_s^n - \mathbf{U}_s\|_{0,\Omega} \end{aligned}$$

and $\|\mathcal{A}\mathbf{U}_s^n - \mathcal{A}\mathbf{U}_s\|_{H^{-1}(\Omega)^d} \rightarrow 0$ as $n \rightarrow \infty$. That is, \mathcal{A} is continuous. □

Theorem 3.5. Suppose that assumptions (A), (B) and (C) hold. Further, if the given data and non-dimensional parameters satisfy the following assumption

$$\frac{2\alpha_1}{c_k} > \frac{k_L c_s \sqrt{c_p} \alpha_4}{\alpha_3 Da}, \tag{3.37}$$

then for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)^d$, the problem $(Q_{w_2}(\mathbf{U}_s))$ has a unique solution $\mathbf{U}_s \in H_0^1(\Omega)^d$. Moreover, if

$$\frac{2\alpha_1}{c_k} > \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da} \tag{3.38}$$

holds, then \mathbf{U}_s satisfies the following a priori estimate

$$\|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da}\right)} \left(\frac{\phi_s\alpha_4}{\alpha_3} + \sqrt{c_p}\|\mathbf{b}_s\|_{0,\Omega}\right). \tag{3.39}$$

Proof. The analysis shown in Lemmas 3 and 4 justifies that \mathcal{A} satisfies the hypothesis of the Browder-Minty theorem (see Theorem A.3, Appendix A). Consequently, the operator Equation (3.32) or the problem (3.28) has a unique solution $\mathbf{U}_s \in H_0^1(\Omega)^d$ for any given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$. Further, if (3.38) holds, then \mathbf{U}_s satisfies the apriory estimate (3.39). Indeed, from (3.28) replacing $\mathbf{Z} = \mathbf{U}_s$ and using Poincare’s inequality, we have

$$B(\mathbf{U}_s, \mathbf{U}_s) = L(\mathbf{U}_s) \leq (\phi_s\|P\|_{0,\Omega} + \sqrt{c_p}\|\mathbf{b}_s\|_{0,\Omega})\|\nabla \mathbf{U}_s\|_{0,\Omega}$$

Making use of the definition of $B(\mathbf{U}_s, \mathbf{U}_s)$ and Korn’s, Hölder’s and Sobolev inequalities and boundedness property of \mathbf{K} , we obtain

$$\|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da}\right)} \left(\frac{\phi_s\alpha_4}{\alpha_3} + \sqrt{c_p}\|\mathbf{b}_s\|_{0,\Omega}\right).$$

The analysis in Subsections 3.4–3.5 describes the existence and uniqueness of problems $(Q_{w_1}(\boldsymbol{\zeta})) - (Q_{w_2}(\mathbf{U}_s))$. In the next subsections, we focus on developing existence and uniqueness results corresponding to $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta})) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ that is the case (b). □

3.6 Case (b): existence and uniqueness of a solution to $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta}))$

Analysis in this subsection is analogous to the arguments in the Subsection 3.4. Thus, we only state the main theorem and omit the proof.

Theorem 3.6. *Suppose that the assumptions (A) and (B) hold. Then for a given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$, the problem $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta}))$ has a unique solution $(\mathbf{V}_f, P) \in \mathbb{Y} = H^1(\Omega)^d \times L^2(\Omega)$ that depends continuously on the given data. Moreover, the solution (\mathbf{V}_f, P) satisfies the following a priori bound*

$$\|\mathbf{V}_f\|_{1,\Omega}^2 + \|P\|_{0,\Omega}^2 \leq \left(\frac{\alpha_4}{\alpha_3}\right)^2. \tag{3.40}$$

Proof. Proof of this theorem follows from Theorem 3.3 and Proposition 3.4. □

3.7 Case (b): existence and uniqueness of a solution to $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$

We note that $\mathbf{K}(\nabla \cdot \mathbf{U}_s)$ is a non-linear function of $\nabla \cdot \mathbf{U}_s$ (see assumption (A4) in Subsection 1.2) which makes $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ a semilinear problem.

By introducing a semilinear form $B(\cdot, \cdot) : H_0^1(\Omega)^d \times H_0^1(\Omega)^d \rightarrow \mathbb{R}$ that is given by

$$B(\mathbf{U}_s, \mathbf{Z}) = 2\alpha_1(D(\mathbf{U}_s) : D(\mathbf{Z}))_\Omega + \alpha_2(\nabla \cdot \mathbf{U}_s, \nabla \cdot \mathbf{Z})_\Omega - \frac{1}{Da}(\mathbf{K}(\nabla \cdot \mathbf{U}_s)\mathbf{V}_f, \mathbf{Z})_\Omega \tag{3.41}$$

and a linear form $L : H_0^1(\Omega)^d \rightarrow \mathbb{R}$ defined as

$$L(\mathbf{Z}) = \phi_s(P, \nabla \cdot \mathbf{Z})_\Omega + (\mathbf{b}_s, \mathbf{Z})_\Omega, \tag{3.42}$$

weak problem $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ can be rewritten as an abstract formulation. For a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, find $\mathbf{U}_s \in H_0^1(\Omega)^d$ such that

$$B(\mathbf{U}_s, \mathbf{Z}) = L(\mathbf{Z}) \text{ for all } \mathbf{Z} \in H_0^1(\Omega)^d. \tag{3.43}$$

Similar to Subsection 3.5 to show existence and uniqueness results for problem (3.43), we will use the Browder-Minty theorem (see Theorem A.3, Appendix A), which is based on the monotone operator approach. We state the main theorem without proof, which can be done similarly to the proof of Theorem 3.5.

Theorem 3.7. *Suppose that assumptions (A), (B) and (C) hold. Further, if the given data and non-dimensional parameters satisfy the following assumption*

$$\frac{2\alpha_1}{c_k} > \frac{k_L c_s \alpha_4}{\alpha_3 Da}, \tag{3.44}$$

then for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)^d$, the problem $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ has a unique solution $\mathbf{U}_s \in H_0^1(\Omega)^d$. Moreover, if

$$\frac{2\alpha_1}{c_k} > \frac{k_2 \sqrt{c_p} \alpha_4}{\alpha_3 Da} \tag{3.45}$$

holds, then \mathbf{U}_s satisfies the following a priori estimate

$$\|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2 \sqrt{c_p} \alpha_4}{\alpha_3 Da}\right)} \left(\frac{\phi_s \alpha_4}{\alpha_3} + \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} \right). \tag{3.46}$$

Proof. The proof of this theorem can be done similarly to the proof of Theorem 3.5. The analysis in Subsections 3.6–3.7 completes the existence and uniqueness of solution to problem $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta})) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$. In the next section, we focus on the development of existence and uniqueness results corresponding to coupled non-linear problems $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ and $(Q_{w_1}(\nabla \cdot \mathbf{U}_s)) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$, respectively, by converting them into a fixed-point problem. \square

4. Case (a): reduction to a fixed-point problem for $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$

We note that for a given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$, the problem $(Q_{w_1}(\boldsymbol{\zeta}))$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ (see Subsection 3.4). Consequently, we can define a mapping $T_1 : H_0^1(\Omega)^d \rightarrow H^1(\Omega)^d \times L^2(\Omega)$ such that $T_1(\boldsymbol{\zeta}) = (\mathbf{V}_f, P)$. Further, for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, the problem $(Q_{w_2}(\mathbf{U}_s))$ has a unique solution $\mathbf{U}_s \in H_0^1(\Omega)^d$ (see Subsection 3.5). Therefore, we can define a mapping $T_2 : H^1(\Omega)^d \times L^2(\Omega) \rightarrow H_0^1(\Omega)^d$ such that $T_2(\mathbf{V}_f, P) = \mathbf{U}_s$. Now, in order to get the fixed-point problem corresponding to $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$, we define a composition map $T = T_2 \circ T_1 : H_0^1(\Omega)^d \rightarrow H_0^1(\Omega)^d$ such that

$$T(\boldsymbol{\zeta}) = (T_2 \circ T_1)(\boldsymbol{\zeta}) = T_2(T_1(\boldsymbol{\zeta})) = T_2(\mathbf{V}_f, P) = \mathbf{U}_s. \tag{4.1}$$

Thus, a fixed-point of mapping T solves the coupled non-linear problem $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ or equivalently, (3.3)–(3.5) when $\boldsymbol{\zeta} = \mathbf{U}_s$. In order to show that the mapping T has a fixed point, we use Schauder’s fixed-point theorem (see Theorem A.2, Appendix A).² Thus, in the following analysis, we prove some results in the form of lemmas to verify the hypotheses of Schauder’s fixed-point theorem.

4.1 Analysis of the fixed-point problem

Throughout this subsection, we assume that hypotheses of Theorem 3.3, Proposition 3.4 and Theorem 3.5 hold. Then, we have

Lemma 5. *Given $r > 0$, let \mathbf{W}_r be a closed and convex subset of $H^1(\Omega)^d$ defined by*

$$\mathbf{W}_r = \{\mathbf{w} \in H_0^1(\Omega)^d : \|\mathbf{w}\|_{H_0^1(\Omega)^d} = \|\nabla \mathbf{w}\|_{0,\Omega} \leq r\},$$

²For the fixed-point approach, we are inspired by the working techniques used in [38, 39].

and assume that the data satisfies

$$\frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da}\right)} \left(\phi_s \frac{\alpha_4}{\alpha_3} + \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} \right) \leq r. \tag{4.2}$$

Then, $T(\mathbf{W}_r) \subseteq \mathbf{W}_r$.

Proof. For any $\mathbf{w} \in \mathbf{W}_r$, using estimate (3.39), we find

$$\begin{aligned} \|T(\mathbf{w})\|_{H_0^1(\Omega)^d} &= \|T_2(T_1(\mathbf{w}))\|_{H_0^1(\Omega)^d} = \|T_2(\mathbf{V}_f, P)\|_{H_0^1(\Omega)^d} = \|\mathbf{U}_s\|_{H_0^1(\Omega)^d} \\ &= \|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da}\right)} \left(\phi_s \frac{\alpha_4}{\alpha_3} + \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} \right). \end{aligned} \tag{4.3}$$

The above estimate (4.3) together with the assumption (4.2) imply $T(\mathbf{w}) \in \mathbf{W}_r$, which proves $T(\mathbf{W}_r) \subseteq \mathbf{W}_r$. \square

Lemma 6. The map $T_1 : H_0^1(\Omega)^d \rightarrow H^1(\Omega)^d \times L^2(\Omega)$ satisfies

$$\|T_1(\boldsymbol{\zeta}) - T_1(\tilde{\boldsymbol{\zeta}})\|_{\mathbb{Y}} \leq \frac{\sqrt{2}k_L\alpha_4c_s}{\alpha_3^2 Da} \|\boldsymbol{\zeta} - \tilde{\boldsymbol{\zeta}}\|_{0,\Omega}. \tag{4.4}$$

Proof. Given $\boldsymbol{\zeta}, \tilde{\boldsymbol{\zeta}} \in H_0^1(\Omega)^d$, let $(\mathbf{V}_f, P), (\tilde{\mathbf{V}}_f, \tilde{P}) \in H^1(\Omega)^d \times L^2(\Omega)$ be the corresponding solutions of $(Q_{w_1}(\boldsymbol{\zeta}))$. Then, Equation (3.7) implies

$$\langle \mathcal{H}_\zeta(\mathbf{V}_f, P), (\mathbf{W}, q) \rangle = 0 \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}, \tag{4.5}$$

$$\langle \mathcal{H}_{\tilde{\zeta}}(\tilde{\mathbf{V}}_f, \tilde{P}), (\mathbf{W}, q) \rangle = 0 \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}. \tag{4.6}$$

Taking the difference between the above equations, we get

$$\langle \mathcal{H}_\zeta(\mathbf{V}_f, P) - \mathcal{H}_{\tilde{\zeta}}(\tilde{\mathbf{V}}_f, \tilde{P}), (\mathbf{W}, q) \rangle = 0 \text{ for all } (\mathbf{W}, q) \in \mathbb{Y}. \tag{4.7}$$

Replacing $\mathbf{W} = \mathbf{V}_f - \tilde{\mathbf{V}}_f$ and $q = P - \tilde{P}$ in the above equation and using the definition of \mathcal{H}_ζ and $\mathcal{H}_{\tilde{\zeta}}$, we get

$$\begin{aligned} &2\|D(\mathbf{V}_f) - D(\tilde{\mathbf{V}}_f)\|_{0,\Omega}^2 + \lambda\|\nabla \cdot \mathbf{V}_f - \nabla \cdot \tilde{\mathbf{V}}_f\|_{0,\Omega}^2 + a_0\|P - \tilde{P}\|_{0,\Omega}^2 \\ &= -\frac{1}{Da}(\mathbf{K}(\boldsymbol{\zeta})(\mathbf{V}_f - \tilde{\mathbf{V}}_f), \mathbf{V}_f - \tilde{\mathbf{V}}_f)_\Omega - \frac{1}{Da}((\mathbf{K}(\boldsymbol{\zeta}) - \mathbf{K}(\tilde{\boldsymbol{\zeta}}))\tilde{\mathbf{V}}_f, \mathbf{V}_f - \tilde{\mathbf{V}}_f)_\Omega. \end{aligned} \tag{4.8}$$

Using Korn's, Hölder's inequalities and estimate (3.23), we obtain

$$\|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega}^2 + \|P - \tilde{P}\|_{0,\Omega}^2 \leq \frac{\alpha_4}{\alpha_3^2 Da} \|\mathbf{K}(\boldsymbol{\zeta}) - \mathbf{K}(\tilde{\boldsymbol{\zeta}})\|_{0,\Omega} \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{L^4(\Omega)}.$$

Further, using Lipschitz continuous property of \mathbf{K} and Sobolev's inequality, we get

$$\left(\|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega}^2 + \|P - \tilde{P}\|_{0,\Omega}^2 \right) \leq \frac{k_L\alpha_4c_s}{\alpha_3^2 Da} \|\boldsymbol{\zeta} - \tilde{\boldsymbol{\zeta}}\|_{0,\Omega} \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega},$$

or,

$$\|(\mathbf{V}_f, P) - (\tilde{\mathbf{V}}_f, \tilde{P})\|_{\mathbb{Y}} = \left(\|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega}^2 + \|P - \tilde{P}\|_{0,\Omega}^2 \right)^{1/2} \leq \frac{\sqrt{2}k_L\alpha_4c_s}{\alpha_3^2 Da} \|\boldsymbol{\zeta} - \tilde{\boldsymbol{\zeta}}\|_{0,\Omega}. \tag{4.9}$$

This establishes (4.4). \square

Lemma 7. The map $T_2 : H^1(\Omega)^d \times L^2(\Omega) \rightarrow H_0^1(\Omega)^d$ satisfies

$$\|T_2(\mathbf{V}_f, P) - T_2(\tilde{\mathbf{V}}_f, \tilde{P})\|_{H_0^1(\Omega)} \leq \beta \left[\|P - \tilde{P}\|_{0,\Omega} + \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega} \right] \tag{4.10}$$

where $\beta = \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_1c_s\alpha_4\sqrt{c_p}}{\alpha_3 Da}\right)} \max \left\{ \phi_s, \frac{k_2\sqrt{c_p}}{Da} \right\}$.

Proof. Given $(\mathbf{V}_f, P), (\tilde{\mathbf{V}}_f, \tilde{P}) \in H^1(\Omega)^d \times L^2(\Omega)$, let $\mathbf{U}_s, \tilde{\mathbf{U}}_s \in H_0^1(\Omega)^d$ be the corresponding solutions of $(Q_{w_2}(\mathbf{U}_s))$. Then, (3.28) implies

$$B(\mathbf{U}_s, \mathbf{Z}) = L(\mathbf{Z}) \text{ for all } \mathbf{Z} \in H_0^1(\Omega)^d, \tag{4.11}$$

$$B(\tilde{\mathbf{U}}_s, \mathbf{Z}) = L(\mathbf{Z}) \text{ for all } \mathbf{Z} \in H_0^1(\Omega)^d. \tag{4.12}$$

Taking the difference between the above equations, we get

$$\begin{aligned} & 2\alpha_1(D(\mathbf{U}_s) - D(\tilde{\mathbf{U}}_s) : D(\mathbf{Z}))_\Omega + \alpha_2(\nabla \cdot \mathbf{U}_s - \nabla \cdot \tilde{\mathbf{U}}_s, \nabla \cdot \mathbf{Z})_\Omega \\ &= \phi_s(P - \tilde{P}, \nabla \cdot \mathbf{Z})_\Omega + \frac{1}{Da}(\mathbf{K}(\mathbf{U}_s)(\mathbf{V}_f - \tilde{\mathbf{V}}_f), \mathbf{Z})_\Omega \\ & \quad + \frac{1}{Da}((\mathbf{K}(\mathbf{U}_s) - \mathbf{K}(\tilde{\mathbf{U}}_s))\tilde{\mathbf{V}}_f, \mathbf{Z})_\Omega. \end{aligned}$$

Replace $\mathbf{Z} = \mathbf{U}_s - \tilde{\mathbf{U}}_s$ and using Cauchy-Schwarz and Hölder’s inequalities, we get

$$\begin{aligned} 2\alpha_1\|D(\mathbf{U}_s) - D(\tilde{\mathbf{U}}_s)\|_{0,\Omega}^2 + \alpha_2\|\nabla \cdot \mathbf{U}_s - \nabla \cdot \tilde{\mathbf{U}}_s\|_{0,\Omega}^2 &\leq \phi_s\|P - \tilde{P}\|_{0,\Omega}\|\nabla \cdot (\mathbf{U}_s - \tilde{\mathbf{U}}_s)\|_{0,\Omega} \\ & \quad + \frac{1}{Da}\|\mathbf{K}(\mathbf{U}_s)\|_{L^\infty(\Omega)}\|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{0,\Omega}\|\mathbf{U}_s - \tilde{\mathbf{U}}_s\|_{0,\Omega} \\ & \quad + \frac{1}{Da}\|\mathbf{K}(\mathbf{U}_s) - \mathbf{K}(\tilde{\mathbf{U}}_s)\|_{0,\Omega}\|\tilde{\mathbf{V}}_f\|_{L^4(\Omega)}\|\mathbf{U}_s - \tilde{\mathbf{U}}_s\|_{L^4(\Omega)}. \end{aligned}$$

Moreover, using Lipschitz property of \mathbf{K} , Korn’s, Poincare’s and Sobolev’s embedding inequalities and estimate (3.23), we get

$$\begin{aligned} & \|T_2(\mathbf{V}_f, P) - T_2(\tilde{\mathbf{V}}_f, \tilde{P})\|_{H_0^1(\Omega)^d} = \|\mathbf{U}_s - \tilde{\mathbf{U}}_s\|_{H_0^1(\Omega)^d} = \|\nabla \mathbf{U}_s - \nabla \tilde{\mathbf{U}}_s\|_{0,\Omega} \\ & \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_L c_s \alpha_4 \sqrt{c_p}}{\alpha_3 Da}\right)} \left[\phi_s\|P - \tilde{P}\|_{0,\Omega} + \frac{k_2 \sqrt{c_p}}{Da}\|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{0,\Omega} \right], \end{aligned}$$

or,

$$\|T_2(\mathbf{V}_f, P) - T_2(\tilde{\mathbf{V}}_f, \tilde{P})\|_{H_0^1(\Omega)^d} \leq \beta \left[\|P - \tilde{P}\|_{0,\Omega} + \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega} \right] \tag{4.13}$$

where $\beta = \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_L c_s \alpha_4 \sqrt{c_p}}{\alpha_3 Da}\right)} \max \left\{ \phi_s, \frac{k_2 \sqrt{c_p}}{Da} \right\}$. □

Lemma 8. The map $T = T_2 \circ T_1 : H_0^1(\Omega)^d \subset L^2(\Omega)^d \rightarrow H_0^1(\Omega)^d$ satisfies

$$\|T(\boldsymbol{\zeta}) - T(\tilde{\boldsymbol{\zeta}})\|_{H_0^1(\Omega)^d} \leq \frac{2\beta k_L \alpha_4 c_s}{\alpha_3^2 Da} \|\boldsymbol{\zeta} - \tilde{\boldsymbol{\zeta}}\|_{0,\Omega} \leq \frac{2\beta k_L \alpha_4 c_s \sqrt{c_p}}{\alpha_3^2 Da} \|\nabla \boldsymbol{\zeta} - \nabla \tilde{\boldsymbol{\zeta}}\|_{0,\Omega}. \tag{4.14}$$

Proof. From (4.13), we have

$$\begin{aligned} & \|T(\boldsymbol{\zeta}) - T(\tilde{\boldsymbol{\zeta}})\|_{H_0^1(\Omega)^d} = \|T_2(\mathbf{V}_f, P) - T_2(\tilde{\mathbf{V}}_f, \tilde{P})\|_{H_0^1(\Omega)^d} \\ & \leq \beta \left[\|P - \tilde{P}\|_{0,\Omega} + \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega} \right] \end{aligned} \tag{4.15}$$

We achieve (4.14) with the help of (4.9) and (4.15). □

Theorem 4.1. The mapping $T : \mathbf{W}_r \subset H_0^1(\Omega)^d \rightarrow \mathbf{W}_r \subset H_0^1(\Omega)^d$ is continuous and $\overline{T(\mathbf{W}_r)}$ is compact.

Proof. The continuity of T follows in a straightforward manner from (4.14). Now, given a sequence $\{\boldsymbol{\zeta}_k\}_{k \in \mathbb{N}}$ of \mathbf{W}_r which is clearly bounded, there exists a sub-sequence $\{\boldsymbol{\zeta}_{k_j}\} \subset \{\boldsymbol{\zeta}_k\}_{k \in \mathbb{N}}$ and $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$ such that $\boldsymbol{\zeta}_{k_j} \rightarrow \boldsymbol{\zeta}$ in $H_0^1(\Omega)^d$. In this way, thanks to the compact embedding of $H_0^1(\Omega)^d$ in $L^2(\Omega)^d$, which implies $\boldsymbol{\zeta}_{k_j} \rightarrow \boldsymbol{\zeta}$ in $L^2(\Omega)^d$, which combined with (4.14) implies that $T(\boldsymbol{\zeta}_{k_j}) \rightarrow T(\boldsymbol{\zeta})$. This proves that $\overline{T(\mathbf{W}_r)}$ is compact. □

Theorem 4.2. *Suppose that the hypotheses of Theorem 3.3, Proposition 3.4 and Theorem 3.5 hold. Then, the mapping T defined in 4.1 has a fixed point $\mathbf{U}_s \in H_0^1(\Omega)^d$, which in turn implies that coupled problem $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ has a solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$. Further, if*

$$\frac{2\beta k_L \alpha_4 c_s \sqrt{c_p}}{\alpha_3^2 Da} < 1 \tag{4.16}$$

then T has a unique fixed point $\mathbf{U}_s \in H_0^1(\Omega)^d$, which in turn implies that coupled problem $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$.

Proof. The above Theorem 4.1 implies that T satisfies all hypotheses of Schauder’s fixed-point theorem (see Theorem A.2, Appendix A). Consequently, the mapping T has a fixed-point $\mathbf{U}_s \in H_0^1(\Omega)$. This implies that coupled problem $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ or equivalently, (3.3)–(3.5) when $\boldsymbol{\zeta} = \mathbf{U}_s$ has a solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$. Further, if (4.12) is true, then $T : \mathbf{W}_r \rightarrow \mathbf{W}_r$ is a strict contraction mapping. That implies T has a unique fixed point $\mathbf{U}_s \in H_0^1(\Omega)^d$ due to Banach’s fixed-point theorem (see p. 415 [34]). This implies that coupled problem $(Q_{w_1}(\mathbf{U}_s)) - (Q_{w_2}(\mathbf{U}_s))$ or equivalently, (3.3)–(3.5) when $\boldsymbol{\zeta} = \mathbf{U}_s$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$. □

Remark 4.3 (Continuous dependence). If the non-dimensional parameters and constants satisfy the following assumption

$$2\alpha^* Da > \frac{k_L \alpha_4 c_s}{\alpha_3}, \quad \frac{4\alpha_1 Da}{c_k} > \left(\frac{c_p k_2^2}{k_1} + \frac{k_L \alpha_4 c_s (c_p + 2\sqrt{c_p})}{\alpha_3} \right), \quad 2\alpha_2 \geq \frac{\phi_s^2}{a_0}, \tag{4.17}$$

where $\alpha^* = \frac{1}{c_k} \min\{2, \frac{k_1}{2Da}\}$. Then, the continuous dependence (3.25) holds. Indeed, we have

$$\begin{aligned} & \| \mathbf{V}_f^1 - \mathbf{V}_f^2 \|_{1,\Omega}^2 + \| \nabla(\mathbf{U}_s^1 - \mathbf{U}_s^2) \|_{0,\Omega}^2 + \| P^1 - P^2 \|_{0,\Omega}^2 \leq \frac{1}{\alpha_6^2} [(\| \mathbf{b}_{f,1} - \mathbf{b}_{f,2} \|_{0,\Omega} \\ & + \sqrt{c_i} \| \mathbf{T}_{\infty,1} - \mathbf{T}_{\infty,2} \|_{0,\partial\Omega})^2 + \| a_{0,1} - a_{0,2} \|_{0,\Omega}^2 + c_p \| \mathbf{b}_{s,1} - \mathbf{b}_{s,2} \|_{0,\Omega}^2], \end{aligned} \tag{4.18}$$

where $\alpha_5 = \min \left\{ \alpha^* - \frac{k_L \alpha_4 c_s}{2\alpha_3 Da}, \left(\frac{2\alpha_1}{c_k} - \frac{c_p k_2^2}{2k_1 Da} - \frac{k_L \alpha_4 c_s (c_p + 2\sqrt{c_p})}{2\alpha_3 Da} \right), \frac{a_0}{2} \right\}$.

5. Case (b): reduction to a fixed-point problem to $(Q_{w_1}(\nabla \cdot \mathbf{U}_s)) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$

We note that for a given $\boldsymbol{\zeta} \in H_0^1(\Omega)^d$, the problem $(Q_{w_1}(\nabla \cdot \boldsymbol{\zeta}))$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ (see Subsection 3.6). Consequently, we can define a mapping $T_1 : H_0^1(\Omega)^d \rightarrow H^1(\Omega)^d \times L^2(\Omega)$ such that $T_1(\boldsymbol{\zeta}) = (\mathbf{V}_f, P)$. Further, for a given pair $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$, the problem $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ has a unique solution $\mathbf{U}_s \in H_0^1(\Omega)^d$ (see Subsection 3.7). Therefore, we can define a mapping $T_2 : H^1(\Omega)^d \times L^2(\Omega) \rightarrow H_0^1(\Omega)^d$ such that $T_2(\mathbf{V}_f, P) = \mathbf{U}_s$. Now, in order to get the fixed-point problem corresponding to $(Q_{w_1}(\nabla \cdot \mathbf{U}_s)) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$, we define a composition map $T = T_2 \circ T_1 : H_0^1(\Omega)^d \rightarrow H_0^1(\Omega)^d$ such that

$$T(\boldsymbol{\zeta}) = T_2(T_1(\boldsymbol{\zeta})) = T_2(\mathbf{V}_f, P) = \mathbf{U}_s. \tag{5.1}$$

Thus, a fixed-point of mapping T solves the coupled non-linear problem $(Q_{w_1}(\nabla \cdot \mathbf{U}_s))$ and $(Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ or equivalently, (3.3)–(3.5) when $\boldsymbol{\zeta} = \nabla \cdot \mathbf{U}_s$. In order to show that the mapping T has a fixed point, we use Schauder’s fixed-point theorem (see Theorem A.2, Appendix A). Thus, in the following analysis, we state some results in the form of lemmas to verify the hypotheses of Schauder’s fixed-point theorem. The proof of the following results is almost similar to the proof presented in Subsection 4.1, so we omit the details.

5.1 Analysis of the fixed-point problem

Throughout this subsection, we assume that hypotheses of Theorems 3.6 and 3.7 hold. Then, we have

Lemma 9. Given $r > 0$, let \mathbf{W}_r be a closed and convex subset of $H^1(\Omega)^d$ defined by

$$\mathbf{W}_r = \{\mathbf{w} \in H_0^1(\Omega)^d : \|\mathbf{w}\|_{H_0^1(\Omega)^d} = \|\nabla \mathbf{w}\|_{0,\Omega} \leq r\}$$

and assume that the data satisfy

$$\frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_2\sqrt{c_p}\alpha_4}{\alpha_3 Da}\right)} \left(\frac{\phi_s\alpha_4}{\alpha_3} + \sqrt{c_p}\|\mathbf{b}_s\|_{0,\Omega}\right) \leq r. \tag{5.2}$$

Then, $T(\mathbf{W}_r) \subseteq \mathbf{W}_r$.

Proof. This lemma is proved similar to Lemma 5. □

Lemma 10. The map $T_1 : H_0^1(\Omega)^d \rightarrow H^1(\Omega)^d \times L^2(\Omega)$ satisfies

$$\|T_1(\boldsymbol{\zeta}) - T_1(\tilde{\boldsymbol{\zeta}})\|_{\mathbb{Y}} \leq \frac{\sqrt{2}k_L\alpha_4c_s}{\alpha_3^2 Da} \|\nabla \cdot \boldsymbol{\zeta} - \nabla \cdot \tilde{\boldsymbol{\zeta}}\|_{0,\Omega}, \tag{5.3}$$

(see Lemma 3.1 for the definition of the space \mathbb{Y}).

Proof. This lemma is proved similar to Lemma 6. □

Lemma 11. The map $T_2 : H^1(\Omega)^d \times L^2(\Omega) \rightarrow H_0^1(\Omega)^d$ satisfies

$$\|T_2(\mathbf{V}_f, P) - T_2(\tilde{\mathbf{V}}_f, \tilde{P})\|_{H_0^1(\Omega)^d} \leq \tilde{\beta} \left[\|P - \tilde{P}\|_{0,\Omega} + \|\mathbf{V}_f - \tilde{\mathbf{V}}_f\|_{1,\Omega} \right], \tag{5.4}$$

where $\tilde{\beta} = \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{k_L\alpha_4c_s}{\alpha_3 Da}\right)} \max \left\{ \phi, \frac{k_2\sqrt{c_p}}{Da} \right\}$.

Proof. This lemma is proved similar to Lemma 7. □

Lemma 12. The map $T = T_2 \circ T_1 : H_0^1(\Omega)^d \rightarrow H_0^1(\Omega)^d$ satisfies

$$\|T(\boldsymbol{\zeta}) - T(\tilde{\boldsymbol{\zeta}})\|_{H_0^1(\Omega)^d} \leq \frac{2\tilde{\beta}k_L\alpha_4c_s}{\alpha_3^2 Da} \|\nabla \cdot \boldsymbol{\zeta} - \nabla \cdot \tilde{\boldsymbol{\zeta}}\|_{0,\Omega} \leq \frac{2\tilde{\beta}k_L\alpha_4c_s}{\alpha_3^2 Da} \|\nabla \boldsymbol{\zeta} - \nabla \tilde{\boldsymbol{\zeta}}\|_{0,\Omega}. \tag{5.5}$$

Proof. This lemma is proved similar to Lemma 8. □

Theorem 5.1. Suppose that the hypotheses of Theorems 3.6 and 3.7 hold and if the following assumption

$$\frac{2\tilde{\beta}k_L\alpha_4c_s}{\alpha_3^2 Da} < 1 \tag{5.6}$$

holds then T has a unique fixed point $\mathbf{U}_s \in H_0^1(\Omega)^d$, which in turn implies that coupled problem $(Q_{w_1}(\nabla \cdot \mathbf{U}_s)) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$.

Proof. If $\frac{4\tilde{\beta}k_L\alpha_4c_s}{\alpha_3^2 Da} < 1$, then Lemma 12 implies $T : \mathbf{W}_r \subset H_0^1(\Omega)^d \rightarrow \mathbf{W}_r \subset H_0^1(\Omega)^d$ is a strict contraction mapping. That implies T has a unique fixed point $\mathbf{U}_s \in H_0^1(\Omega)^d$ due to Banach’s fixed-point theorem. That means the coupled problem $(Q_{w_1}(\nabla \cdot \mathbf{U}_s)) - (Q_{w_2}(\nabla \cdot \mathbf{U}_s))$ or equivalently, the problem (3.3)–(3.5) with $\boldsymbol{\zeta} = \nabla \cdot \mathbf{U}_s$ has a unique solution $(\mathbf{V}_f, P) \in H^1(\Omega)^d \times L^2(\Omega)$ and $\mathbf{U}_s \in H_0^1(\Omega)^d$. □

Remark 5.2.

- If the non-dimensional parameters and the constants satisfy the following constraints

$$2\alpha^* Da > \frac{k_Lc_s\alpha_4}{\alpha_3}, \frac{4\alpha_1 Da}{c_k} > \left(\frac{c_p k_2^2}{k_1} + \frac{3k_Lc_s\alpha_4}{\alpha_3}\right), 2\alpha_2 \geq \frac{\phi_s^2}{a_0}. \tag{5.7}$$

then the continuous dependence (4.18) holds with the modified constant

$$\alpha_6 = \min \left\{ \alpha^* - \frac{k_L c_s \alpha_4}{2\alpha_3 Da}, \left(\frac{2\alpha_1}{c_k} - \frac{c_p k_2^2}{2k_1 Da} - \frac{3k_L c_s \alpha_4}{2\alpha_3 Da} \right), \frac{a_0}{2} \right\}.$$

6. Unbounded K

One may note that theorems in Sections 3, 4 and 5 are proven under the boundedness assumption (i) of (3.1) and Lipschitz continuity (3.2) of $\mathbf{K} = \mathbf{K}(\boldsymbol{\zeta})$. In this section, we would like to relax such assumptions for case (a) $\boldsymbol{\zeta} = \mathbf{U}_s$ and case (b) $\boldsymbol{\zeta} = \nabla \cdot \mathbf{U}_s$. Instead of boundedness property (ii) of (3.1), we assume there exists a constant $k_0 > 0$, such that following sub-linear growth condition holds

$$\|\mathbf{K}(\mathbf{x})\| \leq k_0(1 + \|\mathbf{x}\|) \text{ for all } \mathbf{x} \in \mathbb{R}^d. \tag{6.1}$$

The above growth condition yields for case (a)

$$\|\mathbf{K}(\mathbf{U}_s)\|_{0,\Omega} \leq \sqrt{2}k_0(\sqrt{|\Omega|} + \|\mathbf{U}_s\|_{0,\Omega}). \tag{6.2}$$

We have the following result for case (a).

Theorem 6.1. *Assume that the data and parameters satisfy the assumption (A), (i) of (3.1) and (3.2), (3.37), (3.38) and \mathbf{K} satisfy the growth condition (6.2). Further, if the following parameter constraint*

$$\frac{2\alpha_1}{c_k} > \frac{\sqrt{2c_p} c_s k_0 \alpha_4}{\alpha_3 Da} \tag{6.3}$$

is satisfied, then (3.3)–(3.5) together with $\boldsymbol{\zeta} = \mathbf{U}_s$ has a solution $(\mathbf{V}_f, \mathbf{U}_s, P) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ such that

$$\|(\mathbf{V}_f, P)\|_{\mathbb{Y}} \leq \frac{\alpha_4}{\alpha_3} \tag{6.4}$$

and

$$\|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{\sqrt{2c_p} c_s k_0 \alpha_4}{\alpha_3 Da} \right)} \left[\sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} + \frac{\alpha_4}{\alpha_3} \left(\phi_s + \frac{\sqrt{2|\Omega|} c_s k_0}{Da} \right) \right] \tag{6.5}$$

where $|\Omega|$ denotes the area or volume of the domain Ω .

Proof. We are inspired by the working techniques from [28]. Here, we are dealing with particular non-linear structures of the hydraulic resistivity. We approximate the operators \mathbf{K} with a sequence of uniformly positive definite, bounded operators, $\{\mathbf{K}^m\}_{m \geq 1}$ defined by

$$\mathbf{K}_{ij}^m(\mathbf{U}_s) := \min(m, \mathbf{K}_{ij}(\mathbf{U}_s)), \text{ for all } m \in \mathbb{N}. \tag{6.6}$$

Since \mathbf{K}^m is bounded for each m , Theorem 4.2 implies that there exists a triplet $(\mathbf{V}_f^m, \mathbf{U}_s^m, P^m)$ in $H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ such that

$$\begin{aligned} & 2(D(\mathbf{V}_f^m) : D(\mathbf{W}))_{\Omega} + \lambda(\nabla \cdot \mathbf{V}_f^m, \nabla \cdot \mathbf{W})_{\Omega} - \phi_f(P^m, \nabla \cdot \mathbf{W})_{\Omega} \\ & + \frac{1}{Da}(\mathbf{K}^m(\mathbf{U}_s^m) \mathbf{V}_f^m, \mathbf{W})_{\Omega} + \phi_f(\nabla \cdot \mathbf{V}_f^m, q)_{\Omega} + a_0(P^m, q)_{\Omega} \\ & = (\mathbf{b}_f, \mathbf{W})_{\Omega} + (\mathbf{T}_{\infty}, \mathbf{W})_{\partial\Omega} + (a_0, q)_{\Omega}, \end{aligned} \tag{6.7}$$

$$\begin{aligned} & 2\alpha_1(D(\mathbf{U}_s^m) : D(\mathbf{Z}))_{\Omega} + \alpha_2(\nabla \cdot \mathbf{U}_s^m, \nabla \cdot \mathbf{Z})_{\Omega} = \phi_s(P^m, \nabla \cdot \mathbf{Z})_{\Omega} \\ & + \frac{1}{Da}(\mathbf{K}^m(\mathbf{U}_s^m) \mathbf{V}_f^m, \mathbf{Z})_{\Omega} + (\mathbf{b}_s, \mathbf{Z})_{\Omega}. \end{aligned} \tag{6.8}$$

We desire a uniform bound for the sequence $\{(\mathbf{V}_f^m, \mathbf{U}_s^m, P^m)\}$, which is independent of m . In order to do so, replace (\mathbf{W}, q) by (\mathbf{V}_f^m, P^m) in (6.7). Using the positive definiteness of \mathbf{K}^m and the Cauchy-Schwarz, trace, Korn’s inequalities, we get the following estimate

$$\|(\mathbf{V}_f^m, P^m)\|_{\mathbb{Y}} \leq \frac{\alpha_4}{\alpha_3}. \tag{6.9}$$

One can observe the right-hand side of (6.9) is independent of m . Next, replacing \mathbf{Z} by \mathbf{U}_s^m in (6.8) and use Hölder’s, Poincaré’s, Korn’s, Sobolev inequalities and (6.1), to get

$$\begin{aligned} \frac{2\alpha_1}{c_k} \|\nabla \mathbf{U}_s^m\|_{0,\Omega}^2 &\leq \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} \|\nabla \mathbf{U}_s^m\|_{0,\Omega} + \phi_s \|P^m\|_{0,\Omega} \|\nabla \cdot \mathbf{U}_s^m\|_{0,\Omega} \\ &\quad + \frac{1}{Da} \|\mathbf{K}(\mathbf{U}_s^m)\|_{0,\Omega} \|\mathbf{V}_f^m\|_{L^4(\Omega)} \|\mathbf{U}_s^m\|_{L^4(\Omega)} \end{aligned}$$

$$\begin{aligned} \frac{2\alpha_1}{c_k} \|\nabla \mathbf{U}_s^m\|_{0,\Omega}^2 &\leq \sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} \|\nabla \mathbf{U}_s^m\|_{0,\Omega} + \phi_s \|P^m\|_{0,\Omega} \|\nabla \mathbf{U}_s^m\|_{0,\Omega} \\ &\quad + \frac{c_s}{Da} [\sqrt{2}k_0(\sqrt{\Omega} + \sqrt{c_p} \|\nabla \mathbf{U}_s^m\|_{0,\Omega})] \|\mathbf{V}_f^m\|_{1,\Omega} \|\nabla \mathbf{U}_s^m\|_{0,\Omega} \end{aligned}$$

$$\|\nabla \mathbf{U}_s^m\|_{0,\Omega} \leq \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{\sqrt{2c_p} c_s k_0 \alpha_4}{\alpha_3 Da}\right)} \left[\sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} + \frac{\alpha_4}{\alpha_3} \left(\phi_s + \frac{\sqrt{2|\Omega|} c_s k_0}{Da} \right) \right] \tag{6.10}$$

The estimates (6.9) and (6.10) imply that $(\mathbf{V}_f^m, \mathbf{U}_s^m, P^m) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ uniformly bounded for all $m \geq 1$. Hence, there exists a triplet $(\mathbf{V}_f, \mathbf{U}_s, P) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ and a sub-sequence of $(\mathbf{V}_f^m, \mathbf{U}_s^m, P^m)$ (we denote by the same symbol) such that

$$(\mathbf{V}_f^m, \mathbf{U}_s^m, P^m) \rightharpoonup (\mathbf{V}_f, \mathbf{U}_s, P) \text{ weakly in } H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega), \tag{6.11}$$

and the compact embedding $H^1(\Omega)^d \hookrightarrow L^4(\Omega)^d$ yields

$$(\mathbf{V}_f^m, \mathbf{U}_s^m) \rightarrow (\mathbf{V}_f, \mathbf{U}_s) \text{ strongly in } L^4(\Omega)^d \times L^4(\Omega)^d. \tag{6.12}$$

We seek to pass the limit in (6.7)–(6.8) as $m \rightarrow \infty$. We observe that the weak convergence (6.11) is sufficient to pass the limit in the linear terms of (6.7)–(6.8); however, the non-linear terms demand strong convergence (6.12). Subsequently, the non-linear terms of the problem (6.7)–(6.8), which can be rewritten as

(i) for (6.7)

$$\underbrace{((\mathbf{K}^m(\mathbf{U}_s^m) - \mathbf{K}(\mathbf{U}_s))\mathbf{V}_f^m, \mathbf{W})_\Omega} + \underbrace{(\mathbf{K}(\mathbf{U}_s)(\mathbf{V}_f^m - \mathbf{V}_f), \mathbf{W})_\Omega} + (\mathbf{K}(\mathbf{U}_s)\mathbf{V}_f, \mathbf{W})_\Omega \tag{6.13}$$

(ii) for (6.8)

$$\underbrace{((\mathbf{K}^m(\mathbf{U}_s^m) - \mathbf{K}(\mathbf{U}_s))\mathbf{V}_f^m, \mathbf{Z})_\Omega} + \underbrace{(\mathbf{K}(\mathbf{U}_s)(\mathbf{V}_f^m - \mathbf{V}_f), \mathbf{Z})_\Omega} + (\mathbf{K}(\mathbf{U}_s)\mathbf{V}_f, \mathbf{Z})_\Omega, \tag{6.14}$$

Since \mathbf{U}_s^m converges to \mathbf{U}_s strongly in $L^2(\Omega)^d$, it implies $\mathbf{U}_s^m - \mathbf{U}_s \rightarrow 0$ a.e. in Ω up to a sub-sequence. The fact that \mathbf{K} is Lipschitz guarantees that

$$\mathbf{K}^m(\mathbf{U}_s^m) - \mathbf{K}(\mathbf{U}_s) \rightarrow 0 \text{ a.e. in } \Omega. \tag{6.15}$$

Indeed, we have

$$\begin{aligned} |\mathbf{K}^m(\mathbf{U}_s^m)(\mathbf{x}) - \mathbf{K}(\mathbf{U}_s)(\mathbf{x})| &= |\mathbf{K}^m(\mathbf{U}_s^m)(\mathbf{x}) - \mathbf{K}^m(\mathbf{U}_s)(\mathbf{x}) + \mathbf{K}^m(\mathbf{U}_s)(\mathbf{x}) + \mathbf{K}(\mathbf{U}_s)(\mathbf{x})| \\ &\leq k_L |\mathbf{U}_s^m(\mathbf{x}) - \mathbf{U}_s(\mathbf{x})| + |\mathbf{K}^m(\mathbf{U}_s)(\mathbf{x}) - \mathbf{K}(\mathbf{U}_s)(\mathbf{x})|. \end{aligned}$$

As $\mathbf{U}_s^m - \mathbf{U}_s \rightarrow 0$ a.e. in Ω together with the definition of the truncation function implies $\mathbf{K}^m(\mathbf{U}_s) - \mathbf{K}(\mathbf{U}_s) \rightarrow 0$ a.e. in Ω . This establishes (6.15). Using the bound (6.9) and the convergence result (6.15), we get

$$\lim_{m \rightarrow \infty} ((\mathbf{K}^m(\mathbf{U}_s^m) - \mathbf{K}(\mathbf{U}_s))\mathbf{V}_f^m, \mathbf{W})_\Omega = 0, \quad \lim_{m \rightarrow \infty} ((\mathbf{K}^m(\mathbf{U}_s^m) - \mathbf{K}(\mathbf{U}_s))\mathbf{V}_f^m, \mathbf{Z})_\Omega = 0. \tag{6.16}$$

The bound (6.10) together with estimate (6.2), and the strong convergence $\mathbf{V}_f^m \rightarrow \mathbf{V}_f$ in $L^4(\Omega)^d$ leads to

$$\lim_{m \rightarrow \infty} (\mathbf{K}(\mathbf{U}_s)(\mathbf{V}_f^m - \mathbf{V}_f), \mathbf{W})_\Omega = 0, \quad \lim_{m \rightarrow \infty} (\mathbf{K}(\mathbf{U}_s)(\mathbf{V}_f^m - \mathbf{V}_f), \mathbf{Z})_\Omega = 0 \tag{6.17}$$

Thus, the terms with under braces in (6.13) and (6.14) tend to zero as $m \rightarrow \infty$. Hence, we can say that (6.7)–(6.8) recovers (3.3)–(3.5) as $m \rightarrow \infty$. This completes the proof of the Theorem 6.1. \square

Remark 6.2.

- We note that in the above case i.e. when $\mathbf{K}(\mathbf{U}_s)$ is not bounded, however, satisfies the sub-linear growth condition (6.1), the uniqueness of solutions holds whenever the non-dimensional parameters and constants satisfy the following inequalities:

$$\frac{2\alpha Da}{c_s} > \left[\frac{k_L \sqrt{c_p} \alpha_4}{\alpha_3} + \sqrt{2}k_0(\sqrt{|\Omega|} + \sqrt{c_p} \alpha_7) \right], \quad 2\alpha_2 \geq \frac{\phi_s^2}{a_0}, \tag{6.18}$$

$$\frac{4\alpha_1 Da}{c_k c_s} > \left[\frac{3k_L \sqrt{c_p} \alpha_4}{\alpha_3} + \sqrt{2}k_0(\sqrt{|\Omega|} + \sqrt{c_p} \alpha_7) \right], \tag{6.19}$$

where $\alpha_7 = \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{\sqrt{2}c_p c_s k_0 \alpha_4}{\alpha_3 Da}\right)} \left[\sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} + \frac{\alpha_4}{\alpha_3} \left(\phi_s + \frac{\sqrt{2|\Omega|} c_s k_0}{Da} \right) \right]$.

- In case (b) that is when $\boldsymbol{\zeta} = \nabla \cdot \mathbf{U}_s$, the sub-linear growth condition (6.1) becomes

$$\|\mathbf{K}(\nabla \cdot \mathbf{U}_s)\|_{0,\Omega} \leq \sqrt{2}k_0(\sqrt{|\Omega|} + \|\nabla \cdot \mathbf{U}_s\|_{0,\Omega}). \tag{6.20}$$

The specific structure $\mathbf{K}(\nabla \cdot \mathbf{U}_s) = [\gamma_1 + \gamma_2|\nabla \cdot \mathbf{U}_s|]\mathbf{I}$ falls under this case, and we can clearly see that \mathbf{K} satisfies (6.20). In this case, existence and uniqueness analysis can be developed based on similar lines as in Theorem 6.1. Indeed, we have the following theorem.

Theorem 6.3. Assume that the data and parameters satisfy the assumption (A), (i) of (3.1) and (3.2), (3.44), (3.45) and \mathbf{K} satisfy the growth condition (6.20). Further, if the following parameter constraint

$$\frac{2\alpha_1}{c_k} > \frac{\sqrt{2} c_s k_0 \alpha_4}{\alpha_3 Da} \tag{6.21}$$

is satisfied, then (3.3)–(3.5) together with $\boldsymbol{\zeta} = \nabla \cdot \mathbf{U}_s$ has a solution $(\mathbf{V}_f, \mathbf{U}_s, P) \in H^1(\Omega)^d \times H_0^1(\Omega)^d \times L^2(\Omega)$ such that

$$\|(\mathbf{V}_f, P)\|_{\mathbb{Y}} \leq \frac{\alpha_4}{\alpha_3}, \quad \|\nabla \mathbf{U}_s\|_{0,\Omega} \leq \alpha_8 \tag{6.22}$$

where $|\Omega|$ denotes the area or volume of the domain Ω and

$$\alpha_8 = \frac{1}{\left(\frac{2\alpha_1}{c_k} - \frac{\sqrt{2} c_s k_0 \alpha_4}{\alpha_3 Da}\right)} \left[\sqrt{c_p} \|\mathbf{b}_s\|_{0,\Omega} + \frac{\alpha_4}{\alpha_3} \left(\phi_s + \frac{\sqrt{2|\Omega|} c_s k_0}{Da} \right) \right].$$

Further, the solution is unique and subject to the following constraints

$$\frac{2\alpha Da}{c_s} > \left[\frac{k_L \alpha_4}{\alpha_3} + \sqrt{2}k_0(\sqrt{|\Omega|} + \alpha_8) \right], \quad 2\alpha_2 \geq \frac{\phi_s^2}{a_0}, \tag{6.23}$$

$$\frac{4\alpha_1 Da}{c_k c_s} > \left[\frac{3k_L \alpha_4}{\alpha_3} + \sqrt{2}k_0(\sqrt{|\Omega|} + \alpha_8) \right]. \tag{6.24}$$

Proof. The proof of this theorem is similar to Theorem 6.1, we omit the details. \square

Table 1. Dimensionless poro-elasto-hydrodynamics parameters corresponding to tumour tissue with their value range.

Dimensionless parameter	Range of values	Supporting references
Da	$10^{-4} - 10^{-1}$	[1, 30]
α_t	$0 < \alpha_t \leq 10$	[22]
ϱ_t	$10^2 \leq \varrho_t \leq 10^5$	[30]
ν_p	$0.45 \leq \nu_p \leq 0.49$	[31, 32]
ϕ_f	$0.6 \leq \phi_f \leq 0.8$	[1, 30]
K	$0.00006 \leq K \leq 1.4$	[30]

6.1 Comments on parameter restrictions

It may be noted that the existence and uniqueness of results (e.g. see Theorems 3.3, 3.5, 3.6, 3.7, 4.2, 5.1, 6.1 and 6.3, etc.) that are established in this work hold subject to certain parameter restrictions see e.g. (3.37), (3.38), (3.44), (3.45), (4.16), (4.17), (5.6), (5.7), (6.3), (6.18), (6.19), (6.21), (6.23) and (6.24) etc. Such a situation is typical in the case of multiphase mixture models where extra care needs to be paid to the physical admissibility of the parameters. See also Vromans et al. [29]. Further, some of the parameters in these inequalities do show a dependency on material properties of the tissue, like Poisson ratio, Young’s modulus etc. Thus, one has to ensure that these inequalities are satisfied simultaneously. This certainly depends on the choice of relevant parameters. We have collected data from the existing literature on parameters like Da , α_t , ϱ_t etc. which are relevant to biological tissues. We have then verified that there do exist parameter combinations within the given ranges in Table 1 that obey all the inequalities. This ensures that these assumptions (3.37), (3.38), (3.44), (3.45), (4.16), (4.17), (5.6), (5.7), (6.3), (6.18), (6.19), (6.21), (6.23) and (6.24) etc. can be interpreted from the point of view of various applications. We list the dimensionless poro-elasto-hydrodynamics parameters in Table 1. Further, we choose Ω as a d-dimensional ($d = 3$) sphere of unit radius (in dimension 1 mm) then $|\Omega| = \frac{4\pi}{3}$, $|\partial\Omega| = 4\pi$. We set $\mathbf{b}_i = 0$, $i \in \{f, s\}$, $\mathbf{T}_\infty = (1, 0, 0)$, and choose the numerical values for constants $c_k(\Omega) > 0$, $c_p(\Omega) > 0$, $c_s(d) > 0$, $c_i(\Omega, d) > 0$ as follows: $c_k = 2$ (or 3) for $H_0^1(\Omega)^d$ (or $H^1(\Omega)^d$) [33, 34] and $c_p = 1/2$, $c_s = 1/2$ [35], $c_t = 2$ [34]. To verify the inequalities (3.37), (3.38), (3.44), (3.45), (4.16), (4.17), (5.6), (5.7), (6.3) and (6.21), we use the following combination of parameters from Table 1: $Da = 2 \times 10^{-2}$, $\alpha_t = 1$, $\varrho_t = 10^4$, $\nu_p = 0.45$, $k_1 = 0.5$, $k_2 = 1.4$, $\phi_s = 0.4$, $L_r A_r = 1$, $k_L = 2 \times 10^{-3}$, and $0 < k_0 \leq 1$. Further, the restrictions (6.18), (6.19) and (6.23), (6.24), which ensure the uniqueness for arbitrary \mathbf{K} , hold when we choose $Da = 10^{-1}$, and $k_0 = 10^{-2}$ with the remaining parameters as chosen above. Note that the choice of parameters mentioned above may not be unique; there could be other parameter combinations for which these restrictions hold true.

7. Summary

In this work, we have modelled the poro-elasto-hydrodynamics that mimic an in-vitro solid tumour using biphasic mixture theory. We simplified the generic biphasic mixture equations using certain assumptions based on the biological context and treated hydraulic resistivity as anisotropic, which depends on the deformation. This made our model non-linear and coupled. We derive an equivalent variational (or weak) formulation and developed existence and uniqueness results using the Galerkin method, monotone operator theory and fixed-point theory. The detailed analysis is done by considering two cases: (a) $\mathbf{K}(\boldsymbol{\zeta}) = \mathbf{K}(\mathbf{U}_s)$ (b) $\mathbf{K}(\nabla \cdot \boldsymbol{\zeta}) = \mathbf{K}(\nabla \cdot \mathbf{U}_s)$. In both cases, we first developed existence and uniqueness analysis for auxiliary linear and semilinear sub-problems using the Galerkin method and monotone operator theory. Then, we convert the corresponding coupled non-linear problem to the fixed-point problem in both cases. Further, to develop the existence of solutions for the corresponding fixed-point problems,

we used the Schauder fixed-point theorem. Uniqueness is assured via the Banach contraction theorem. For the case where \mathbf{K} is not bounded but satisfies the sub-linear growth condition, we have developed the existence and uniqueness results. Moreover, we have collected certain realistic ranges of parameters involved in the model and ensured that the theoretical restrictions derived by us are compatible with these parameter ranges.

Acknowledgement. We are grateful for the valuable constructive suggestions provided by the anonymous referees, which greatly enhanced the quality of our paper. In particular, we greatly appreciate the suggestions on alternative methodologies to prove one of the main results. M. Alam, one of the authors, is thankful to Dr. Michael Eden and Dr. Debajyoti Choudhuri for their input regarding the analysis.

Competing interests. The authors declare that there is no conflict of interest.

References

- [1] Dey, B. & Raja Sekhar, G. P. (2016) Hydrodynamics and convection enhanced macromolecular fluid transport in soft biological tissues: Application to solid tumor. *J. Theor. Biol.* **395**, 62–86.
- [2] Byrne, H. (1999) Using mathematics to study solid tumour growth. In: *Proceedings of the 9th General Meetings of European Women in Mathematics*, New York, Hindawi Publishing, pp. 81–107.
- [3] Shelton, S. E. (2011) Mechanistic Modeling of Cancer Tumor Growth Using a Porous Media Approach, Master thesis, University of North Carolina.
- [4] Araujo, R. P. & McElwain, D. S. (2004) A history of the study of solid tumour growth: the contribution of mathematical modelling. *Bull. Math Biol.* **66**, 1039–1091.
- [5] Preziosi, L. & Tosin, A. (2009) Multiphase modelling of tumour growth and extracellular matrix interaction: mathematical tools and applications. *J. Math. Biol.* **58** (4-5), 625–656.
- [6] Please, C. P., Pettet, G. & McElwain, D. L. S. (1998) A new approach to modelling the formation of necrotic regions in tumours. *Appl. Math. Lett.* **11**, 89–94.
- [7] Ambrosi, D. & Preziosi, L. (2002) On the closure of mass balance models for tumor growth. *Math. Models Methods Appl. Sci.* **12**, 737–754.
- [8] Byrne, H. & Preziosi, L. (2003) Modelling solid tumour growth using the theory of mixtures. *Math. Med. Biol.* **20** (4), 341–366.
- [9] Sumets, P. P., Cater, J. E., Long, D. S. & Clarke, R. J. (2015) A boundary-integral representation for biphasic mixture theory, with application to the post-capillary glycocalyx. *Proc. R. Soc. A Math. Phys. Eng. Sci.* **471**, 20140955.
- [10] Barbeiro, S. & Wheeler, M. F. (2010) A priori error estimates for the numerical solution of a coupled geomechanics and reservoir flow model with stress-dependent permeability. *Comput. Geosci.* **14**, 755–768.
- [11] Jäger, W., Mikelić, A. & Neuss-Radu, M. (2011) Homogenization limit of a model system for interaction of flow, chemical reactions, and mechanics in cell tissues. *SIAM J. Math. Anal.* **43** (3), 1390–1435.
- [12] Cao, Y., Chen, S. & Meir, A. J. (2014) Steady flow in a deformable porous medium. *Math. Method Appl. Sci.* **37**, 1029–1041.
- [13] Alam, M., Dey, B. & Raja Sekhar, G. P. (2018) Mathematical analysis of hydrodynamics and tissue deformation inside an isolated solid tumor. *Theor. Appl.* **45** (2), 253–278.
- [14] Alam, M., Dey, B. & Raja Sekhar, G. P. (2019) Mathematical modeling and analysis of hydroelastodynamics inside a solid tumor containing deformable tissue. *ZAMM-J. Appl. Math. Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik* **e201800223** (5).
- [15] Alam, M., Byrne, H. & Raja Sekhar, G. P. (2021) Existence and uniqueness results on biphasic mixture model for an in-vivo tumor. *Appl. Anal.* **101** (15), 5442–5468.
- [16] Barry, S. & Aldis, G. (1990) Comparison of models for flow induced deformation of soft biological tissue. *J. Biomech.* **23** (7), 647–654.
- [17] Holmes, M. & Mow, V. (1990) The nonlinear characteristics of soft gels and hydrated connective tissues in ultrafiltration. *J. Biomech.* **23** (11), 1145–1156.
- [18] Ateshian, G. A. & Weiss, J. A. (2010) Anisotropic hydraulic permeability under finite deformation. *J. Biomech. Eng.* **132** (11), 111004.
- [19] Federico, S. & Herzog, W. (2008) On the anisotropy and inhomogeneity of permeability in articular cartilage. *Biomech. Model. Mechanobiol.* **7** (5), 367–378.
- [20] Giverso, C. & Preziosi, L. (2019) Influence of the mechanical properties of the necrotic core on the growth and remodelling of tumour spheroids. *Int. J. Nonlinear Mech.* **108**, 20–32.
- [21] Barry, S. I., Parkerf, K. H. & Aldis, G. K. (1991) Fluid flow over a thin deformable porous layer. *Zeitschrift für angewandte Mathematik und Physik ZAMP* **42** (5), 633–648.
- [22] Netti, P. A., Baxter, L. T., Boucher, Y., Skalak, R. & Jain, R. K. (1997) Macro-and microscopic fluid transport in living tissues: application to solid tumors. *AICHE J.* **43** (3), 818–834.
- [23] Wang, W. & Parker, K. H. (1995) The effect of deformable porous surface layers on the motion of a sphere in a narrow cylindrical tube. *J. Fluid Mech.* **283**, 287–305.

[24] Sun, S., Riviere, B. & Wheeler, M. F. (2002) A combined mixed finite element and discontinuous Galerkin method for miscible displacement problem in porous media. In: *Recent Progress in Computational and Applied PDEs*, Springer, pp. 323–351.

[25] Damiano, E. R., Duling, B. R., Ley, K. & Skalak, T. C. (1996) Axisymmetric pressure-driven flow of rigid pellets through a cylindrical tube lined with a deformable porous wall layer. *J. Fluid Mech.* **314**, 163–189.

[26] Gad-el Hak, M. (1995) Stokes hypothesis for a Newtonian, isotropic fluid. *J. Fluids Eng.* **117** (1), 3–5.

[27] Rajagopal, K. (2013) A new development and interpretation of the Navier-Stokes fluid which reveals why the Stokes assumption is inapt. *Int. J. Nonlinear Mech.* **50**, 141–151.

[28] Çeşmelioglu, A. & Rivière, B. (2012) Existence of a weak solution for the fully coupled Navier–Stokes/Darcy-transport problem. *J. Differ. Equations* **252** (7), 4138–4175.

[29] Vromans, A. J., van de Ven, A. A. F. & Muntean, A. (2019) Parameter delimitation of the weak solvability for a pseudo-parabolic system coupling chemical reactions, diffusion and momentum equations. *Adv. Math. Sci. Appl.* **28**, 273–311.

[30] Dey, B., Raja Sekhar, G. P. & Mukhopadhyay, S. K. (2018) In vivo mimicking model for solid tumor towards hydromechanics of tissue deformation and creation of necrosis. *J. Biol. Phys.* **44** (3), 361–400.

[31] Roose, T., Chapman, S. J. & Maini, P. K. (2007) Mathematical models of avascular tumor growth. *SIAM Rev.* **49**, 179–208.

[32] Islam, M., Tang, S., Liverani, C., Saha, S., Tasciotti, E. & Righetti, R. (2020) Non-invasive imaging of Young’s modulus and Poisson’s ratio in cancers in vivo. *Sci. Rep.* **10**, 1–12.

[33] Bernstein, B. & Toupin, R. (1960) Korn inequalities for the sphere and circle. *Arch. Ration. Mech. Anal.* **6** (1), 51–64.

[34] Salsa, S. (2016) *Partial Differential Equations in Action: From Modelling to Theory*, Springer, Vol. **99**.

[35] Attouch, H., Buttazzo, G. & Michaille, G. (2014). *Variational Analysis in Sobolev and BV Spaces: Applications to PDEs and Optimization*, SIAM.

[36] Temam, R. (2001) *Navier-Stokes Equations: Theory and Numerical Analysis*, American Mathematical Society, **343**.

[37] Zeidler, E. (2013) *Nonlinear Functional Analysis and Its Applications: II/B: Nonlinear Monotone Operators*, Springer Science & Business Media.

[38] Camano, J., Gatica, G. N., Oyarzúa, R. & Tierra, G. (2016) An augmented mixed finite element method for the Navier–Stokes equations with variable viscosity. *SIAM J. Numer. Anal.* **54** (2), 1069–1092.

[39] Caucao, S., Gatica, G. N., Oyarzúa, R. & Sánchez, N. (2020) A fully-mixed formulation for the steady double-diffusive convection system based upon Brinkman–Forchheimer equations. *J. Sci. Comput.* **85** (2), 37.

Appendix A

*Function spaces and useful results:*³ Let Ω be a bounded, open subset of \mathbb{R}^d , $\{d = 2, 3\}$. $L^2(\Omega)$ is the space of all measurable functions u defined on Ω for which

$$\|u\|_{0,\Omega} = \left(\int_{\Omega} |u|^2 d\Omega \right)^{1/2} < +\infty, \tag{A1}$$

In (A1), $\|\cdot\|_{0,\Omega}$ defines a norm on $L^2(\Omega)$. For any $\mathbf{u} = (u_1, u_2, \dots, u_d) \in L^2(\Omega)^d$, $\|\mathbf{u}\|_{0,\Omega}$ is defined as

$$\|\mathbf{u}\|_{0,\Omega} = \left(\int_{\Omega} \sum_{i=1}^d |u_i|^2 d\Omega \right)^{1/2}, \tag{A2}$$

and for any element $\mathbf{K} = (K_{ij})_{1 \leq i, j \leq d} \in (L^2(\Omega))^{d \times d}$, we define the norm of \mathbf{K} as

$$\|\mathbf{K}\|_{0,\Omega} = \left(\int_{\Omega} \sum_{i=1}^d \sum_{j=1}^d |K_{ij}|^2 d\Omega \right)^{1/2}. \tag{A3}$$

The symbols $(\cdot, \cdot)_{\Omega}$, and $(\cdot, \cdot)_{\partial\Omega}$ denote inner products in $L^2(\Omega)$, $L^2(\Omega)^d$, and $(L^2(\Omega))^{d \times d}$ and in the corresponding trace spaces $L^2(\partial\Omega)$, $L^2(\partial\Omega)^d$, and $(L^2(\partial\Omega))^{d \times d}$, respectively. For any two functions \mathbf{u} and \mathbf{v} , the inner products $(\cdot, \cdot)_{\Omega}$, and $(\cdot, \cdot)_{\partial\Omega}$ are defined as

$$(\mathbf{u}, \mathbf{v})_{\Omega} = \int_{\Omega} \mathbf{u} \cdot \mathbf{v} d\Omega, \quad (\mathbf{u}, \mathbf{v})_{\partial\Omega} = \int_{\partial\Omega} \mathbf{u} \cdot \mathbf{v} d\sigma.$$

The first-order Sobolev space is defined as

$H^1(\Omega)^d = \{\mathbf{u} \in L^2(\Omega)^d | \nabla \mathbf{u} \in (L^2(\Omega))^{d \times d}\}$ and the norm of a function $\mathbf{u} \in H^1(\Omega)^d$ is defined as

$$\|\mathbf{u}\|_{1,\Omega} = (\|\mathbf{u}\|_{0,\Omega}^2 + \|\nabla \mathbf{u}\|_{0,\Omega}^2)^{1/2}. \tag{A4}$$

³see [34] for function spaces and preliminaries.

$H_0^1(\Omega)^d$ denotes the space of functions in $H^1(\Omega)^d$ with zero trace. The dual space of $H_0^1(\Omega)^d$ is denoted by $H^{-1}(\Omega)^d$.

To show the existence of a solution, we rely on the following results.

Lemma A.1. (p.164 [36]) *Let \mathbb{X} be a finite-dimensional Hilbert space with scalar product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, and let G be a continuous mapping from \mathbb{X} into itself such that*

$$\langle G(x), x \rangle > 0 \text{ for } \|x\| = r_0 > 0.$$

Then there exists $x \in \mathbb{X}$, with $\|x\| \leq r_0$ such that

$$\langle G(x), x \rangle = 0.$$

Theorem A.2. (Schauder's (see p. 417 [34])) *Let X be a Banach space. Assume that:*

- (i) $A \subset X$ is closed and convex.
- (ii) $T : A \rightarrow A$ is continuous.
- (iii) $\overline{T(A)}$ is compact in X .

Then T has a fixed point $x^ \in A$.*

Theorem A.3. (Browder-Minty (see p. 557 [37])) *Let X be a real, separable, reflexive Banach space and let $T : X \rightarrow X^*$ (the dual of X) be bounded, continuous and strongly monotone. Then*

$$T(u) = g$$

has a unique solution for each $g \in X^$.*