

contribute to the provision of mental health and learning disabilities in an acute trust given the background of medical and mental health training. The impact of the strategy now needs to be assessed to ensure that it delivers improved services.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Improving Adherence to Prolactin Monitoring Guidelines in Acute Psychiatric Wards

Dr Hannah Campling, Dr Kimia Ghaderi and Dr Jayana Patel  
North London Foundation Trust, London, United Kingdom

doi: [10.1192/bjo.2025.10348](https://doi.org/10.1192/bjo.2025.10348)

**Aims:** The purpose of this project was to improve clinician's understanding and adherence to prolactin monitoring guidelines among doctors, reducing unnecessary testing and ensuring appropriate management of hyperprolactinaemia in patients on antipsychotic medications.

Prolactin monitoring in patients on antipsychotic medications is crucial for detecting and managing potential side effects. However, inconsistent adherence to monitoring guidelines can lead to missed diagnoses, unnecessary testing, and suboptimal patient management, in appropriate cessation of antipsychotics, or unnecessary addition of aripiprazole. This quality improvement project aimed to assess and improve clinicians' knowledge and measure the adherence to prolactin monitoring guidelines across three acute psychiatric wards in a North London Hospital.

**Methods:** A retrospective audit examining prolactin monitoring practices of patient records from June–July 2024 was conducted across three inpatient wards in a North London Hospital. Key metrics included frequency of symptom inquiry, appropriateness of testing, and adherence to management guidelines. Following the audit, an educational intervention was implemented at North London Mental Health Trust Academic programme, consisting of a presentation and pre/post-teaching surveys to assess knowledge improvement. Flowcharts summarising the guidelines were subsequently displayed in doctors' offices.

**Results:** The audit revealed low rates of symptom inquiry (7.4% in Sunflower, 6.25% in Tulip, 21.1% in Daisy) and high rates of unnecessary testing (44.4% in Sunflower, 81.25% in Tulip, 73.7% in Daisy) among patients on antipsychotics. Guideline adherence for managing raised prolactin levels was poor across all wards. The pre-teaching survey (n=37) demonstrated significant knowledge gaps, average 45% correct responses, particularly regarding age-specific monitoring and indications for testing. Post-intervention, a marked improvement in knowledge was observed across all domains in the post-teaching survey (n=16) with an average of 84% correct responses. For instance, correct responses regarding age-specific monitoring improved from 22% to 68.75% for women and from 11% to 81.25% for men.

**Conclusion:** This quality improvement project identified significant gaps in clinicians' knowledge and adherence to prolactin monitoring guidelines. The educational intervention demonstrated substantial improvements in clinicians' understanding of appropriate

monitoring practices. Ongoing efforts, including the display of guideline flowcharts and plans for reassessment, aim to sustain these improvements. Future work will focus on measuring long-term adherence to guidelines and its impact on patient outcomes and resource utilisation.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Management of Metabolic Health Syndrome: A Pilot Study in the Early Intervention Service

Dr Elishba Chacko and Dr Reem Abed  
SHSC, Sheffield, United Kingdom

doi: [10.1192/bjo.2025.10349](https://doi.org/10.1192/bjo.2025.10349)

**Aims:** Educating patients about metabolic side effects.

Empowering patients with knowledge and skills to make informed lifestyle choices.

Implementing personalised lifestyle interventions to improve metabolic health parameters.

Monitoring progress to facilitate long-term adherence to healthy behaviours.

**Methods:** Patients between 18–65 years.

Symptoms of metabolic syndrome i.e. high blood pressure, low HDL, truncal obesity, high triglycerides, impaired fasting glucose.

Patients currently/historically on antipsychotic medication.

Patients who have at least a year left in the service were included in the pilot.

Relative stability in mental health i.e. ability to engage with physical health appointments.

**Results:** The pilot concluded that patients benefited from tailored lifestyle interventions, giving them a sense of purpose and accountability.

There were significant changes in waist/circumference ratio; with noted improvement. Waist mean change = –5 cm (–6%); Waist: height mean change = –0.03 (–6%).

There were significant changes in weight: 5 individuals lost weight and improved their BMI; 2 individuals improved from overweight >normal.

There were no significant changes in biochemical markers.

A larger sample is required for a longer duration to study the impact of lifestyle interventions.

**Conclusion:** Small sample size – risk of bias, limited generalizability. Measurements for waist circumference might be prone to error as there is variation in the method of measurement i.e. over or under clothing. Blood results were inconclusive, perhaps the focus of the second phase should be waist circumference since there was marked change and measurable.

Clients would benefit from maintaining a food/mood diary and attending a briefing group at the start of the study to understand the basic principles of nutrition and the digestion process.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.