

RESEARCH ARTICLE

Log-concavity and relative log-concave ordering of compound distributions

Wanwan Xia¹  and Wenhua Lv²

¹School of Physical and Mathematical Sciences, Nanjing Tech University, Nanjing, Jiangsu, China

²School of Mathematical Sciences, Chuzhou University, Chuzhou, Anhui, China

Corresponding author: Wanwan Xia; Emails: 201910006533@njtech.edu.cn; whl@chzu.edu.cn

Keywords: Binomial distribution; Compound distribution; Entropy; Log-concavity; Negative binomial distribution

MSC: 60E05; 60E15

Abstract

In this paper, we compare the entropy of the original distribution and its corresponding compound distribution. Several results are established based on convex order and relative log-concave order. The necessary and sufficient condition for a compound distribution to be log-concave is also discussed, including compound geometric distribution, compound negative binomial distribution and compound binomial distribution.

1. Introduction

The entropy $H(X)$ of a random variable X measures the uncertainty of X . In this paper, we only consider discrete random variables. Let X be a discrete random variable with probability mass function (pmf) $\{x_1, \dots, x_n; p_1, \dots, p_n\}$, that is,

$$p_i = \mathbb{P}(X = x_i), \quad i = 1, \dots, n,$$

with $p_i \geq 0$, $i = 1, \dots, n$, and $\sum_{i=1}^n p_i = 1$. Here, n may be finite or infinite. The Shannon entropy of X is defined by [1]:

$$H(X) = - \sum_{i=1}^n p_i \log p_i.$$

The comparisons between distributions with respect to Shannon entropy are regarded as a measure of variability or dispersion. In insurance risk theory, similar comparisons are often established for compound distributions. The random variables corresponding to the compound distributions can be recorded as $S = \sum_{i=1}^M X_i$, which are extensively used in applied settings. For example, in [3], S can be used to model the total claim amount, M is the number of claims, and the X_i are the sizes of claims.

Our results are closely related to those of [8], who established entropy comparison results concerning compound distributions of random variables taking nonnegative integers based on convex ordering and log-concavity. We recall the following definitions. First, denote $\mathbb{N} = \{0, 1, 2, \dots\}$.

Definition 1.1. A sequence $\{h_n, n \in \mathbb{N}\}$ is said to be log-concave (LC) if $h_n \geq 0$ for $n \in \mathbb{N}$, and

$$h_n^2 \geq h_{n+1}h_{n-1}, \quad n \geq 1.$$

A log-concave sequence $\{h_n, n \in \mathbb{N}\}$ does not have internal zeros, i.e., there does not exist $i < j < k$ such that $h(i)h(k) \neq 0$ and $h(j) = 0$. A random variable X taking values in \mathbb{N} is said to be log-concave, if its pmf $\{f_n, n \in \mathbb{N}\}$ is log-concave.

Definition 1.2. For an integer $n \geq 2$, a positive sequence $\{h_i, 0 \leq i \leq n\}$ is called ultra-log-concave of order n (ULC(n)) if,

$$\frac{h_{i+1}^2}{\binom{n}{i+1}^2} \geq \frac{h_i}{\binom{n}{i}} \frac{h_{i+2}}{\binom{n}{i+2}}, \quad 0 \leq i \leq n - 2.$$

A random variable X taking values in \mathbb{N} is said to be ULC(n), if its pmf $\{f_i, 0 \leq i \leq n\}$ is ULC(n). Equivalently, X is ULC(n) if the sequence $\{f_i / \binom{n}{i}, 0 \leq i \leq n\}$ is log-concave.

Definition 1.3. A random variable X taking values in \mathbb{N} is said to be ultra-log-concave (ULC), if the support of X is an interval on \mathbb{N} , and its pmf $\{f_i, i \in \mathbb{N}\}$ satisfies:

$$(i + 1)f_{i+1}^2 \geq (i + 2)f_i f_{i+2}, \quad i \geq 0.$$

Equivalently, X is ULC if the sequence $\{i!f_i, i \in \mathbb{N}\}$ is log-concave.

In fact, both ULC(n) and ULC can be defined in terms of the relative log-concave order ([5]).

Definition 1.4. Let f and g be two pmfs on \mathbb{N} . Then f is relative log-concave to g , written as $f \leq_{lc} g$, if

- (1) the support of f and g are both intervals on \mathbb{N} ;
- (2) $\text{supp}(f) \subseteq \text{supp}(g)$;
- (3) f_i/g_i is log-concave on $i \in \text{supp}(f)$.

From the above definitions, we have $X \in \text{ULC}(n)$ is equivalent to $X \leq_{lc} B(n, p)$, and $X \in \text{ULC}$ is equivalent to $X \leq_{lc} \text{Poi}(\lambda)$, where $p \in (0, 1)$ and $\lambda > 0$. Also, we have the following inclusion relationship $\text{ULC}(1) \subseteq \text{ULC}(2) \subseteq \dots \subseteq \text{ULC} \subseteq \text{LC}$.

Definition 1.5. For random variables X and Y on \mathbb{N} , X is smaller than Y in the convex order, written as $X \leq_{cx} Y$, if $\mathbb{E}[\psi(X)] \leq \mathbb{E}[\psi(Y)]$ for all convex functions ψ on \mathbb{N} , provided the expectations exist.

The convex order compares the spread or variability of two distributions. Actually, if $X \leq_{cx} Y$ and both X and Y have finite means, we have $\mathbb{E}[X] = \mathbb{E}[Y]$ and $\text{Var}(X) \leq \text{Var}(Y)$. Further properties of the convex order can be found in [4]. Since entropy ordering also compares the variability of distributions, it is reasonable to expect some connection with convex ordering. For compound distributions, [7–9] draw the following conclusion.

Theorem 1.1 Suppose X and Y are two absolutely continuous or nonnegative integer-valued random variables.

- (1) [8],[9] If $X \leq_{lc} Y$ and $\mathbb{E}[X] = \mathbb{E}[Y]$, then $X \leq_{cx} Y$.
- (2) [7],[8] If $X \leq_{cx} Y$, and Y has a log-concave pdf or pmf, then $H(X) \leq H(Y)$.

Theorem 1.1 (2) provides a new method to prove entropy inequality. Compared with direct proof, it is relatively easier to establish convex ordering and to verify log-concave condition. Many conclusions in this paper involve **Theorem 1.1**.

The contributions and the outline of this paper are as follows.

- (i) In Section 2, based on Theorem 1.1, we give a direct proof for Lemma 3(2) in [6]. Our proofs are different since [6] constructed a Markov chain whose limiting distribution is a Binomial distribution.
- (ii) It is interesting to compare the original distribution and its corresponding compound distribution in the sense of the log-concave order. [8] considered the Poisson and Binomial distributions. Similar result is established in Section 3 for the Negative Binomial distribution.
- (iii) [8] obtained the necessary and sufficient conditions for a compound Poisson distribution to be log-concave. In Section 4, we establish necessary and sufficient conditions under which a compound Negative Binomial distribution or a compound Binomial distribution is log-concave.
- (iv) The preservation of convex order under compound operation was investigated by [8]. In Section 5, we consider whether the log-concave order is preserved under compound operation.

2. The entropy of ULC distribution

Suppose that \mathcal{F} is the set that contains all pmfs for nonnegative integer-valued random variables. Consider two operators S_p and T_n defined on \mathcal{F} [6]:

- (1) For all $p \in (0, 1)$, S_p maps pmf $f = \{f_i, i \in \mathbb{N}\}$ to another pmf $g = \{g_i, i \in \mathbb{N}\}$, where

$$g_i = pf_{i-1} + (1 - p)f_i, \quad i \geq 0,$$

and define $f_i = 0$ for all $i < 0$. Suppose that f is the pmf of X , $Z \sim B(1, p)$, independent of X , then $S_p f$ is the pmf of $X + Z$.

- (2) For all $n > 1$, T_n maps pmf $g = \{g_i, i = 0, \dots, n\}$ to another pmf $f = \{f_i, i = 0, \dots, n - 1\}$, where

$$f_i = \frac{n - i}{n}g_i + \frac{i + 1}{n}g_{i+1}, \quad i = 0, \dots, n - 1. \tag{2.1}$$

Denote g as the pmf of Y . Given Y , consider hypergeometric distribution $(n, Y, n - 1)$, that is, suppose there are n balls in an empty box, in which the number of white balls is Y . Now take $n - 1$ balls out of the box randomly without putting them back, and define random variable X as the number of white balls in the taken balls. Then $T_n g$ is the pmf of X .

Operators S_p and T_n satisfies [6]:

- (1) If $b_{n,p} = \{b(i, n, p), i = 0, \dots, n\}$ is the pmf of $B(n, p)$, then

$$S_p b_{n,p} = b_{n+1,p}, \quad T_{n+1} b_{n+1,p} = b_{n,p}, \quad n \geq 1.$$

So, $T_{n+1} \circ S_p b_{n,p} = b_{n,p}$.

- (2) $ULC(n)$ contains all the pmfs that are Ultra-log-concave of order n . Moreover, we have $S_p f \in ULC(n + 1)$ and $T_n f \in ULC(n - 1)$ for all $f \in ULC(n)$.

[6] proved that if X has pmf $f \in ULC(n)$ with $\mathbb{E}[X] = np, p \in (0, 1)$, and $Z \sim B(1, p)$, independent of X , and if Y is a hypergeometric distribution with parameters $(n + 1, X + Z, n)$, then $H(X) \leq H(Y)$. The proof uses the properties of operators S_p and T_n . By constructing a Markov chain whose limiting distribution is $B(n, p)$ and showing that the entropy never decreases along the iterations of this Markov chain. Here, we give another method to prove the non-decreasing property of the corresponding entropy based on Theorem 1.1.

Proposition 2.1. *Suppose that X has a pmf $f \in ULC(n)$, $\mathbb{E}[X] = np, p \in (0, 1)$, $Z \sim B(1, p)$, where Z is independent of X . Let Y be a hypergeometric distribution with parameters $(n + 1, X + Z, n)$. Then $X \leq_{cx} Y$.*

Proof. Define differential operation $\Delta h_i = h_i - h_{i-1}$ for any sequence $\{h_i\}$, and let $g = \{g_i, i = 0, \dots, n\}$ denote the pmf of Y . By (2.1), we have

$$g_i = \frac{n + 1 - i}{n + 1} (pf_{i-1} + qf_i) + \frac{i + 1}{n + 1} (pf_i + qf_{i+1}), \quad q = 1 - p. \tag{2.2}$$

Hence,

$$f_i - g_i = \frac{1}{n + 1} \Delta [p(n - i)f_i - q(i + 1)f_{i+1}] = \Delta(u_i h_i),$$

where

$$u_i = p - q \frac{(i + 1)f_{i+1}}{(n - i)f_i}, \quad h_i = \frac{(n - i)f_i}{n + 1}, \quad 0 \leq i < n,$$

and $u_i h_i = 0, i = -1, n$. Therefore, for any convex function sequence $\psi = \{\psi_i\}$, we have

$$\mathbb{E}[\psi(Y)] - \mathbb{E}[\psi(X)] = \sum_{i=0}^n \psi_i (g_i - f_i) = - \sum_{i=0}^n \psi_i \Delta(u_i h_i) = \sum_{i=0}^{n-1} (\psi_{i+1} - \psi_i) u_i h_i.$$

Since $h_i \geq 0$ and $\sum_{i=0}^n h_i = nq/(n + 1)$, h can be modified to a probability function. On the other hand, $f \in \text{ULC}(n)$ means that u_i is increasing in $i \in \{0, \dots, n - 1\}$; the convexity of ψ means that $\psi_{i+1} - \psi_i$ is increasing in i , and $\sum_{i=0}^{n-1} u_i h_i = 0$. Hence, by Chebyshev rearrangement theorem, we have $\sum_{i=0}^{n-1} (\psi_{i+1} - \psi_i) u_i h_i \geq 0$, that is, $\mathbb{E}[\psi(Y)] \geq \mathbb{E}[\psi(X)]$. □

Under the assumptions of Proposition 2.1, $Y \in \text{ULC}(n)$, and the corresponding pmf is log-concave. Hence, by Theorem 1.1, $X \leq_{\text{cx}} Y$ leads to $H(X) \leq H(Y)$.

3. Log-concave ordering between the original distribution and its corresponding compound distribution

Suppose that S is a nonnegative integer-valued random variable, defined by:

$$S = \sum_{i=1}^M X_i,$$

where $\{X_n, n \geq 1\}$ is a sequence of independent and identically distributed (iid) nonnegative integer-valued random variables, and M is a counting random variable independent of all X_i 's. Let f and h be the pmfs of X_i and M , respectively. The distribution of S is called a compound distribution with its pmf denoted by $c_h(f)$. If $M \sim \text{Poi}(\lambda), B(n, p)$ or $\text{NB}(\alpha, p)$, then the distribution of S is called compound Poisson distribution, compound Binomial distribution or compound Negative Binomial distribution. For $\alpha = 1$, $\text{NB}(1, p)$ reduces to the geometric distribution $\text{Geo}(p)$.

In the following, the pmfs of $\text{Poi}(\lambda), B(n, p), \text{NB}(\alpha, p)$ and $\text{Geo}(p)$ distributions are denoted by $\text{poi}(\lambda), b(n, p), \text{nb}(\alpha, p)$ and $\text{geo}(p)$, respectively. Similarly, the pmfs of the corresponding compound distributions are denoted by $c_{\text{poi}(\lambda)}(f), c_{b(n,p)}(f), c_{\text{nb}(\alpha,p)}(f)$ and $c_{\text{geo}(p)}(f)$. Denote by μ_f the mean of a distribution with pmf f . Then the means of the distributions with pmf $c_{\text{poi}(\lambda)}(f), c_{b(n,p)}(f), c_{\text{nb}(\alpha,p)}(f)$ are $\lambda\mu_f, n p \mu_f$ and $\alpha(1 - p)\mu_f/p$, respectively.

Proposition 3.1. ([8]) *Let f be a pmf on \mathbb{N} .*

- (1) *If the compound Poisson distribution $c_{\text{poi}(\lambda)}(f)$ is non-degenerate and log-concave, and $\lambda, \lambda^* > 0$, then $\text{poi}(\lambda^*) \leq_{\text{lc}} c_{\text{poi}(\lambda)}(f)$.*
- (2) *If the compound Binomial distribution $c_{\text{b}(n,p)}(f)$ is non-degenerate and log-concave, and $p, p^* \in (0, 1)$, then $\text{b}(n, p^*) \leq_{\text{lc}} c_{\text{b}(n,p)}(f)$.*

By Theorem 1.1 and Proposition 3.1, we have

$$\lambda^* = \lambda\mu_f \implies H(\text{poi}(\lambda^*)) \leq H(c_{\text{poi}(\lambda)}(f)), \tag{3.1}$$

$$p^* = p\mu_f \implies H(\text{b}(n, p^*)) \leq H(c_{\text{b}(n,p)}(f)). \tag{3.2}$$

Here, $\lambda^* = \lambda\mu_f$ ensures that two pmfs $\text{poi}(\lambda^*)$ and $c_{\text{poi}(\lambda)}(f)$ have the same mean. Similarly, $p^* = p\mu_f$ ensures that two pmfs $\text{b}(n, p^*)$ and $c_{\text{b}(n,p)}(f)$ have the same mean. Suppose that $M \sim \text{B}(n, p)$, $M^* \sim \text{B}(n, p^*)$, and $\{X_i, i \geq 1\}$ are iid random variables with a common pmf f . Assume that all random variables considered here and below are independent of each other. The explanation of (3.2) is as follows, and the explanation of (3.1) can be given similarly. We consider two following cases:

- (i) Suppose $\mu_f \geq 1$. Then $p^* \geq p$ since $p^* = p\mu_f$ in (3.2). Let $\{I_i, i \geq 1\}$ be a sequence of iid Bernoulli random variables with $\mathbb{P}(I_i = 1) = p/p^* \in (0, 1]$. Since $\sum_{i=1}^{M^*} I_i$ has the same distribution as M , it follows that the pmf of $\sum_{i=1}^{M^*} I_i X_i$ is $c_{\text{b}(n,p^*)}(\tilde{f}) = c_{\text{b}(n,p)}(f)$, where \tilde{f} is the pmf of $I_i X_i$, given by:

$$\tilde{f} = \frac{p}{p^*}f + \left(1 - \frac{p}{p^*}\right)\delta_0,$$

where δ_0 is the pmf of a degenerate random variable $Z=0$. Notice that $\mu_{\tilde{f}} = p\mu_f/p^* = 1$, and the uncertainty of $\sum_{i=1}^{M^*} I_i X_i$ is obviously stronger than that of M^* . Thus, (3.2) holds.

- (ii) Suppose $\mu_f \in (0, 1)$. In this case, $p^* < p$. Let $\{I_i, i \geq 1\}$ be a sequence of iid Bernoulli random variables with $\mathbb{P}(I_i = 1) = p^*/p \in (0, 1]$. Then $\sum_{i=1}^M I_i \sim \text{B}(n, p^*)$, and the pmf of $\sum_{i=1}^M X_i$ is $c_{\text{b}(n,p)}(f)$. Note that $\mathbb{E}[I_i] = \mathbb{E}[X_i] = p^*/p$ and $I_i \leq_{\text{cx}} X_i$ for each i . Thus, the uncertainty of $\sum_{i=1}^M X_i$ is obviously stronger than $\sum_{i=1}^M I_i$, that is (3.2).

To obtain the similar result for a compound Negative Binomial distribution, we need the recursive expression for the corresponding pmf.

Lemma 3.2. *Denote the pmf of $c_{\text{nb}(\alpha,p)}(f)$ as g . Then*

$$(n + 1)g_{n+1} = \frac{q}{1 - qf_0} \sum_{j=0}^n [(\alpha - 1)j + n + \alpha] f_{j+1} g_{n-j}, \quad n \geq 0. \tag{3.3}$$

Proof. In [3], it is assumed that $f_0 = 0$. We consider a more general situation, $f_0 \geq 0$. The pmf of $\text{NB}(\alpha, p)$ is denoted by $\{p_n, n \geq 0\}$, that is

$$p_n = \binom{\alpha + n - 1}{n} p^\alpha q^n, \quad n \geq 0, \quad q = 1 - p,$$

so

$$\frac{p_n}{p_{n-1}} = a + \frac{b}{n}, \quad n \geq 1,$$

where $a = q, b = (\alpha - 1)q$. Hence, for all $n \geq 0$,

$$\sum_{j=0}^{n+1} \left(a + \frac{bj}{n+1} \right) f_j g_{n+1-j} = qf_0 g_{n+1} + \frac{q}{n+1} \sum_{j=0}^n [(\alpha - 1)j + n + \alpha] f_{j+1} g_{n-j}. \tag{3.4}$$

Denote by $f^{(k)} = \{f_i^{(k)}, i \in \mathbb{N}\}$ the pmf of $\sum_{i=1}^k X_i$, where X_1, \dots, X_k are iid random variables with a common pmf f . On the other hand, by using $\sum_{j=0}^{n+1} j f_j f_{n+1-j}^{(k)} = \frac{n+1}{k+1} f_{n+1}^{(k+1)}$, we have

$$\begin{aligned} \sum_{j=0}^{n+1} \left(a + \frac{bj}{n+1} \right) f_j g_{n+1-j} &= \sum_{j=0}^{n+1} \left(a + \frac{bj}{n+1} \right) f_j \sum_{k=0}^{\infty} p_k f_{n+1-j}^{(k)} \\ &= \sum_{k=0}^{\infty} p_k \sum_{j=0}^{n+1} \left(a + \frac{bj}{n+1} \right) f_j f_{n+1-j}^{(k)} \\ &= \sum_{k=0}^{\infty} p_k \left(a f_{n+1}^{(k+1)} + \frac{b}{n+1} \sum_{j=0}^{n+1} j f_j f_{n+1-j}^{(k)} \right) \\ &= \sum_{k=0}^{\infty} p_k \left(a + \frac{b}{k+1} \right) f_{n+1}^{(k+1)} \\ &= \sum_{k=1}^{\infty} p_k f_{n+1}^{(k)} \\ &= p_0 f_{n+1}^{(0)} + \sum_{k=1}^{\infty} p_k f_{n+1}^{(k)} \quad [f_{\ell}^{(0)} = 0, \ell \geq 1] \\ &= g_{n+1}. \end{aligned} \tag{3.5}$$

By (3.4) and (3.5), we conclude (3.3). □

Proposition 3.3. *Let f be a pmf defined on \mathbb{N} . If $\alpha^* \geq \alpha \in (0, 1]$, $p, p^* \in (0, 1)$, and if the compound Negative Binomial distribution $c_{nb(\alpha,p)}(f)$ is non-degenerate and log-concave, then $nb(\alpha^*, p^*) \leq_{lc} c_{nb(\alpha,p)}(f)$.*

Proof. Denote $g = c_{nb(\alpha,p)}(f)$. Since g is non-degenerate and log-concave, we have $g_n > 0$ for $n \in \mathbb{N}$ and

$$\frac{g_{n-j}}{g_{n-j-1}} \geq \frac{g_n}{g_{n-1}}, \quad 0 < j < n. \tag{3.6}$$

Since $\alpha \in (0, 1)$ and $\alpha^* \geq \alpha$, it follows that:

$$\frac{(\alpha - 1)j + n + \alpha}{(\alpha - 1)j + n - 1 + \alpha} \geq \frac{n + \alpha}{n + \alpha - 1} \geq \frac{n + \alpha^*}{n + \alpha^* - 1}, \quad j \geq 1. \tag{3.7}$$

In view of (3.6) and (3.7), we have

$$\begin{aligned}
 (n + 1)g_{n+1} &\geq \frac{q}{1 - qf_0} \sum_{j=0}^n [(\alpha - 1)j + n + \alpha] f_{j+1} g_{n-j-1} \cdot \frac{g_n}{g_{n-1}} \\
 &\geq \frac{g_n}{g_{n-1}} \cdot \frac{q}{1 - qf_0} \sum_{j=0}^{n-1} [(\alpha - 1)j + n + \alpha] f_{j+1} g_{n-j-1} \\
 &\geq \frac{g_n}{g_{n-1}} \cdot \frac{q}{1 - qf_0} \sum_{j=0}^{n-1} [(\alpha - 1)j + n - 1 + \alpha] \frac{\alpha^* + n}{\alpha^* + n - 1} f_{j+1} g_{n-j-1} \\
 &\geq \frac{g_n(\alpha^* + n)}{g_{n-1}(\alpha^* + n - 1)} \cdot \frac{q}{1 - qf_0} \sum_{j=0}^{n-1} [(\alpha - 1)j + (n - 1) + \alpha] f_{j+1} g_{n-j-1} \\
 &= \frac{g_n(\alpha^* + n)}{g_{n-1}(\alpha^* + n - 1)} \cdot ng_n, \quad n \geq 1.
 \end{aligned}$$

The above inequality can be simplified to:

$$\left[\frac{g_n}{\binom{\alpha^* + n - 1}{n}} \right]^2 \leq \frac{g_{n-1}}{\binom{\alpha^* + n - 2}{n-1}} \cdot \frac{g_{n+1}}{\binom{\alpha^* + n}{n+1}}, \quad n \geq 1,$$

that is, $g_n / \binom{\alpha^* + n - 1}{n}$ is log-convex, so $\text{nb}(\alpha^*, p^*) \leq_{\text{lc}} g$. □

Corollary 3.4. *Suppose that f is a pmf defined on \mathbb{N} with the mean $\mu > 0$. For $\alpha \in (0, 1]$, if there exists $\alpha^* > \alpha$ and $p^* \in (0, 1)$ such that:*

$$\frac{\alpha^*(1 - p^*)}{p^*} = \frac{\alpha\mu(1 - p)}{p},$$

and $c_{\text{nb}(\alpha,p)}(f)$ is non-degenerate and log-concave, then $H(\text{nb}(\alpha^*, p^*)) \leq H(c_{\text{nb}(\alpha,p)}(f))$.

Proof. By Proposition 3.3, $\text{nb}(\alpha^*, p^*) \leq_{\text{lc}} c_{\text{nb}(\alpha,p)}(f)$. It is easy to verify that the means of $\text{nb}(\alpha^*, p^*)$ and $c_{\text{nb}(\alpha,p)}(f)$ are equal. Thus, the desired result follows from Theorem 1.1(1) directly. □

Proposition 3.5. *Suppose that $\alpha > 0$ and $p \in (0, 1)$. If $c_{\text{nb}(\alpha,p)}(f)$ is non-degenerate and log-concave, then $\text{poi}(\lambda) \leq_{\text{lc}} c_{\text{nb}(\alpha,p)}(f)$ for any $\lambda > 0$. In particular, when $\lambda^* = \alpha\mu(1 - p)/p$, we have $H(\text{poi}(\lambda^*)) \leq H(c_{\text{nb}(\alpha,p)}(f))$.*

Proof. The notations are the same as in the proof of Proposition 3.3. Obviously, $g_n > 0$ for $n \geq 0$ and by the log-concavity of g , we have

$$\frac{g_{n-j}}{g_{n+1-j}} \leq \frac{g_n}{g_{n+1}}, \quad j = 0, \dots, n.$$

Hence, for $n \geq 0$,

$$\begin{aligned}
 (n + 1)g_{n+1} &\leq \frac{q}{1 - qf_0} \sum_{j=0}^n [(\alpha - 1)j + n + \alpha] f_{j+1} g_{n+1-j} \cdot \frac{g_n}{g_{n+1}} \\
 &\leq \frac{g_n}{g_{n+1}} \cdot \frac{q}{1 - qf_0} \sum_{j=0}^{n+1} [(\alpha - 1)j + n + 1 + \alpha] f_{j+1} g_{n+1-j}
 \end{aligned}$$

$$= \frac{g_n}{g_{n+1}} \cdot (n + 2)g_{n+2},$$

that is, $n!g_n$ is log-concave in $n \in \mathbb{N}$. Thus, $\text{poi}(\lambda) \leq_{lc} c_{\text{nb}(\alpha,p)}(f)$. The rest can be derived directly from [Theorem 1.1](#). □

4. Log-concavity of a compound distribution

[8] proved that if f is log-concave, then $c_{\text{poi}(\lambda)}(f)$ is log-concave if and only if $\lambda f_1^2 \geq 2f_2$. In order to show that [Propositions 3.3, 3.5](#) and [Corollary 3.4](#) are meaningful, we need to investigate the log-concavity of compound Negative Binomial distribution. Firstly, we study the necessary and sufficient condition for a compound Geometric distribution to be log-concave.

Proposition 4.1. *Suppose f is a pmf defined on \mathbb{N} such that $f_1 > 0$. For $p \in (0, 1)$, we have $c_{\text{geo}(p)}(f) \in \text{LC}$ if and only if $f_k = 0$ for all $k \geq 2$.*

Proof. Denote the pmf of $\text{Geo}(p)$ as $\{p_n, n \geq 0\}$. Then the pmf of $c_{\text{geo}(p)}(f)$ is g . By [\(3.3\)](#), we have:

$$g_{n+1} = \eta \sum_{j=0}^n f_{j+1}g_{n-j}, \quad n \geq 0, \tag{4.1}$$

where $\eta = q/(1 - qf_0) > 0$.

First of all, observe that

- (1) $g_0 = p_0 + \sum_{n=1}^{\infty} p_n f_0^n = p + \sum_{n=1}^{\infty} p q^n f_0^n = p/(1 - qf_0) = p\eta/q > 0$;
- (2) $f_1 \neq 0 \implies g_n > 0$ for $n \geq 0$.

(\implies) Prove that $f_n = 0$ for all $n \geq 2$ by induction. For $n=2$, by [\(4.1\)](#), we have $g_1 = \eta f_1 g_0, g_2 = \eta[f_1 g_1 + f_2 g_0]$. Substituting in $g_1^2 \geq g_0 g_2$, we have $f_2 g_0 = 0$, that is, $f_2 = 0$ and $g_1^2 = g_0 g_2$.

Now assume that $f_n = 0$ for $n = 2, \dots, k$ and $g_{k-1}^2 = g_{k-2} g_k$. Notice that

$$g_{k-1} = \eta f_1 g_{k-2}, \quad g_k = \eta f_1 g_{k-1}, \quad g_{k+1} = \eta[f_1 g_k + f_{k+1} g_0].$$

Substituting in $g_k^2 \geq g_{k-1} g_{k+1}$, we have

$$f_1^2 g_{k-1}^2 \geq f_1^2 g_k g_{k-2} + f_1 f_{k+1} g_0 g_{k-2}.$$

So, $f_1 f_{k+1} g_0 g_{k-2} = 0$. Thus, $f_{k+1} = 0$ and in the meantime, $g_k^2 = g_{k-1} g_{k+1}$. The necessity is proved by induction.

(\impliedby) Suppose that $f_k = 0$ for all $k \geq 2$. By [\(4.1\)](#), we have

$$g_{n+1} = \eta f_1 g_n = (\eta f_1)^{n+1} g_0, \quad n \geq 0. \tag{4.2}$$

It is obvious that $g_n^2 = g_{n-1} g_{n+1}$ for $n \geq 1$ and, hence, $g \in \text{LC}$. □

Remark 4.2.

- (1) By [\(4.2\)](#), we have

$$c_{\text{geo}(p)}(f) \in \text{LC} \implies c_{\text{geo}(p)}(f) = \text{geo}\left(\frac{p}{1 - qf_0}\right).$$

(2) Assume that $f_n = 0$ for $n \geq 2$ with $f_1 > 0$. Then $g := c_{nb(\alpha,p)}(f) \in LC$ if and only if $\alpha \geq 1$. In fact, if $f_n = 0$ for $n \geq 2$, then (3.3) can be simplified to:

$$g_{n+1} = \frac{n + \alpha}{n + 1} \eta f_1 g_n, \quad n \geq 0.$$

It is easy to verify that

$$g_{n+1}^2 \geq g_n g_{n+2} \iff \frac{\alpha - 1}{n + 1} \geq \frac{\alpha - 1}{n + 2},$$

that is, $\alpha \geq 1$.

Proposition 4.3. Suppose that $f \in LC$, and denote $q = 1 - p$ and $\eta = q / (1 - qf_0)$. Then $g := c_{nb(\alpha,p)}(f) \in LC$ if and only if

$$(\alpha - 1)\eta f_1^2 \geq 2f_2. \tag{4.3}$$

The proof of Proposition 4.3 is postponed to Appendix A.1.

Remark 4.4. Suppose that $f \in LC$, $0 < \alpha \leq \alpha^*$ and $p \geq p^* > 0$. If $g := c_{nb(\alpha,p)}(f) \in LC$, then $c_{nb(\alpha^*,p^*)}(f) \in LC$.

Next, the necessary and sufficient condition for a compound Binomial distribution to be log-concave is discussed. To this end, we first give the recursive expression of its corresponding pmf.

Lemma 4.5. Denote $g := c_{b(n,p)}(f)$. Then the pmf $\{g_k, k \in \mathbb{N}\}$ has recursive expression as follows:

$$(k + 1)g_{k+1} = \delta \sum_{j=0}^k [(n + 1)j + n - k] f_{j+1} g_{k-j}, \quad k \geq 0, \tag{4.4}$$

where $\delta = p / (pf_0 + q)$ and $q = 1 - p$.

Proof. The proof of (4.4) is similar to (3.3) since

$$\frac{p_k}{p_{k-1}} = -\frac{p}{q} + \frac{(n + 1)p}{qk} = a + \frac{b}{k}, \quad \forall k \geq 1.$$

Here, (3.5) still holds,

$$\sum_{j=0}^{k+1} \left(a + \frac{bj}{k + 1} \right) f_j g_{k+1-j} = g_{k+1}, \quad k \geq 0, \tag{4.5}$$

and (3.4) is replaced by the following formula

$$\begin{aligned} & \sum_{j=0}^{k+1} \left(a + \frac{bj}{k + 1} \right) f_j g_{k+1-j} \\ &= -\frac{p}{q} f_0 g_{n+1} + \frac{p}{q(n + 1)} \sum_{j=0}^k [(n + 1)j + n - k] f_{j+1} g_{k-j}, \quad k \geq 0. \end{aligned} \tag{4.6}$$

Thus, (4.4) follows from (4.5) and (4.6). □

Proposition 4.6. Denote $g := c_{b(n,p)}(f)$, and let $f \in LC$. If $g \in LC$, then

$$(n + 1)\delta f_1^2 \geq 2f_2, \tag{4.7}$$

where $\delta = p/(pf_0 + q)$ and $q = 1 - p$. In particular, for $n = 1$, (4.7) is also the sufficient condition for $c_{b(1,p)}(f) \in LC$. But for $n \geq 2$, (4.7) is not the sufficient condition for $g \in LC$.

Proof. (1) By lemma 4.5, we have $g_1 = n\delta f_1 g_0$ and $g_2 = [(n - 1)\delta f_1 g_1 + 2n\delta f_2 g_0]/2$. In view of $g_1^2 \geq g_0 g_2$, we obtain (4.7).

(2) For $n = 1$, (4.7) holds, that is $\delta f_1^2 \geq f_2$. Based on the random expression of the random variable corresponding to g , we have $g_0 = q + pf_0$ and $g_k = pf_k$ for $k \geq 1$. To prove $g \in LC$, we only need to prove $g_1^2 \geq g_0 g_2$, that is, (4.7).

(3) Take a counterexample to illustrate: assume $n = 2$, and $c_{b(2,p)}(f) = c_{b(1,p)}(f) * c_{b(1,p)}(f)$. Denote $h = c_{b(1,p)}(f)$, by (2), we have $h_0 = q + pf_0$, $h_k = pf_k$, $k \geq 1$. Especially, take $p = 20/23$,

$$f = \left(\frac{1}{20}, \frac{1}{5}, \frac{3}{10}, \frac{9}{20}, 0, 0, \dots \right).$$

Then

$$h = \left(\frac{4}{23}, \frac{4}{23}, \frac{6}{23}, \frac{9}{23}, 0, 0, \dots \right).$$

Hence,

$$g = h * h = \left(g_0, g_1, g_2, \frac{120}{23^2}, \frac{108}{23^2}, \frac{108}{23^2}, g_6, \dots \right).$$

Obviously, $g_4^2 < g_3 g_5$, which means that g is not log-concave. On the other hand, $\delta = p/(pf_0 + q) = 5$, $f \in LC$, and $3\delta f_1^2 = 3/5 = 2f_2$, that is (4.7) holds. Therefore, (4.7) is not the sufficient condition for $g \in LC$. □

5. The relative log-concavity

Lemma 2 in [8] states that, for pmfs f, g, f^* and g^* defined on \mathbb{Z}_+ , if $f \leq_{cx} f^*$ and $g \leq_{cx} g^*$, then $c_g(f) \leq_{cx} c_{g^*}(f^*)$. On the other hand, $g \leq_{lc} \text{poi}(\lambda)$ for any $g \in ULC$ and $\lambda > 0$. Connected with Theorem 1.1, it is easy to establish the following proposition.

Proposition 5.1.

- (1) [8] If $g \in ULC$ and $\mu_g = \lambda$, then $c_g(f) \leq_{cx} c_{\text{poi}(\lambda)}(f)$.
- (2) [8],[2] If $g \in ULC$, $\mu_g = \lambda > 0$ and $c_{\text{poi}(\lambda)}(f) \in LC$, then $H(c_g(f)) \leq H(c_{\text{poi}(\lambda)}(f))$.

Notice that $g \in ULC(n) \iff g \leq_{lc} b(n, p)$ for all $p \in (0, 1)$, and that $g \in LC \iff g \leq_{lc} \text{geo}(\lambda)$ for all $\lambda > 0$. The following two propositions are easily established by Theorem 1.1.

Proposition 5.2.

- (1) If $g \in ULC(n)$ and $\mu_g = np$, then $c_g(f) \leq_{cx} c_{b(n,p)}(f)$.

(2) If $g \in \text{ULC}(n)$, $\mu_g = np$ and $c_{b(n,p)}(f) \in \text{LC}$, then $H(c_g(f)) \leq H(c_{b(n,p)}(f))$.

Proposition 5.3.

- (1) If $g \in \text{LC}$ and $\mu_g = (1 - \lambda)/\lambda$, then $c_g(f) \leq_{\text{cx}} c_{\text{geo}(\lambda)}(f)$.
- (2) If $g \in \text{LC}$, $\mu_g = (1 - \lambda)/\lambda$ and $c_{\text{geo}(\lambda)}(f) \in \text{LC}$, then $H(c_g(f)) \leq H(c_{\text{geo}(\lambda)}(f))$.

Naturally, the following problems will arise:
Question

- (1) If $g \in \text{ULC}$ and $\mu_g = \lambda$, is it correct $c_g(f) \leq_{\text{lc}} c_{\text{poi}(\lambda)}(f)$?
- (2) If $g \in \text{ULC}(n)$ and $\mu_g = np$, is it correct $c_g(f) \leq_{\text{lc}} c_{b(n,p)}(f)$?
- (3) If $g \in \text{LC}$ and $\mu_g = (1 - \lambda)/\lambda$, is it correct that $c_g(f) \leq_{\text{lc}} c_{\text{geo}(\lambda)}(f)$?

The following three counterexamples show that the assertions in Question 1 are negative.

Example 5.1. $g \in \text{ULC}$, $\mu_g = \lambda$ and $f \in \text{LC} \not\Rightarrow c_g(f) \leq_{\text{lc}} c_{\text{poi}(\lambda)}(f)$.

Suppose that $g = B(2, \lambda/2)$ and $0 < \lambda < 2$, f satisfies $f_j = 0$ for $j \geq 3$, and denote $s = c_g(f)$ and $t = c_{\text{poi}(\lambda)}(f)$. Then

$$\begin{aligned} s_0 &= \left(1 - \frac{\lambda}{2}\right)^2 + \lambda \left(1 - \frac{\lambda}{2}\right) f_0 + \frac{\lambda^2}{4} f_0^2, & s_1 &= \lambda \left(1 - \frac{\lambda}{2}\right) f_1 + \frac{\lambda^2}{2} f_0 f_1, \\ s_2 &= \lambda \left(1 - \frac{\lambda}{2}\right) f_2 + \frac{\lambda^2}{2} f_0 f_2 + \frac{\lambda^2}{4} f_1^2, & s_3 &= \frac{\lambda^2}{2} f_1 f_2, & s_4 &= \frac{\lambda^2}{4} f_2^2, \end{aligned}$$

and

$$\begin{aligned} t_0 &= e^{\lambda(f_0-1)}, & t_1 &= \lambda f_1 e^{\lambda(f_0-1)}, & t_2 &= \frac{\lambda}{2} (\lambda f_1^2 + 2f_2) e^{\lambda(f_0-1)}, \\ t_3 &= \lambda^2 f_1 \left(\frac{1}{6} \lambda f_1^2 + f_2\right) e^{\lambda(f_0-1)}, & t_4 &= \frac{\lambda^2}{4} \left(\frac{1}{6} \lambda^2 f_1^4 + 2\lambda f_1^2 f_2 + 2f_2^2\right) e^{\lambda(f_0-1)}. \end{aligned}$$

Especially, we take $\lambda = 1$ and $f = (f_0, f_1, f_2) = (1/3, 1/3, 1/3)$, that is, f is a discrete uniform distribution on $\{0, 1, 2\}$. Obviously, $f \in \text{ULC}$. By calculation,

$$\left(\frac{s_0}{t_0}, \frac{s_1}{t_1}, \frac{s_2}{t_2}, \frac{s_3}{t_3}, \frac{s_4}{t_4}\right) = \left(\frac{4}{9}, \frac{2}{3}, \frac{9}{14}, \frac{9}{19}, \frac{54}{145}\right) e^{2/3}.$$

It can be verified that

$$\left(\frac{s_1}{t_1}\right)^2 - \frac{s_0}{t_0} \frac{s_2}{t_2} > 0, \quad \left(\frac{s_3}{t_3}\right)^2 - \frac{s_2}{t_2} \frac{s_4}{t_4} = -0.015 < 0.$$

Hence, s_k/t_k is not log-concave, so $s \not\leq_{\text{lc}} t$.

Example 5.2. $g \in \text{LC}$, $\mu_g = (1 - \lambda)/\lambda$ and $f \in \text{LC} \not\Rightarrow c_g(f) \leq_{\text{lc}} c_{\text{geo}(\lambda)}(f)$.

Suppose that f and g satisfy $f_j = g_j = 0$ for $j \geq 3$, and denote $s = c_g(f)$ and $t = c_{\text{geo}(\lambda)}(f)$. Then

$$\begin{aligned} s_0 &= g_0 + g_1 f_0 + g_2 f_0^2, & s_1 &= g_1 f_1 + 2g_2 f_0 f_1, \\ s_2 &= g_1 f_2 + g_2 (2f_0 f_2 + f_1^2), & s_3 &= 2g_2 f_1 f_2, & s_4 &= g_2 f_2^2, \end{aligned}$$

and

$$t_1 = \eta f_1 t_0, \quad t_2 = \eta(f_1 t_1 + f_2 t_0), \quad t_3 = \eta(f_1 t_2 + f_2 t_1), \quad t_4 = \eta(f_1 t_3 + f_2 t_2),$$

with $\eta = (1 - \lambda)/[1 - (1 - \lambda)f_0]$. If we take $f = (f_0, f_1, f_2) = (1/3, 1/3, 1/3) \in \text{LC}$, then

$$g = (g_0, g_1, g_2) = \left(\frac{7\lambda - 3}{4\lambda}, \frac{1 - \lambda}{2\lambda}, \frac{1 - \lambda}{4\lambda} \right).$$

Given $\lambda = 40/77$, g is log-concave, that is, $g \leq_{\text{lc}} \text{Geo}(\lambda)$. It is easy to verify that

$$\left(\frac{s_3}{t_3} \right)^2 - \frac{s_2 s_4}{t_2 t_4} = -0.082 < 0,$$

hence, s_k/t_k is not log-concave, so $s \not\leq_{\text{lc}} t$.

Example 5.3. $g \in \text{ULC}(n)$, $\mu_g = np$, $p \in (0, 1)$ and $f \in \text{LC} \not\Rightarrow c_g(f) \leq_{\text{lc}} c_{\text{b}(n,p)}(f)$.

Suppose that $f = (f_0, f_1, f_2) = (1/3, 1/3, 1/3) \in \text{LC}$ and $g = (g_0, g_1, g_2) = (1 - 3p/2, p, p/2)$. If $p \in (1/2, 2/3)$, then $g \in \text{ULC}(2)$. Denote $s = c_g(f)$ and $t = c_{\text{b}(2,p)}(f)$. Then s_j and t_j can be calculated as in Example 5.2. Choose $p = 7/12$. Then

$$\left(\frac{s_3}{t_3} \right)^2 - \frac{s_2 s_4}{t_2 t_4} = -0.1728 < 0,$$

hence, s_k/t_k is not log-concave, so $s \not\leq_{\text{lc}} t$.

Acknowledgements. The authors are grateful to the Associate Editor and two anonymous referees for their comprehensive reviews of an earlier version of this paper.

Funding statement. W. Xia was supported by the Natural Science Foundation of Jiangsu Higher Education Institutions of China (No. 20KJB110014). W. Lv was supported by the Scientific Research Project of Chuzhou University (No. zrzj2021019) and by the Scientific Research Program for Universities in Anhui Province (No. 2023AH051573).

References

- [1] Cover, T.M. & Thomas, J.A. (2006). 2nd Edn. *Elements of information theory*. New York: Wiley-Interscience.
- [2] Johnson, O., Kontoyiannis, I. & Madiman, M. (2008). On the entropy and log-concavity of compound Poisson measures. *arXiv:0805.4112*.
- [3] Panjer, H.H. (1981). Recursive evaluation of a family of compound distributions. *Astin Bulletin* 12(1): 22–26.
- [4] Shaked, M. & Shanthikumar, J.G. (2007). *Stochastic orders*. New York: Springer.
- [5] Whitt, W. (1985). Uniform conditional variability ordering of probability distributions. *Journal of Applied Probability* 22(3): 619–633.
- [6] Yu, Y. (2008a). On the maximum entropy properties of the Binomial distribution. *IEEE Transactions on Information Theory* 54(7): 3351–3353.
- [7] Yu, Y. (2008b). On an inequality of Karlin and Rinott concerning weighted sums of i.i.d. random variables. *Advances in Applied Probability* 40(4): 1223–1226.
- [8] Yu, Y. (2009). On the entropy of compound distributions on nonnegative integers. *IEEE Transactions on Information Theory* 55(8): 3645–3650.
- [9] Yu, Y. (2010). Relative log-concavity and a pair of triangle inequalities. *Bernoulli* 16(2): 459–470.

Appendix A. Proof of Proposition 4.3

Proof. For reading convenience, use $P = g$ as pmf, and set $P_i = 0$ for $i < 0$. Denote $r_j = \eta f_{j+1}$, and rewrite (3.3) as:

$$(n + 1)P_{n+1} = \sum_{j=0}^n [(\alpha - 1)j + n + \alpha]r_j P_{n-j}, \quad \forall n \geq 0. \tag{A.1}$$

(\implies) By (A.1), we have $P_1 = \alpha r_0 P_0, P_2 = [(1 + \alpha)r_0 P_1 + 2\alpha r_1 P_0]/2$. Since $P \in LC$, it follows that $P_1^2 \geq P_0 P_1$, and thus,

$$P_1 \cdot \alpha r_0 P_0 \geq \frac{1}{2} [(1 + \alpha)r_0 P_1 + 2\alpha r_1 P_0] P_0.$$

So (4.3) can be proved directly by simplify the above equation.

(\impliedby) Suppose (4.3) holds. First, notice the following equations:

$$\begin{aligned} \alpha P_n P_{n+1} &= \sum_{j=0}^n [(\alpha - 1)j + n + \alpha](n + \alpha)r_j P_{n-j} P_n \\ &\quad - \sum_{j=0}^{n-1} [(\alpha - 1)j + n - 1 + \alpha](n + 1 + \alpha)r_j P_{n-1-j} P_{n+1}, \\ (n + 1)^2 P_{n+1}^2 &= \sum_{k=0}^n \sum_{l=0}^n [(\alpha - 1)k + n + \alpha][(\alpha - 1)l + n + \alpha]r_k r_l P_{n-k} P_{n-l}, \\ n(n + 2)P_{n+1}^2 &= (n + 1)^2 P_{n+1}^2 - P_{n+1}^2, \\ n(n + 2)P_n P_{n+2} &= \sum_{k=0}^{n-1} \sum_{l=0}^{n+1} [(\alpha - 1)k + n - 1 + \alpha][(\alpha - 1)l + n + 1 + \alpha] \times r_k r_l P_{n-1-k} P_{n+1-l}. \end{aligned}$$

The first equation follows from the fact that the right hand side equals $(n + \alpha)P_n \cdot (n + 1)P_{n+1} - (n + 1 + \alpha)P_{n+1} \cdot nP_n = \alpha P_n P_{n+1}$. Define

$$J_n^{(1)} = (\alpha r_0 P_n - P_{n+1})P_{n+1}. \tag{A.2}$$

Then,

$$\begin{aligned} &n(n + 2)[P_{n+1}^2 - P_n P_{n+2}] \\ &= (\alpha r_0 P_n P_{n+1} - P_{n+1}^2) + (n + 1)^2 P_{n+1}^2 - n(n + 2)P_n P_{n+2} - \alpha r_0 P_n P_{n+1} \\ &= J_n^{(1)} + \sum_{k=0}^n \sum_{l=0}^n [(\alpha - 1)k + n + \alpha][(\alpha - 1)l + n + \alpha]r_k r_l P_{n-k} P_{n-l} \\ &\quad - \sum_{k=0}^{n-1} \sum_{l=0}^{n+1} [(\alpha - 1)k + n - 1 + \alpha][(\alpha - 1)l + n + 1 + \alpha]r_k r_l P_{n-1-k} P_{n+1-l} \\ &\quad - r_0 \sum_{j=0}^n [(\alpha - 1)j + n + \alpha](n + \alpha)r_j P_{n-j} P_n \\ &\quad + r_0 \sum_{j=0}^{n-1} [(\alpha - 1)j + n - 1 + \alpha](n + 1 + \alpha)r_j P_{n-1-j} P_{n+1} \end{aligned}$$

$$\begin{aligned}
 &= J_n^{(1)} + \sum_{k=0}^n \sum_{l=1}^n [(\alpha - 1)k + n + \alpha][(\alpha - 1)l + n + \alpha]r_k r_l P_{n-k} P_{n-l} \\
 &\quad - \sum_{k=0}^{n-1} \sum_{l=1}^{n+1} [(\alpha - 1)k + n - 1 + \alpha][(\alpha - 1)l + n + 1 + \alpha]r_k r_l P_{n-1-k} P_{n+1-l} \\
 &= J_n^{(1)} + \sum_{k=0}^n \sum_{l=0}^n [(\alpha - 1)k + n + \alpha][(\alpha - 1)(l + 1) + n + \alpha]r_k r_{l+1} P_{n-k} P_{n-1-l} \\
 &\quad - \sum_{k=0}^n \sum_{l=0}^n [(\alpha - 1)k + n - 1 + \alpha][(\alpha - 1)(l + 1) + n + 1 + \alpha]r_k r_{l+1} P_{n-1-k} P_{n-l} \\
 &= J_n^{(1)} + \sum_{k=0}^n \sum_{l=0}^n [\alpha(k + 1) + n - k][\alpha(l + 2) + n - l - 1]r_k r_{l+1} P_{n-k} P_{n-1-l} \\
 &\quad - \sum_{k=0}^n \sum_{l=0}^n [\alpha(k + 1) + n - 1 - k][\alpha(l + 2) + n - l]r_k r_{l+1} P_{n-1-k} P_{n-l} \\
 &= J_n^{(1)} + J_n^{(2)} + J_n^{(3)} + J_n^{(4)},
 \end{aligned}$$

where $J_n^{(2)}, J_n^{(3)}$ and $J_n^{(4)}$ are defined by

$$\begin{aligned}
 J_n^{(2)} &= \alpha^2 \sum_{k=0}^n \sum_{l=0}^n (k + 1)(l + 2)r_k r_{l+1} [P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}] \\
 &= \alpha^2 \sum_{k \geq l} [P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}] [(k + 1)(l + 2)r_k r_{l+1} - (l + 1)(k + 2)r_l r_{k+1}], \\
 J_n^{(3)} &= \sum_{k=0}^n \sum_{l=0}^n [(n - k)(n - l - 1)P_{n-k} P_{n-1-l} - (n - 1 - k)(n - l)P_{n-1-k} P_{n-l}]r_k r_{l+1} \\
 &= \sum_{k \geq l} [(n - k)(n - l - 1)P_{n-k} P_{n-1-l} - (n - 1 - k)(n - l)P_{n-1-k} P_{n-l}](r_k r_{l+1} - r_l r_{k+1}), \\
 J_n^{(4)} &= \alpha \sum_{k=0}^n \sum_{l=0}^n r_k r_{l+1} [(k + 1)(n - l - 1) + (n - k)(l + 2)]P_{n-k} P_{n-1-l} \\
 &\quad - \alpha \sum_{k=0}^n \sum_{l=0}^n r_k r_{l+1} [(k + 1)(n - l) + (n - k - 1)(l + 2)]P_{n-1-k} P_{n-l}.
 \end{aligned}$$

Define function $h(k, l) = (k + 1)(n - l) + (n - k - 1)(l + 2)$, which satisfies $h(k, l) = h(l, k)$. Therefore, $J_n^{(4)}$ can be simplified to:

$$\begin{aligned}
 J_n^{(4)} &= \alpha \sum_{k=0}^n \sum_{l=0}^n r_k r_{l+1} [(h(k, l)) + l - k + 1]P_{n-k} P_{n-1-l} - h(k, l)P_{n-1-k} P_{n-l}] \\
 &= \alpha \sum_{k=0}^n \sum_{l=0}^n r_k r_{l+1} h(k, l)(P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}) \\
 &\quad + \alpha \sum_{k=0}^n \sum_{l=0}^n r_k r_{l+1} (l - k + 1)P_{n-k} P_{n-1-l} \\
 &= \alpha \sum_{k \geq l} h(k, l)(P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l})(r_k r_{l+1} - r_l r_{k+1})
 \end{aligned}$$

$$\begin{aligned}
 & + \alpha \sum_{k=0}^n \sum_{l=1}^{n+1} r_k r_l (l - k) P_{n-k} P_{n-l} \\
 & = \alpha \sum_{k \geq l} h(k, l) (P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}) (r_k r_{l+1} - r_l r_{k+1}) + \alpha \sum_{k=0}^n r_k r_0 k P_{n-k} P_n \\
 & \geq \alpha \sum_{k \geq l} h(k, l) (P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}) (r_k r_{l+1} - r_l r_{k+1}). \tag{A.3}
 \end{aligned}$$

Then, we prove the log-concavity of $\{P_n, n \geq 0\}$ by induction. For $k = 1$, it is the same as (4.3). Now assume $P_k^2 \geq P_{k-1} P_{k+1}$ for all $k \leq n$. To prove $P_{n+1}^2 \geq P_n P_{n+2}$, it suffices to prove:

$$J_n^{(1)} + J_n^{(2)} + J_n^{(3)} + J_n^{(4)} \geq 0.$$

In fact, $J_n^{(\nu)} \geq 0$ for all $\nu \in \{1, 2, 3, 4\}$. Details are as follows.

- $\nu = 1$: By the assumption of induction, we have $\alpha r_0 = P_1/P_0 \geq P_2/P_1 \geq P_{n+1}/P_n$ and, hence, $J_n^{(1)} \geq 0$ due to (A.2).
- $\nu = 2$: $f \in LC$ implies that $\{(i + 1)r_i, i \geq 0\}$ is also log-concave. Hence,

$$(k + 1)(l + 2)r_k r_{l+1} \geq (l + 1)(k + 2)r_l r_{k+1}, \quad k \geq l.$$

Furthermore, $P_{n-k} P_{n-1-l} \geq P_{n-1-k} P_{n-l}$ for $k \geq l$, which holds by the assumption of induction. So, $J_n^{(2)} \geq 0$.

- $\nu = 3$: The hypothesis means that P_k is log-concave in $k \in \{0, 1, \dots, n + 1\}$, implying that kP_k is log-concave in $k \in \{0, 1, \dots, n + 1\}$. Therefore,

$$(n - k)(n - l - 1)P_{n-k} P_{n-1-l} \geq (n - 1 - k)(n - l)P_{n-1-k} P_{n-l}, \quad k \geq l.$$

Obviously, $r_k r_{l+1} \geq r_l r_{k+1}$ for $k \geq l$. Thus, $J_n^{(3)} \geq 0$.

- $\nu = 4$: By the definition of $h(k, l)$, we have

$$h(k, l) \geq 0, \quad \forall k \leq n - 1; \quad h(n, n - 1) = 0; \quad h(n, l) > 0, \quad \forall l \leq n - 2.$$

Applying (A.3), we have

$$\begin{aligned}
 J_n^{(4)} & \geq \alpha \sum_{l \leq k \leq n-1} h(k, l) (P_{n-k} P_{n-1-l} - P_{n-1-k} P_{n-l}) (r_k r_{l+1} - r_l r_{k+1}) \\
 & \quad + \alpha \sum_{l=0}^n h(n, l) P_0 P_{n-1-l} (r_n r_{l+1} - r_l r_{n+1}) \geq 0.
 \end{aligned}$$

Based on the above discussion, the log-concavity of $\{P_n, n \geq 0\}$ is proved by induction. □