

Marine Record

Cite this article: Inteca G, Silva I and Farooq H (2025) The first record of green turtle (*Chelonia mydas*) nesting attempt in Pemba, Cabo Delgado, northern Mozambique. *Journal of the Marine Biological Association of the United Kingdom* **105**, e37, 1–5. <https://doi.org/10.1017/S0025315425000323>

Received: 26 June 2024
Revised: 17 February 2025
Accepted: 2 March 2025

Keywords:


beach; green sea-turtle; Mozambique; nest; Pemba; threats

Corresponding author: Gélica Inteca;
Email: gelica.inteca@unilurio.ac.mz

© The Author(s), 2025. Published by Cambridge University Press on behalf of Marine Biological Association of the United Kingdom. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



The first record of green turtle (*Chelonia mydas*) nesting attempt in Pemba, Cabo Delgado, northern Mozambique

Gélica Inteca¹ , Isabel Silva¹ and Harith Farooq^{1,2,3}

¹Faculty of Natural Sciences, Lúrio University, Pemba, Mozambique; ²Center for Macroecology, Evolution and Climate, Globe Institute, University of Copenhagen, Copenhagen, Denmark and ³Gothenburg Global Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden

Abstract

We document the first recorded nesting attempt by a green sea-turtle (*Chelonia mydas*) in Pemba located in the Cabo Delgado Province of northern Mozambique – an urban area with a population of 200,529 inhabitants. This sighting expands the known nesting range of sea-turtles by 40 km south of the Quirimbas Archipelago, highlighting Pemba's potential as a critical nesting site. In July 2021, local fishermen were observed by a tourism operator killing the turtle and selling its meat. Consequently, the absence of eggs at the presumed nesting site suggests that the turtle was intercepted before it could lay eggs. This incident highlights the threats to sea-turtles and the need for conservation while emphasising the importance of collaboration among communities, government, and researchers to protect nesting activity.

Introduction

Five species of sea-turtles are found in Africa: the hawksbill (*Eretmochelys imbricata*), loggerhead (*Caretta caretta*), leatherback (*Dermochelys coriacea*), green (*Chelonia mydas*), and olive ridley (*Lepidochelys olivacea*). According to the International Union for Conservation of Nature's Red List (IUCN, 2023), all of these species are listed as threatened, with conservation statuses ranging from Vulnerable to Critically Endangered. The East Coast of Africa plays a critical role for global marine turtle conservation, providing essential nesting, migratory, and foraging habitats for both juvenile and adult turtles in the Western Indian Ocean region (van de Geer et al., 2022). Furthermore, sea-turtles significantly contribute to the livelihoods and socio-economic development of coastal communities, though the full scope of their impact is yet to be fully documented (van de Geer et al., 2022).

Of the five sea-turtle species that occur along the African Coast, all nest in Mozambique, but the green (*Chelonia mydas*, Linnaeus, 1758) and hawksbill (*Eretmochelys imbricata*, Fitzinger 1843) species are the only ones documented to nest in the northern region (Farooq et al., 2022; Fernandes et al., 2017; Louro et al., 2006, 2011; Pereira, 2006; van de Geer et al., 2022). The green sea-turtle has major nesting sites in Mozambique, primarily located in the archipelagos of Quirimbas (Anastácio et al., 2014; Garnier et al., 2012; van de Geer et al., 2022), Primeiras e Segundas, and Bazaruto (Fernandes et al., 2017; Pilcher and Williams, 2019), all of which are protected areas. To the best of our knowledge, nesting outside these areas in the country has not been documented so far. The threats affecting sea-turtles in Mozambique include hunting (Fernandes et al., 2017; Pilcher and Williams, 2019), nest foraging (Anastácio et al., 2014), and pollution (Inteca et al., 2023). Evidence of sea-turtle mortality, such as the discovery of carapaces and bones on beaches and the sale of products made from turtle scutes, has also been observed in local markets across the country (MICOA, 2012; Pilcher and Williams, 2019). Additionally, it has been suggested, although not officially confirmed, that local communities may consume sea-turtle eggs (van de Geer et al., 2022). Reports of sea-turtle mortality and in-water occurrence in Pemba – a city located in the northernmost province of Mozambique, have been documented since 1974 with more recent sightings in 2018 and 2022 (Farooq et al., 2022; Hughes, 1974; Pilcher and Williams, 2019). Despite these reports, no nesting activity was ever documented from Pemba. Thus, our observation is the first recorded nesting attempt by a green sea-turtle (*Chelonia mydas*) in Pemba located in the Cabo Delgado Province of northern Mozambique – an urban area with a population of 200,529 inhabitants (INE, 2019). Given that Pemba's beaches are popular among tourists, this situation presents a unique opportunity for local tourism operators to engage in conservation efforts. Our observation extends the known nesting range of sea-turtles in the northern part of the country by 40 km south of the Quirimbas Archipelago.




Figure 1. Maps depicting the location and context of green turtle nesting in northern Mozambique. (A) a detailed map of Cabo Delgado Province showing known green turtle nesting sites (blue circles) (Vamizi Island [11.01° S, 40.72° E], Quirimbas National Park [12.15° S, 40.56° E], Sencar Island [12.48° S, 40.64° E], Rolas Island [12.15° S, 40.56° E], Matemo Island [12.19° S, 40.60° E], Ibo Island [12.33° S, 40.58° E], Mefunvo Island [12.55° S, 40.60° E], Quilaleia Island [12.53° S, 40.60° E], Mucojo Beach [12.07° S, 40.48° E], Namau Beach [12.78° S, 40.56° E], Guludo Beach [12.13° S, 40.48° E], Naunde Beach [12.17° S, 40.48° E]) (Costa, 2007; Fernandes et al., 2020; Van de geer et al., 2022), and the location of Pemba (red rectangle), from where we report the current nesting attempt occurred. (B) A map of Africa focusing on Mozambique for regional context. (C) A map of Mozambique highlighting Cabo Delgado Province in the northern part of the country.

This highlights the potential significance of Pemba, an urbanised coastal city outside protected areas, as a nesting site.

Observation site

On July 20, 2021, our team members were approached by a local tourist operator who observed fishermen selling sea-turtle meat in baskets on Maringanhas' Beach in the northwest part of Pemba, Cabo Delgado Province, northern Mozambique, Southern

Africa (Figure 1A–C). The operator had witnessed the turtle being killed on-site shortly beforehand. Our team promptly visited the location, measured the turtle's carapace (Figure 2A) and tracks (Figure 2B and C), and identified the species. A total of 57 days later, on September 15, 2021, we revisited the site to look for signs of nesting but no signs were found.

Maringanha Beach (12.9745° S, 40.5872° E) is characterised by its vast sandy expanse, typical coastal vegetation, low step dunes (<1 m) (Figure 2D), a few scattered houses, and absence of street




Figure 2. Photographs taken on the beach where the observation occurred. (A) Carapace of the green turtle; (B) faeces and blood from the green turtle; (C) tracks of the green turtle; (D) the place where the female was trying to dig the nest; photos taken by António Brito (tourist operator) on July 21, 2021. Due to the fishermen's concerns regarding the involvement of the authorities, and to avoid jeopardising our opportunity to gather information, we were given only a brief window to photograph the specimen.




Figure 3. A map depicting the location and environmental context of Maringanha in Pemba, Cabo Delgado, showing the intensity of night lights in Pemba on July 21, 2021, highlighting the low light pollution in Maringanha (log10-transformed data from the VIIRS-DNB global cloud-free composites (Eogdata, 2021)).

lighting. The area predominantly supports artisanal fishing, which is a primary source of income for the local community.

Results and discussion

Identifying turtles by colour can be challenging due to their high colour variability. Instead, a reliable identification method involves examining the shape of the head, bill, body, and carapace. We identified the specimen as a green sea-turtle (*Chelonia mydas*) based on the number of scutes on the carapace (Figure 2A), and the composition of the turtle's faeces, which consisted of sea-grass (Figure 2B) (Eckert et al., 1999). This diet is consistent with adult green sea-turtles occurring at warmer sites with temperatures above 25°C, where Pemba is located (Esteban et al., 2020).

The carapace measured 87 cm in curved carapace width and 94 cm in curved carapace length. Additionally, the symmetrical tracks with parallel flipper marks measured 86 cm in width (Figure 2C). These measurements place the individual within the lower size range for adult female green sea-turtles, which typically measure between 90 and 120 cm in length (Bourjea et al., 2007). No eggs were discovered at the presumed nesting site, suggesting that the turtle was intercepted before laying eggs.

The nearest known nesting areas for green sea-turtles are approximately 40–60 km north, including both mainland and islands within Quirimbas National Park, such as Ibo Island, Quilalea Island, Sencar Island, Rolas Island, Muchojo Beach, Namau Beach, and Naunde Beach, Figure 1 (Costa, 2007).

This observation represents the southernmost documented nesting attempt in northern Mozambique, expanding the known nesting range of the species beyond the protected Quirimbas Archipelago to include one of the most densely populated cities in the northern part of the country. While green sea-turtles typically prefer nesting in secluded locations (Hendrickson, 1958; Medeiros et al., 2022), exceptions have been recorded in the southern mainland of Tanzania, as well as in Kiunga, Watamu, and Mombasa in Kenya, and along the north coast of Somalia (van de Geer et al., 2022). But despite the high population density of Pemba city in Mozambique, the recent nesting attempt at Maringanha Beach suggests that the habitat may still be suitable for sea-turtle nesting.

The relatively low level of artificial lighting at this site (Figure 3) and the presence of a wide sandy shoreline may have contributed to its viability as a potential nesting location. Artificial light pollution is a well-known factor that disrupts sea-turtle nesting behaviour by deterring females from emerging and disorienting hatchlings (Martins et al., 2017; Witherington, 1992). The low lighting incidence on the beach – approximately 2.72 Digital Number (DN) on the day it attempted to nest (Eogdata, 2021) – suggests that light pollution may not have been a major deterrent in this case. Values of DN range from 0 to 63, with 3 and 5.5 serving as thresholds for moderate and high DN, respectively. Although sea-turtles have been observed to nest at light levels up to 5.5 DN, most nesting activity typically occurs at DN levels of 3 or below (Colman et al., 2020).

Beyond light pollution, other environmental factors may influence nesting suitability in Pemba. The City is known for its suitable beach morphology, featuring a wide, sandy shoreline and a low-lying dune system ideal for sea-turtle nesting (Caldas and Rita, 2014). However, due to the high human activity in the area, sea-turtle nests are likely to be uncovered and the eggs consumed by local communities (Pilcher and Williams, 2019; van de Geer et al., 2022). Green sea-turtles in Mozambique are also threatened by

beach development (Anastácio et al., 2014), by-catch from trawling and artisanal fishing (Pilcher and Williams, 2019), and marine litter (Inteca et al., 2023). Despite these challenges, our observation suggests that Pemba could be a viable nesting site for green sea-turtles, provided there is adequate enforcement and monitoring. This incident highlights the ongoing threats to sea-turtle populations in the region and the urgent need for conservation measures. The fact that it was a local tourism operator that witnessed and reported the event showcases the potential role of tourism stakeholders in collaboration with local communities, government, and researchers in wild life monitoring and conservation efforts. Our findings emphasise the importance of enhanced protection for coastal and marine ecosystems, which are essential for biodiversity and may also support sustainable ecotourism initiatives.

Acknowledgements. We acknowledge António Brito, the touristic operator who, after seeing the group of fishermen selling sea-turtle meat and tracks on the site, immediately communicated with us. Harith Farooq was supported by The Danish Independent Research Council (grant no. 0165-00018B).

Author contributions. Gélica Inteca: Data collection, Writing – Original draft preparation and Editing, Isabel Silva: Reviewing, Harith Farooq: Editing, Writing and Reviewing.

Competing interests. The authors declare that they have no conflict of interest.

References

- Anastácio R, Santos C, Lopes C, Moreira H, Souto L, Ferrão J, Garnier J and Pereira MJ (2014) Reproductive biology and genetic diversity of the green turtle (*Chelonia mydas*) in Vamizi Island, Mozambique. *SpringerPlus* 3(1), 1–16.
- Bourjea J, Frappier J, Quillard M, Ciccione S, Roos D, Hughes G and Grizel H (2007) Mayotte Island: Another important green turtle nesting site in the southwest Indian Ocean. *Endangered Species Research* 3, 273–282.
- Caldas P and Rita A (2014) *Habitat selection and climate change impacts on green turtles from Poilão Island, Guinea-Bissau*, 14.
- Colman LP, Lara PH, Bennie J, Broderick AC, de Freitas JR, Marcondes A, Witt MJ and Godley BJ (2020) Assessing coastal artificial light and potential exposure of wildlife at a national scale: The case of marine turtles in Brazil. *Biodiversity and Conservation* 29(4), 1135–1152.
- Costa A (2007) *Status and Management of Marine Turtles in the Quirimbas National Park. A Report Prepared for the Quirimbas National Park*. WWF Mozambique Coordination Office. Marine Programme-Marine Turtle Tagging and Monitoring Programme. Maputo. January 2007, 1–13.
- Eckert KL, Bjørndal KA, Abreu-Grobois FA, Donnelly M, Schroeder B and Murphy S (1999) Research and management techniques for the conservation of sea turtles. *Population Surveys (Ground and Aerial) on Nesting Beaches* 301(4), 713–1401.
- Eogdata. (2021) *Index of/nighttime_light/monthly_notile/v10/2021/202107/vcm-cfg*. https://eogdata.mines.edu/nighttime_light/monthly_notile/v10/2021/202107/vcm-cfg/ (accessed 12 October 2023).
- Esteban N, Mortimer JA, Stokes HJ, Laloë JO, Unsworth RKF and Hays GC (2020) A global review of green turtle diet: Sea surface temperature as a potential driver of omnivory levels. *Marine Biology* 167(12), 1–17. (Springer Science and Business Media Deutschland GmbH).
- Farooq H, Nanvonamquitxo C, Nassongole B, Conradie W, Bills R, Soares A and Antonelli A (2022) Shedding light on a biodiversity dark spot: Survey of amphibians and reptiles of Pemba region in northern Mozambique. *Herpetological Conservation and Biology* 17(2), 423–432.
- Fernandes RS, Inteca GE, Williams JL, Taju A, Muaves I and Pereira MA (2020) *Monitoring, Tagging and Conservation of Marine Turtles in Mozambique: Annual Report 2018/19* (Issue January). doi:10.13140/RG.2.2.19667.73764

- Fernandes RS, Williams JL, Valladolid SG, Muaves L, Louro CMM and Pereira MAM** (2017) Monitoring, tagging and conservation of marine turtles in Mozambique: Annual Report 2016/17. *Ctv I* (September), 1–30.
- Garnier J, Hill N, Guissamulo A, Silva I, Witt M and Godley B** (2012) Status and community-based conservation of marine turtles in the northern Querimbas Islands (Mozambique). *Oryx* **46**(3), 359–367.
- Hendrickson RJ** (1958) The Green Sea Turtle. *Chelonia Mydas* (Linn.) In Malaya and Sarawak Department of Zoology. University of Malaya. Singapore chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/http://seaturtle.org/library/HendricksonJR_1958_ProcZoolSocLond.pdf (accessed 11 August 2021).
- Hughes G** (1974) *The sea turtles of South-East Africa*. 1–96. <http://www.ori.org.za/uploads/files/oriinvrep36.pdf> (accessed 10 August 2021).
- INE.** (2019) Resultados definitivos Censo 2017. IV Recenseamento Geral da População e Habitação de Moçambique. [Www.ine.gov.mz/iv-rgph-2017/mocambique/censo-2017-br-ochura-dos-resultados-definitivos-do-iv-rgph-nacional.pdf/view](http://www.ine.gov.mz/iv-rgph-2017/mocambique/censo-2017-br-ochura-dos-resultados-definitivos-do-iv-rgph-nacional.pdf/view) (accessed 28 September 2021).
- Inteca G, Hagy B, Silva I, Amoda C, Cululo A and Farooq H** (2023) The tourism industry keeps beaches clean in Mozambique. *Marine Pollution Bulletin*, **196**, 115615.
- IUCN.** (2023) *IUCN Red List of Threatened Species*. <https://www.iucnredlist.org/> (accessed 17 May 2023).
- Louro CMM, Videira E, Pereira M and Fernandes R** (2011) *Monitoring, Tagging and Conservation of Marine Turtles in Mozambique: Annual Report 2011/12. July*.
- Louro C, Pereira M and Costa A** (2006) *Report on the Conservation Status of Marine Turtles in Mozambique* (Issue January).
- Martins KA, Souza Pereira PD, Silva-Casarin R and Nogueira Neto AV** (2017) The influence of climate change on coastal erosion vulnerability in northeast Brazil. *Coastal Engineering Journal* **59**(2), 1740007–1.
- Medeiros L, Chaloupka M and Bolten AB** (2022) Tracking green turtle nesting trends at a remote oceanic rookery. *Marine Biology* **169**(5), 68.
- MICOA.** (2012) *República de Moçambique Ministério Para a Coordenação da Acção Ambiental. Centro de Desenvolvimento Sustentável para as Zonas Costeiras Tartarugas Marinhas Importância Social, Económica e Ecológica. Maputo. 2012.* 1.
- Pereira MAM** (2006) National Overview and Assessment on Marine Litter Related Activities: Mozambique. The Report Submitted to the Western Indian Ocean Marine Science Association (WIOMSA), 1–13.
- Pilcher NJ and Williams J** (2019) Status, scope and trends of the legal and illegal international trade in marine turtles, its conservation impacts, management options and mitigation priorities. *Eighteenth Meeting of the CITES Conference of the Parties 18* (January), 106.
- van de Geer CH, Bourjea J, Broderick AC, Dalleau M, Fernandes RS, Harris LR, Inteca GE, Kiponda FK, Louro CMM, Mortimer JA, Msangameno D, Mwasi LD, Nel R, Okemwa GM, Olendo M, Pereira MAM, Rees AF, Silva I, Singh S and Godley BJ** (2022) Marine turtles of the African east coast: Current knowledge and priorities for conservation and research. *Endangered Species Research* **47**, 297–331.
- Witherington BE** (1992) Herpetologists' league behavioral responses of nesting sea turtles to artificial lighting. *Herpetologica* **48**(1), 31–39.