

P01-319

THERAPEUTIC EFFECT OF RISPERIDONA INJECTABLE OF LONG EVOLUTION IN TEENAGERS

D. López Marco, R. Consuegra Sánchez, I. Martínez Pérez, A. Gómez Poveda

Unit Mental Health Infanto Juvenil, Center of Mental Health Cartagena, Murcia, Spain

The pharmacological treatment with oral Risperidona is used for years in infanto-juvenile psychiatry. In many occasions difficulty exists for the fulfillment; well for the appearance of adverse effects or for rejection to the capture or for difficulty of familiar fulfillment.

Aims: To expose from the clinical and welfare expectation, the experience with injectable Risperidona of long duration with three cases clinical whose principal diagnosis is of Mental Delay Moderated with alteration of the associate behavior.

Material and methods:

Case 1: 14-year-old Teenager of age attended in our CSM for Mental Delay and Disorder of conduct to familiar and school level.

CI=50. It has received diverse antipsychotic and stabilizing treatments of oral form being the discontinuous capture. For one year in treatment with injectable Risperidona 50mgr/14 days, answering favorably to the treatment, diminishing the conducts heteroagresivas that he was presenting to the beginning.

Case 2: 15-year-old Teenager of age late Mental and disorder of the disturbing behavior with episodes of excitation, irritability and loss of control.

Not clinical stabilization with oral treatment for lack of familiar adherence. We introduce injectable Risperidona, 7,5mgr/14 days, pupil to the family to mint of the importance of the treatment and the facility of administration. It improved the adherence to the center and therefore, clinical improvement.

Case 3: Preadolescente and 13 years and age with compatible symptoms late Mental Moderates do and autistic features with important estereotipias and autoagresions.

For a few months in treatment with Risperidona inyectable 37,5mgr/14 days with clinical improvement.