

Comparison of enteric protozoan infections in four Australian hospitals: variable tests and variable results

STEPHANIE M. FLETCHER-LARTEY*¹, DAVID ANDRESEN²,
SEBASTIAN VAN HAL³, JUAN MERIF⁴, DAMIEN STARK⁵,
WILLIAM RAWLINSON^{4,6}, JOHN HARKNESS⁵ and JOHN ELLIS⁷

¹ Public Health Unit, South Western Sydney Local Health District, PO Box 38, Liverpool, NSW 1871, Australia

² St. Vincent's Hospital, 390 Victoria Street, Darlinghurst, NSW 2010, Australia

³ Department of Microbiology and Infectious Diseases, Royal Prince Alfred Hospital, Camperdown, NSW 2050, Australia

⁴ Microbiology Department, South Eastern Area Laboratory Service (SEALS), Prince of Wales Hospital, Randwick, NSW, Australia

⁵ Division of Microbiology, SydPath, St. Vincent's Hospital, 390 Victoria Street, Darlinghurst, NSW 2010, Australia

⁶ School of Medical Sciences, The University of New South Wales, Sydney, Randwick, NSW 2031, Australia

⁷ School of Life Sciences, University of Technology, Sydney, P.O. Box 123, Broadway, NSW, Australia

(Received 2 November 2015; revised 19 May 2016; accepted 20 May 2016)

SUMMARY

There is limited evidence of the prevalence of enteric protozoan infections in developed settings. We estimated the prevalence of enteric protozoa and evaluated the outcome of testing algorithms used in hospital settings in Sydney, Australia. This retrospective study assessed microbiological data from four public clinical laboratories. Pooled data from the four hospitals revealed the most common enteric protozoan detected was *Blastocystis* spp. in an average of 5.4% of cases, followed by *Giardia intestinalis* (1.1%) and *Dientamoeba fragilis* (0.8%). Protozoan detection rates between hospitals were significantly different and could be based on multiple factors. The modified iron haematoxylin staining method, consistently detected higher rates of *Blastocystis* spp., and *G. intestinalis* in comparison with microscopy of wet preparations, as well as higher rates of *G. intestinalis* and *Cryptosporidium* when compared with enzyme immunoassay. The study concludes that there are multiple factors that contribute to the variability in protozoa detection rates in metropolitan hospitals, including widespread variability in the testing protocols for enteric protozoa, individual and population characteristics. A gold standard approach for diagnosis of enteric protozoa is recommended. Molecular diagnostic methods such as polymerase chain reaction would provide consistency across laboratories and yield more reliable estimates of the actual prevalence of enteric protozoa.

Key words: enteric protozoa, gastrointestinal illness, algorithms, molecular diagnosis, diagnostics, *Dientamoeba fragilis*, *Blastocystis* spp., *G. intestinalis*.

INTRODUCTION

Enteric protozoa are important causes of infectious diseases affecting people in developing as well as developed countries (WHO, 2008). Compared with developing countries, relatively few enteric protozoa are included in operational surveillance systems in developed countries. Where these are included, they are mainly seen as indicators of foodborne and waterborne diseases outbreaks (Cretikos *et al.* 2008b; WHO, 2008; Stark *et al.* 2009; Kucerova *et al.* 2010; Dixon *et al.* 2011; Sokolova *et al.* 2011). However, evidence suggests that while some enteric protozoa such as *Entamoeba* spp., *Cryptosporidium* spp. and *Giardia intestinalis* are more frequently identified in diarrhoeal cases in developing regions, like Asia and sub-Saharan

Africa (Nair *et al.* 2010; Fletcher *et al.* 2011); others like *Blastocystis* spp. and *Dientamoeba fragilis* appear to be more prevalent in the developed countries (Roberts *et al.* 2011; Fletcher *et al.* 2012). In developed settings, however, enteric protozoa are often ignored as a cause of diarrhoea due to the often mistaken belief that better hygiene practices are occurring.

In developed settings, bacterial cultures are usually considered initially in the diagnosis for acute diarrhoeal illnesses, while parasitic infections are more likely to be considered in patients with chronic symptoms, appropriate travel histories or other risk factors (Ribes *et al.* 2004). However, laboratory-based surveillance has been used as an important tool for estimating the burden of infectious diseases in several countries worldwide (Flint *et al.* 2005). In Australia, for example, Cryptosporidiosis and Giardiasis are the only parasitic gastrointestinal diseases included in the infectious disease surveillance (Cretikos *et al.* 2008a, Costello *et al.* 2009). Estimates of the actual prevalence of enteric

* Corresponding author. South Western Sydney Local Health District, P.O. Box 38, Liverpool 1871, New South Wales, Australia. Telephone: +61 2 8778 0855. Fax: +61 2 8778 0838. E-mail: stephanie.fletcher@sswahs.nsw.gov.au

protozoa in industrialized countries is often affected by: (i) the lack of routine testing for these parasites and (ii) the lack of sensitive diagnostic techniques to detect them in clinical specimens, while carrier stages and sub-clinical infections are often not diagnosed (Ng *et al.* 2011).

The actual burden of parasitic infections affecting Australians is relatively unknown. Anecdotal evidence suggests that the prevalence is relatively low; however, some individuals are at increased risk of infection. Parasitic infections are considered to be common among Aboriginal communities especially in children under 5 years of age (Commonwealth of Australia, 2000, Currie and Brewster, 2001). Reports indicate that men who have sex with men are at increased risk of infection (Stark, 2007; Stark *et al.* 2008). However, it is assumed that the estimated prevalence of protozoan infections is relatively similar regardless of the testing protocol employed. However, a scientific assessment of this has neither been done; nor has a gold standard approach been determined for diagnosis of enteric protozoan disease. This information is needed for early and accurate diagnosis to aid in the optimal management of parasitic diseases. This not only allows initiation of appropriate therapy, but also implementation of health and hygiene education and control measures in the patients' home and community.

Here we summarize a multi-centre study to determine the relative prevalence of enteric protozoan infections from clinical specimens examined at four public hospitals in Sydney, and the comparison of the outcome of different testing algorithms for the detection of enteric protozoa. Finally, this study suggests that molecular methods should be employed as a gold standard approach for clinical diagnosis of enteric protozoa.

METHODS

Setting and study sites

Four hospitals, Liverpool Hospital (A), The Children's Hospital at Westmead (B), St. Vincent's Hospital (C), Sydney, Prince of Wales Hospital (D), all located in different geographic areas across Sydney were included in the study. These facilities were included based on the population served, and represented a cross-section of different socio-economic and cultural influences across the Sydney metropolitan region. Hospital A, is a tertiary referral hospital for Southwestern Sydney; hospital B is a stand-alone service dedicated to paediatrics attracting referrals on a State-wide basis; hospital C is a major public and a principal referral hospital attracting referrals on a State-wide and national basis; and hospital D is a major teaching hospital and one of 13 principal referral hospitals for adults based in

Sydney's eastern suburb that also serves all of New South Wales. Each hospital provided a fully accredited laboratory service, providing comprehensive biomedical laboratory services.

Ethical approval for this study was received from the Human Research Ethics Committees of each hospital and the University of Technology, Sydney (UTS).

Microbiology methods

All four hospitals routinely tested for enteric organisms in persons who presented with gastrointestinal symptoms. On average, each laboratory tested one stool sample per patient, with between 45 and 89% of these specimens being loose – but not taking the shape of the container. Generally speaking, each laboratory used standard methods for the identification and isolation of enteric pathogens. Additionally, in all hospitals, stools were processed by a wet preparation in saline, and examined for white blood cells, red blood cells, cysts, ova and parasites (COP); bacteriological pathogens were identified using standard culturing methods and each hospital had specific criteria for the testing of viruses (Fletcher *et al.* 2015). A summary of the various tests done for parasitic agents is presented in Supplementary Table S1.

Parasitology

Hospital A. Stool specimens were routinely collected in sodium acetate acetic acid formalin (SAF) fixative (Oxoid Australia), and processed by a direct wet preparation. Light microscopy was routinely performed on all stool specimens. In the instances where no clinical information was received and the patient was an adult or age ≤ 10 years old, or the specimen was not received in SAF, then a *Giardia/Cryptosporidium* screen enzyme immunoassay (EIA) (ProSpecT™ *Giardia/Cryptosporidium* Microplate Assay) was performed. A 10% suspension of stool was prepared in 10% formalin (for *G. intestinalis* and *Cryptosporidium*) and the EIA was performed in accordance with the manufacturer's instructions and without modification. A full COP test was done on all positive microscopy and EIA results using an IHS with modified acid-fast stain.

Hospital B. Light microscopy of a direct saline preparation was performed on all stool specimens. Concentration techniques were performed routinely for persons with a history of overseas travel, prolonged diarrhoea illness (>7 days), attendees at refugee clinics and on specific requests for COP test by the clinician. When a COP test was requested and if any parasites were seen in the wet preparation, a sample of stool was placed into SAF fixative (Oxoid Australia) using a 1:5 ratio and processed

for fecal concentration and stained using the fecal parasite concentrator (Evergreen Scientific, LA, CA), which uses centrifugation at 500 g for 10 min and examined for COP using oil immersion. Alternatively, the fixed smear was prepared for permanent staining by the iron haematoxylin technique. Additionally, each stool specimen had a *Cryptosporidium* smear done routinely using a Modified Kinyoun's acid-fast stain (Cold).

Hospital C. Direct wet preparation and light microscopy were performed routinely on all stool specimens. The wet preparation was examined under a low-power objective (10×) and then scanned under the high dry (40×) objective. All stool specimens were emulsified in SAF fixative (Oxoid Australia) using a 1:3 ratio; and then the samples were centrifuged at 500 g for 10 min. Samples were then processed for permanent staining by a modified iron haematoxylin staining technique (mIHS) technique incorporating a carbol fuchsin step to stain for acid-fast organisms (*Isospora*, *Cryptosporidium* and *Cyclospora*). Stool samples also underwent direct DNA extraction using a QIAamp DNA stool minikit (Qiagen, Hilden, Germany) using a portion of fresh stools sample for the identification of *Entamoeba* spp. These methods have been previously described by Stark and colleagues (Stark *et al.* 2010c; Banik *et al.* 2011; Roberts *et al.* 2011).

Hospital D. Direct wet preparation microscopy was only conducted on patients at risk to parasitic infection as indicated in the clinical history and for patients with recent overseas travel or on request by the clinicians. The wet preparation was examined by light microscopy under low-power objective (10×) and then scanned under the high dry (40×) objective. A sample of stool was also placed into SAF fixative (Meridian Bioscience, Inc., Cincinnati, Ohio) followed by fecal concentration using the Mini Parasep[®]SF concentration kit (DiaSys Europe LTD, Laboratory Diagnostics PTY LTD). In addition, an EIA was performed routinely as a screening test for the detection of *G. intestinalis* and *Cryptosporidium* (ProSpecT[™] *Giardia/Cryptosporidium* Microplate Assay) and the detection of *Entamoeba histolytica/dispar* (ProSpecT[™] *Entamoeba histolytica*, Remel). A 10% suspension of stool was prepared in 10% formalin (for *G. intestinalis* and *Cryptosporidium*) and in specimen buffer provided in kit (for *E. histolytica*) and the EIA was performed in accordance with the manufacturer's instructions and without modification. All positive findings from the EIAs were confirmed by microscopy (i.e. iodine preparation and acid-fast stain). Samples testing positive on the *E. histolytica* EIA that could not be confirmed by direct microscopy (i.e. iodine preparation) were sent to a reference laboratory for permanent stain preparation and

examination. In order to detect *Cryptosporidium* oocysts, smears were made directly from feces and stained by the Ziehl–Neelsen based on the procedures described elsewhere (Collins and Lyne, 1984).

Data extraction and analysis

Each hospital provided a spread sheet containing de-identified microbiology test results for the period January 2007 to December 2010 (hospital C's data were for 2008–2010). The data were then arranged by medical record number, and date of service/stool request in ascending order. For each hospital, the testing protocols were consulted to determine the number of specimens tested for COP/intestinal parasites. The laboratory data were placed into a Statistical Package for the Social Science (SPSS) database and duplicate results removed to avoid duplicate counting of specimens. Duplicates were considered to be any stool specimen from the same individual that was collected on the same date and had the same request number. For the purposes of this analysis, each individual stool sample and results were counted. Positivity was calculated on the basis of one organism per specimen. The percentage positivity rate was calculated as the total number of stool samples positive for an enteric organism divided by the total number of specimens tested. Odds ratios were calculated for each specific test conducted to measure the association between the detection of pathogens (outcome) at different hospitals (exposure). A laboratory survey identified laboratory procedures and information captured on laboratory request forms.

RESULTS

The laboratory survey identified laboratory procedures and information captured on laboratory request forms in the four hospitals. The date of sample collection, age and gender were reported routinely on all requests in all hospitals. Differential diagnosis was reported only sometimes in two hospitals and rarely in the other two. Signs and symptoms were only reported sometimes at all sites, and date of onset of illness was rarely reported except by hospital D, where it was routinely done. Across all hospitals, between 1 and 10% of stool specimens received were formed, while 45–89% were unformed (loose but not taking the shape of the container). Only 10–50% of specimens were considered liquid (taking the shape of the container).

Tests for enteric parasites were conducted on 2138 individual specimens from 1518 persons at hospital A; 11097 specimens from 5229 persons at hospital B; 8613 specimens tested from 6273 persons at hospital C; and 6078 specimens tested from 3772 persons at hospital D.

Table 1. Overall prevalence of enteric protozoa from Cyst, Ova and Parasite test, Hospital A, 2007–2010

Organism identified	Total single specimen tested (n)	%	Total multiple specimen tested (n)	%	Total specimen	% of Overall specimen tested
<i>Blastocystis</i> spp.	92	6.1	30	4.8	122	5.7
<i>Giardia intestinalis</i>	21	1.4	4	0.6	25	1.2
<i>Cryptosporidium</i> spp.	7	0.5	3	0.5	10	0.5
<i>Dientamoeba fragilis</i>	7	0.5	0	0.0	7	0.3
<i>Entamoeba hartmanni</i>	5	0.2	1	0.2	6	0.3
<i>Entamoeba histolytica/dispar</i>	3	0.1	1	0.2	4	0.2
<i>Chilomastix mesnili</i>	1	0.1	2	0.3	3	0.1
<i>Endolimax nana</i>	1	0.1	8	1.3	9	0.4
<i>Enteromonas hominis</i>	0	0.0	1	0.2	1	0.1
Subtotal Protozoa positive	137	9.2	50	8.1	187	8.8
Other pathogens	15	1.1	5	0.8	20	1.0
Samples Positive	152	10.0	55	8.9	207	9.7
Samples Negative	1366	90.0	565	91.1	1931	90.3
Total samples tested	1518	100.0	620	29.0	2138	100.0

Table 2. Overall prevalence of enteric protozoa from cyst, ova and parasite test, hospital B 2007–2010

Organism identified	Total single specimen tested (n)	%	Total multiple specimen tested (n)	%	Total specimen	% of Overall specimen tested
<i>Giardia intestinalis</i>	38	0.7	10	0.2	48	0.47
<i>Blastocystis</i> spp.	29	0.6	11	0.2	40	0.4
<i>Dientamoeba fragilis</i>	19	0.4	2	0.0	21	0.21
<i>Cryptosporidium</i> spp.	8	0.2	1	0.0	9	0.09
Subtotal Protozoa	94	1.9	24	0.5	118	1.2
Other pathogens	838	15.9	101	2.1	939	9.3
Samples positive	932	17.8	115	2.3	1057	10.5
Samples negative	4297	82.2	4769	97.4	9066	89.5
Total samples tested	5229	100.0	4894	100.0	10123	100.0

Enteric parasites summary

Enteric protozoa were identified in an average of 3.6% (95% CI 1.1–11.2%) of specimens from the four hospitals. Across the four hospitals, the most common enteric protozoon detected was *Blastocystis* spp., identified in an average of 5.4% (95% CI 5.0–5.7%) of cases, followed by *G. intestinalis* 1.1% (95% CI 1.0–1.2%), *D. fragilis* in 0.8% (95% CI 0.7–1.0%), *E. histolytica/dispar* in 0.5% (95% CI 0.4–0.6%), *Cryptosporidium* spp. 0.3% (95% CI 0.3–0.4%), *Cyclospora* 0.1% (95% CI 0.02–0.1%). Non-pathogenic protozoa, including *Entamoeba* spp., *Enteromonas hominis* and *Iodamoeba butschlii* were found in <1% of cases, respectively.

At hospital A, 29% of patients submitted multiple specimens, with an enteric protozoon found in 8%. At hospital B, 48% of patients submitted multiple specimens, with enteric protozoa detected in <1%. At hospital C, 38% of patients submitted multiple specimens and protozoa were detected in 8.5%. At hospital D, 38% of patients submitted multiple specimens and 0.5% tested positive.

The results for hospital A are summarized in Table 1. A total 9% (187/2138) of stool specimens examined had an enteric protozoon identified.

Overall, *Blastocystis* spp. (5.71% or 122/2138) and *G. intestinalis* (1.17% or 25/2138) were most frequently detected. At hospital B, an enteric protozoon was detected in 1% (70/10123) of stools (Table 2); *G. intestinalis* (0.5% or 48/10123) and *Blastocystis* spp. (0.4% or 40/10123) were most frequently identified. One or more protozoa were found in 12% (1003/8613) specimens at hospital C (Table 3). *Blastocystis* spp., 7% (571/8613), *Giardia* in 2% (141/8613) and *D. fragilis* in 1% (100/8613) were most frequently detected. At hospital D, 1% (78/6078) of stool specimens tested positive for one or more enteric protozoa (Table 4). A total of 1% (77/3772) tested positive for the *Giardia/Cryptosporidium* coproantigen test (by EIA). However, only 56% (43) of these were confirmed by microscopy of wet preparation to be *G. intestinalis* and 20 (26.0%) confirmed to be *Cryptosporidium* spp.

Comparison of results based on testing protocols

Approximately 2.5% (95% CI 2.3–2.7%) of protozoon infections was detected by permanent staining (IHS or mIHS), 1.1% (95% CI 1.0–1.2%) by microscopy of wet preparations and 0.6% (95% CI 0.5–0.7%) by EIA combined with microscopy.

Table 3. Overall prevalence of enteric protozoa from cyst, ova and parasite test, hospital C for 2008–2010

Organism identified	Total single specimen tested (n)	%	Total multiple specimen tested (n)	%	Total specimen	% of Overall specimen tested
<i>Blastocystis</i> spp.	429	5.0	142	4.3	571	6.6
<i>Giardia intestinalis</i>	109	5.7	32	1.0	141	1.6
<i>Dientamoeba fragilis</i>	71	3.7	29	0.9	100	1.2
<i>Endolimax nana</i>	34	0.4	30	0.9	64	0.7
<i>Cryptosporidium</i> spp.	33	0.4	4	0.1	37	0.4
<i>Entamoeba coli/hartmanni</i>	21	0.2	15	0.5	36	0.4
<i>Entamoeba histolytica/dispar</i>	47	0.8	4	0.1	51	0.6
<i>Enteromonas hominis</i>	7	0.1	9	0.3	16	0.2
<i>Cyclospora</i>	5	0.1	0	0.0	5	0.1
<i>Iodameba</i>	2	0.02	17	0.5	19	0.2
<i>Chilomastix</i>	1	0.01	0	0.0	1	0.01
Subtotal Protozoa	758	18.4	282	12.1	1041	12.1
Other pathogens	1196	19.1	402	17.2	1598	18.5
Samples positive	1954	31.2	684	29.3	2639	30.6
Samples negative	4319	68.8	1656	70.7	5974	69.4
Total samples tested	6273		2340	38.4	8613	

Table 4. Overall prevalence of enteric protozoa from cyst, ova and parasite test, hospital D for 2007–2010

Organism identified	Total single specimen tested (n)	%	Total multiple specimen tested (n)	%	Total specimen	% of Overall specimen tested
<i>Giardia intestinalis</i>	38	1.0	5	0.2	43	0.7
<i>Cryptosporidium</i> spp.	16	0.4	4	0.2	20	0.3
<i>Blastocystis</i> spp.	5	0.0	1	0.04	6	0.1
<i>Entamoeba histolytica/dispar</i>	3	0.0	2	0.1	5	0.1
<i>Entamoeba coli/hartmanni</i>	2	0.1	1	0.04	3	0.1
<i>Endolimax nana</i>	1	0.0	0	0.0	1	0.02
Sub-total Protozoa	65	1.7	13	0.6	78	1.3
Other organisms	2	0.1	0	0.0	2	0.03
Samples positive	67	1.8	13	0.6	80	1.33
Samples negative	3705	98.2	2293	99.4	5998	98.7
Total	3772		2306	37.9	6078	100.0

The mean difference between tests conducted at each hospital is presented in Table 5. *Blastocystis* spp. was more frequently detected at hospital A when compared with hospitals B and D (mean difference 5.3 and 5.6%, respectively; $P = 0.0002$), and at hospital C when compared with hospitals B and D (mean difference >6% each; $P < 0.0001$). A higher rate of detection was also observed for *Giardia* at hospital A when compared with hospitals B and D (mean diff. 0.7%; $P < 0.0002$ and 0.5%, $P < 0.05$, respectively) and hospital C when compared with hospitals B and D (mean different 1.2 and 0.93%; respectively; $P < 0.0001$). In addition, hospital C diagnosed significantly more *E. histolytica/dispar* when compared with hospital D (mean diff. 0.5%; $P < 0.0002$).

The combination of microscopy of wet preparation and EIA detected the prevalence of *Cryptosporidium* spp., in an average of 0.4% [95% CI 0.3–0.5%; OR 1.4 (0.7–3.0)] and *G. intestinalis* in 0.9% [95% CI 0.7–1.1%; OR 1.7 (1.0–2.7)] with no significant differences in detection between

hospitals A and D ($P > 0.05$). Microscopy of wet preparation detected *Blastocystis* spp., in an average of 2.9% (95% CI 2.5–3.4%) of cases, with significantly higher detection rates at hospital A compared with hospital B (OR 14.4; 95% CI 10.1–20.7; $P < 0.0001$). Permanent staining with IHS or mIHS detected *D. fragilis* in an average of 1.1% (95% CI 0.9–1.4%) of cases between hospitals A and C; with hospital C detecting significantly higher rates by employing a mIHS (OR 3.6; 95% CI 1.7–7.7; $P < 0.001$).

DISCUSSION

We present the prevalence of enteric protozoa amongst persons seeking care for gastrointestinal illnesses in Sydney across four major public hospitals. The study reveals that while all four laboratories performed direct microscopy on stool specimens for the detection of cyst, ova and parasites, different approaches are used for different species and tests for some protozoa are not routinely done.

Table 5. Mean difference in protozoa detection rates amongst four Sydney hospitals

Organisms detected	Hospital A/B Mean difference % (s-value) P-value	Hospital A/C Mean difference % (s-value) P-value	Hospital A/D Mean difference % (s-value) P-value	Hospital B/C Mean difference % (s-value) P-value	Hospital B/D Mean difference % (s-value) P-value	Hospital C/D Mean difference % (s-value) P-value
<i>Blastocystis</i> spp.	5.3% (1.9-5) <0.0002	0.9% (1.6) 0.060	5.6% (18.0) 0.0002	6.2% (23.9) <0.0001	0.3% (3.433) 0.0006	6.5% (20.1) <0.0001
<i>Chilomastix mesnili</i>	0.1 (3)	NT	NT	NT	NT	NT
<i>Cryptosporidium</i> spp.	0.4% (4.1) <0.0002	0.04% (0.2) 0.406	0.1% (0.9) 0.361	0.34% (4.7) <0.0001	0.2% (3.501) <0.0005	0.1% (1.0) <0.167
<i>Cyclospora</i>	NT	NT	0.03 (2)	0.0 (0)	NT	NT
<i>Dientamoeba fragilis</i>	0.1% (1.1) 0.291	0.83% (3.5) 0.0003	0.0 (0)	0.95% (8.1) <0.0001	0.0 (0)	0.0 (0)
<i>Endolimax nana</i>	0.0 (0)	0.03% (1.6) 0.052	0.4% (N/C)	0.0 (0)	0.0 (0)	0.7% (N/C)
<i>Entamoeba hartmanni/coli</i>	0.0 (0)	0.14% (0.9) 0.181	0.2% (N/C)	0.0 (0)	0.0 (0)	0.37% (N/C)
<i>Entamoeba histolytica/dispar</i> complex	0.0 (0)	0.4% (N/C)	0.1% (N/C)	0.0 (0)	0.0 (0)	0.6% (4.9) <0.0002
<i>Enteromonas hominis</i>	0.0 (0)	0.1% (N/C)	0.0 (0)	0.0 (0)	0.0 (0)	0.0 (0)
<i>Giardia intestinalis</i>	0.7% (3.8) <0.0002	0.47% (1.6) 0.058	0.5% (2.0) 0.043	1.2% (7.9) <0.0001	0.2% (1.924) 0.054	0.9% (5.0) <0.0001

N/C, not calculated. Cells contained fewer than 5, hence estimates could not be calculated. NT, test not done. Total specimen tested hospital A, 2138; hospital B, 10123; hospital C, 8163; and hospital D, 6078.

We found that, while the prevalence of enteric protozoa species is relatively low in this population, widespread variability in the testing protocols as well as individual and population characteristics may influence protozoan detection rates in this population. Progress towards development of a gold standard approach for diagnosis of disease is warranted.

The rate of detection of enteric parasites between the four hospitals varied. Generally, *Blastocystis* spp. and *G. intestinalis* were the most common enteric protozoa identified in patients. An age relationship was observed with *Giardia* prevalence, with higher rates detected in the 0–5 age group, compared with higher rates of *Blastocystis* spp. in the over 5 age group. This age relationship has been previously described in this population (Fletcher *et al.* 2014) and may indicate that children are more frequently exposed to giardiasis risk factors in this setting, as described previously (Fletcher *et al.* 2012; Yoder *et al.* 2012).

The proportion of stool specimens positive for an enteric parasite between hospitals ranged from a low of 1% in hospitals B and D, to a high of 11.6% at hospital C. The difference in detection rates between Hospitals may be due to various factors. Individual hospitals had different testing criteria for enteric protozoa, hence not all stool specimen were tested for enteric protozoa. Both hospitals A and C tested for a wider range of pathogens routinely (see Tables 1 and 3), including non-pathogenic species, which may be driven by the high-risk populations served including recent migrants (A), men who have sex with men, and HIV/AIDs infected persons (C). Secondly, the composition of the population seen at each hospital could have influenced test results if risk factors were unequally distributed in the population (Mohr and Mohr, 1992). Previously published data on a randomly selected subset of this population indicated that there were minor differences in the sex distribution between hospitals except for hospital C, where there was slightly more males. However, there were significant differences between the age distribution, particularly in the under 5 years age group between hospitals; based on the population served by hospitals. Enteric protozoa were more prevalent in children under 5 years of age in this population (Fletcher *et al.* 2015). Age associated risk factors influencing the underlying prevalence of the condition in this population may therefore account for some variations in detection rates between hospitals. On the other hand, some hospitals did not routinely test for COP if the prevalence of protozoa was relatively low in the population, and was likely to generate many false negatives. The testing protocols may therefore be secondary to a perceived prevalence within high-risk groups in the wider population (e.g. men who have sex with men, recent immigrants and lower socio-economic groups). Both hospitals A and C had specific clinics that catered to high-risk groups.

The differences in detection rates could also be associated with the different diagnostic techniques and handling practices between hospitals. According to Libman *et al.* (2008), significant variations in specimen handling and processing practices between laboratories can affect the assessment of the diagnostic processes. This study found that where wet preparation microscopy of fresh or fixed stool (SAF) specimens, detected a lower prevalence of protozoa. One disadvantage of using the microscopy method only is its low sensitivity to detect protozoa, which lead to false-negative results (Stark *et al.* 2010a; Roberts *et al.* 2011). This is particularly true for protozoan such as *D. fragilis* that requires special staining techniques to detect its nuclear structure (Stark *et al.* 2010b; Stark *et al.* 2011).

The mIHS method consistently detected significantly higher rates of *Blastocystis* spp. and *G. intestinalis* at hospital C and significantly higher rates of *D. fragilis* when compared with the IHS-only method at hospital A and wet preparation microscopy at hospital B. Microscopy remains a widely used tool for protozoan detection, even in low prevalence settings such as Sydney (Bruijnesteijn Van Coppenraet *et al.* 2009; Ghoshal *et al.* 2016). While molecular methods are more sensitive for pathogen detection, these tools are still not widely available or routinely employed even in developed settings (Fletcher *et al.* 2012; Ghoshal *et al.* 2016).

This study has various limitations. The data were collected retrospectively since approval was not obtained for prospective data collection. The authors are mindful that the incidence figures for each hospital should be compared with caution based on the differences between the testing protocols, and their ability to detect protozoa. The authors cannot exclude that some of the differences found between the four hospitals can be explained by actual differences in parasite prevalence in the underlying populations tested. One potential bias of this study is that the hospitals that followed specific criteria for stool testing for protozoa, identified higher rates of protozoa, regardless of the test used as evidenced by *Blastocystis* and *Giardia* rates at hospital A. This is further impacted by the relatively low prevalence of some organisms in the population, small proportion of persons who seek medical attention and even fewer who get tested.

Notwithstanding, these results are useful to local and state health authorities to guide disease surveillance activities for these organisms, aid in the understanding of the epidemiology of protozoan infections in Sydney, and provide the basis for setting research priorities and planning interventions. The development of a gold standard approach for diagnosis of enteric protozoa, which addresses issues such as relatively low incidence of some species such as *Cryptosporidium* and *Cyclospora*; the difficulties in diagnosis (e.g. *D. fragilis*) and differentiation (e.g.

Entamoeba spp.) of some species, is therefore warranted.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/pao.2016.7>.

ACKNOWLEDGEMENTS

We acknowledge the medical records and laboratory staff that extracted the data for each hospital.

FINANCIAL SUPPORT

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

CONFLICT OF INTEREST

None.

ETHICAL STANDARDS

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

REFERENCES

- Banik, G. R., Barratt, J. L. N., Marriott, D., Harkness, J., Ellis, J. T. and Stark, D.** (2011). A case-controlled study of *Dientamoeba fragilis* infections in children. *Parasitology* **138**, 819–823.
- Bruijnesteijn Van Coppenraet, L. E. S., Wallinga, J. A., Ruijs, G. J. H. M., Bruins, M. J. and Verweij, J. J.** (2009). Parasitological diagnosis combining an internally controlled real-time PCR assay for the detection of four protozoa in stool samples with a testing algorithm for microscopy. *Clinical Microbiology and Infection* **15**, 869–874.
- Collins, C. and Lyne, P.** (1984). *Microbiological Methods*, London, Butterworths.
- Commonwealth of Australia** (2000). *Nutrition in Aboriginal and Torres Strait Islander Peoples*, National Health and Medical Research Council (NHMRC), Canberra, ACT, Australia.
- Costello, A., Abbas, M., Allen, A., Ball, S., Bell, S., Bellamy, R., Friel, S., Groce, N., Johnson, A., Kett, M., Lee, M., Levy, C., Maslin, M., McCoy, D., Mcguire, B., Montgomery, H., Napier, D., Pagel, C., Patel, J., De Oliveira, J. A. P., Redclift, N., Rees, H., Rogger, D., Scott, J., Stephenson, J., Twigg, J., Wolff, J. and Patterson, C.** (2009). Managing the health effects of climate change. *Lancet* **373**, 1693–1733.
- Cretikos, M., Telfer, B. and Mcanulty, J.** (2008a). Enteric disease outbreak reporting, New South Wales, Australia, 2000 to 2005. *New South Wales Public Health Bulletin* **19**, 3–7.
- Cretikos, M., Telfer, B. and Mcanulty, J.** (2008b). Evaluation of the system of surveillance for enteric disease outbreaks, New South Wales, Australia, 2000 to 2005. *New South Wales Public Health Bulletin* **19**, 8–14.
- Currie, B. J. and Brewster, D. R.** (2001). Childhood infections in the tropical north of Australia. *Journal of Paediatrics and Child Health* **37**, 326–330.
- Dixon, B., Parrington, L., Cook, A., Pintar, K., Pollari, F., Kelton, D. and Farber, J.** (2011). The potential for zoonotic transmission of *Giardia duodenalis* and *Cryptosporidium* spp. from beef and dairy cattle in Ontario, Canada. *Veterinary Parasitology* **175**, 20–26.
- Fletcher, S. M., Stark, D. and Ellis, J.** (2011). Prevalence of gastrointestinal pathogens in sub-Saharan Africa; systematic review and meta-analysis. *Journal of Public Health in Africa* **2**, 127–137.

- Fletcher, S. M., Stark, D., Harkness, J. and Ellis, J. (2012). Enteric protozoa in the developed world: a public health perspective. *Clinical Microbiology Reviews* **25**, 420–449.
- Fletcher, S., Caprarelli, G., Merif, J., Andresen, D., Hal, S. V., Stark, D. and Ellis, J. (2014). Epidemiology and geographical distribution of enteric protozoan infections in Sydney, Australia. *Journal of Public Health Research* **3**, 298.
- Fletcher, S., Sibbritt, D., Stark, D., Harkness, J., Rawlinson, W., Andresen, D., Van Hal, S., Merif, J. and Ellis, J. (2015). Descriptive epidemiology of infectious gastrointestinal illnesses in Sydney, Australia, 2007–2010. *Western Pacific Surveillance and Response* **6**. doi: <http://dx.doi.org/10.5365%2Fwpsar.v6i4.345>
- Flint, J. A., Van Duynhoven, Y. T., Angulo, F. J., Delong, S. M., Braun, P., Kirk, M., Scallan, E., Fitzgerald, M., Adak, G. K., Sockett, P., Ellis, A., Hall, G., Gargouri, N., Walke, H. and Braam, P. (2005). Estimating the burden of acute gastroenteritis, food-borne disease, and pathogens commonly transmitted by food: an international review. *Clinical Infectious Diseases* **41**, 698–704.
- Ghoshal, U., Dey, A., Ranjan, P., Khanduja, S., Agarwal, V. and Ghoshal, U. C. (2016). Identification of opportunistic enteric parasites among immunocompetent patients with diarrhoea from Northern India and genetic characterisation of *Cryptosporidium* and Microsporidia. *Indian Journal of Medical Microbiology* **34**, 60–66.
- Kucerova, Z., Sokolova, O. I., Demyanov, A. V., Kvac, M., Sak, B., Kvetonova, D. and Secor, W. E. (2010). Microsporidiosis and cryptosporidiosis in HIV/AIDS patients in St. Petersburg, Russia: serological identification of microsporidia and *Cryptosporidium parvum* in sera samples from HIV/AIDS patients. *AIDS Research and Human Retroviruses* **27**, 13–15.
- Libman, M. D., Gyorkos, T. W., Kokoskin, E. and Maclean, J. D. (2008). Detection of pathogenic protozoa in the diagnostic laboratory: result reproducibility, specimen pooling, and competency assessment. *Journal of Clinical Microbiology* **46**, 2200–2205.
- Mohr, E. and Mohr, I. (1992). Statistical analysis of the incidence of positives in the examination of parasitological specimens. *Journal of Clinical Microbiology* **30**, 1572–1574.
- Nair, G., Ramamurthy, T., Bhattacharya, M., Krishnan, T., Ganguly, S., Saha, D., Rajendran, K., Manna, B., Ghosh, M., Okamoto, K. and Takeda, Y. (2010). Emerging trends in the etiology of enteric pathogens as evidenced from an active surveillance of hospitalized diarrhoeal patients in Kolkata, India. *Gut Pathogens* **2**, 4.
- Ng, J., Yang, R., Whiffin, V., Cox, P. and Ryan, U. (2011). Identification of zoonotic *Cryptosporidium* and *Giardia* genotypes infecting animals in Sydney's water catchments. *Experimental Parasitology* **128**, 138–144.
- Ribes, J. A., Seabolt, J. P. and Overman, S. B. (2004). Point prevalence of cryptosporidium, cyclospora, and isospora infections in patients being evaluated for diarrhea. *American Journal of Clinical Pathology* **122**, 28–32.
- Roberts, T., Barratt, J., Harkness, J., Ellis, J. and Stark, D. (2011). Comparison of microscopy, culture, and conventional polymerase chain reaction for detection of *Blastocystis* sp. in clinical stool samples. *American Journal of Tropical Medicine and Hygiene* **84**, 308–312.
- Sokolova, O. I., Demyanov, A. V., Bowers, L. C., Didier, E. S., Yakovlev, A. V., Skarlato, S. O. and Sokolova, Y. Y. (2011). Emerging microsporidia infections in Russian HIV-infected patients. *Journal of Clinical Microbiology* **49**, 2102–2108.
- Stark, D. E. A. (2007). Prevalence of enteric protozoa in human immunodeficiency Virus (HIV)-positive and HIV negative men who have sex with men from Sydney, Australia. *American Journal of Tropical Medicine and Hygiene* **76**, 549–552.
- Stark, D., Van Hal, S., Matthews, G., Marriott, D. and Harkness, J. (2008). Invasive Amebiasis in men who have sex with men, Australia. *Emerging Infectious Diseases* **14**, 1141–1143.
- Stark, D., Barratt, J. L. N., Van Hal, S., Marriott, D., Harkness, J. and Ellis, J. T. (2009). Clinical significance of enteric protozoa in the immunosuppressed human population. *Clinical Microbiology Reviews* **22**, 634–650.
- Stark, D., Barratt, J., Roberts, T., Marriott, D., Harkness, J. and Ellis, J. (2010a). Comparison of microscopy, two xenic culture techniques, conventional and real-time PCR for the detection of *Dientamoeba fragilis* in clinical stool samples. *European Journal of Clinical Microbiology & Infectious Diseases* **29**, 411–416.
- Stark, D., Barratt, J., Roberts, T., Marriott, D., Harkness, J. and Ellis, J. (2010b). A review of the clinical presentation of dientamoebiasis. *American Journal of Tropical Medicine and Hygiene* **82**, 614–619.
- Stark, D., Schuller, M., Sloots, T. P., James, G. S., Halliday, C. L. and Carter, I. W. J. (2010c). *Entamoeba histolytica* PCR for Clinical Microbiology. Springer, Netherlands.
- Stark, D., Al-Qassab, S. E., Barratt, J. L. N., Stanley, K., Roberts, T., Marriott, D., Harkness, J. and Ellis, J. T. (2011). Evaluation of multiplex tandem real-time PCR for detection of *Cryptosporidium* spp, *Dientamoeba fragilis*, *Entamoeba histolytica*, and *Giardia intestinalis* in clinical stool samples. *Journal of Clinical Microbiology* **49**, 257–262.
- Who (2008). *The Global Burden of Disease: 2004 Update Geneva*, World Health Organization (WHO) Press, Switzerland, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
- Yoder, J. S., Gargano, J. W., Wallace, R. M. and Beach, M. J. (2012). Giardiasis surveillance – United States, 2009–2010. *Morbidity and Mortality Weekly Report* **61**, 13–23.