

Differential Harnack estimates for a weighted nonlinear parabolic equation under a super Perelman–Ricci flow and implications

Ali Taheri  and Vahideh Vahidifar

School of Mathematical and Physical Sciences, University of Sussex,
 Falmer, Brighton, UK (a.taheri@sussex.ac.uk; v.vahidifar@sussex.ac.uk)

(Received 17 January 2023; accepted 24 August 2023)

In this paper, we derive new differential Harnack estimates of Li–Yau type for positive smooth solutions to a class of nonlinear parabolic equations in the form

$$\mathcal{L}_\phi^a[w] := \left[\frac{\partial}{\partial t} - \mathbf{a}(x, t) - \Delta_\phi \right] w(x, t) = \mathcal{G}(t, x, w(x, t)), \quad t > 0,$$

on smooth metric measure spaces where the metric and potential are time dependent and evolve under a (k, m) -super Perelman–Ricci flow. A number of consequences, most notably, a parabolic Harnack inequality, a class of Hamilton type global curvature-free estimates and a general Liouville type theorem together with some consequences are established. Some special cases are presented to illustrate the strength of the results.

Keywords: Smooth metric measure spaces; ϕ -Laplacian; Perelman–Ricci flow; gradient estimates; Bakry–Émery tensor; Harnack inequality; Liouville type results

2020 *Mathematics Subject Classification:* 53E20; 53C23; 53C21; 58J60; 58J35; 60J60

1. Introduction

In this paper, we study gradient estimates for positive smooth solutions to a class of nonlinear parabolic equations on smooth metric measure spaces with evolving geometry. Whilst gradient estimates have been studied over the years for linear and nonlinear equations on static manifolds or for equations of mainly Schrödinger and heat types on evolving manifolds, the investigation of nonlinear parabolic equations on smooth metric measure spaces with evolving metrics and potentials is relatively new and recent. These problems pose interesting challenges and technicalities whilst having huge scope for applications as already known and further discussed below.

To this end let (M, g) be a complete (smooth) Riemannian manifold of dimension $n \geq 2$ with Riemannian volume measure dv_g and let $d\sigma = \omega dv_g$ be a positive weighted measure on M with weight function $\omega > 0$ and potential $\phi = -\log \omega$. The triple $(M, g, d\sigma)$ is called a smooth metric measure space or a weighted manifold or a manifold with density (see [8, 21] for background and § 3).

Our aim here is to prove differential Harnack estimates for positive smooth solutions $w = w(x, t)$ to the following nonlinear parabolic equation on $(M, g, d\sigma)$ where the metric tensor g and potential ϕ evolve under a (k, m) -super Perelman–Ricci flow:

$$\begin{cases} \mathcal{L}_\phi^a[w](x, t) = \left[\frac{\partial}{\partial t} - \mathbf{a}(x, t) - \Delta_\phi \right] w(x, t) = \mathcal{G}(t, x, w(x, t)), & t > 0, \\ \frac{1}{2} \frac{\partial g}{\partial t}(x, t) + \text{Ric}_\phi^m(g)(x, t) \geq -kg(x, t), & m \geq n, k \geq 0, \\ \text{Ric}_\phi^m(g)(x, t) = \text{Ric}(g)(x, t) + \nabla_{g(t)} \nabla_{g(t)} \phi(x, t) - \frac{\nabla_{g(t)} \phi \otimes \nabla_{g(t)} \phi}{m - n}. \end{cases} \quad (1.1)$$

To describe the above system in more detail we first note that the differential operator Δ_ϕ in (1.1) is the ϕ -Laplacian associated with the triple $(M, g, d\sigma)$ (also known as the weighted or drifting or Witten Laplacian). It is a natural extension of the Riemannian Laplacian to the smooth metric measure space setting whose action on $v \in \mathcal{C}^2(M)$ can be described by

$$\Delta_\phi v = \Delta v - \langle \nabla \phi, \nabla v \rangle = e^\phi \text{div}(e^{-\phi} \nabla v). \quad (1.2)$$

Here, Δ , div and ∇ are the usual Laplace–Beltrami, divergence and gradient operators associated with the metric tensor g respectively. Naturally, when the metric tensor g and potential ϕ are time dependent this (spatial) differential operator is time dependent too (in that its coefficients depend explicitly on the time variable). Continuing further and referring again to the first equation in (1.1), the time evolution operators

$$\mathcal{L}_\phi^a = \frac{\partial}{\partial t} - \mathbf{a}(x, t) - \Delta_\phi, \quad \mathcal{L}_\phi = \mathcal{L}_\phi^0 = \frac{\partial}{\partial t} - \Delta_\phi, \quad \mathcal{L}_\phi^a = \mathcal{L}_\phi - \mathbf{a}, \quad (1.3)$$

are the \mathbf{a} -weighted (and weighted) heat operators with $\mathbf{a} = \mathbf{a}(x, t)$ a sufficiently smooth function of the space–time variables (x, t) . The nonlinearity $\mathcal{G} = \mathcal{G}(t, x, w)$ on the right-hand side here is a sufficiently smooth function depending on both the space–time variables and the dependent variable w . We shall present later on, various examples of such nonlinearities from different contexts ranging from conformal geometry, relativity and mathematical physics to applications of mathematics in materials and biological sciences, each representing a different phenomenon whilst depicting a corresponding singular or regular behaviour on its domain.

The differential inequality and equation on the second and third lines in system (1.1) respectively describe the evolution of the geometry of the triple $(M, g, d\sigma)$. Indeed the inequality in the second line in (1.1) describes the evolution of the generalized Ricci curvature tensor that in turn should be interpreted in the sense of symmetric $(2, 0)$ tensors whilst the equation on the third line gives the formulation of this generalized Ricci tensor that in turn involves the usual Ricci curvature tensor associated with g , the Hessian of ϕ and a rank-one symmetric tensor involving the gradient of ϕ (see § 3 for more on notation and background). Here $m \geq n$ is a constant (not necessarily an integer) having the role of a dimension and $k \geq 0$ is a fixed constant.

It is evident that the static case, that is, the case with time-independent metrics and potentials, constitutes a special case of system (1.1). In this event, the differential inequality on the second line in the system reduces to a spatial lower bound on the (time independent) generalized Ricci tensor (see § 3 for more), that is,

$$\mathcal{R}ic_\phi^m(g) = \mathcal{R}ic(g) + \nabla\nabla\phi - \frac{\nabla\phi \otimes \nabla\phi}{m-n} \geq -kg. \tag{1.4}$$

Here gradient estimates for positive solutions to linear and nonlinear heat type equations have been studied extensively starting from the seminal paper of Li and Yau [29] (see also [28]). In the nonlinear setting perhaps the first equation to be considered is the one with a logarithmic type nonlinearity (see e.g., [27, 33, 52])

$$\mathcal{L}_\phi^3[w] = \left[\frac{\partial}{\partial t} - a(x, t) - \Delta_\phi \right] w = A(x, t)w \log w. \tag{1.5}$$

The interest in such problems originates partly from its natural links with gradient Ricci solitons and partly from links with geometric and functional inequalities on manifolds, notably, the logarithmic Sobolev and energy-entropy inequalities [7, 8, 22, 49]. Recall that a Riemannian manifold (M, g) is said to be a gradient Ricci soliton if there exists a smooth function ϕ on M and a constant $\lambda \in \mathbb{R}$ such that (cf. [13, 16, 32])

$$\mathcal{R}ic_\phi(g) = \mathcal{R}ic(g) + \nabla\nabla\phi = \lambda g. \tag{1.6}$$

The notion is a generalization of an Einstein manifold and has a fundamental role in the analysis of singularities of the Ricci flow [24, 55].

Another prominent class of nonlinear equations rooted in conformal geometry and studied extensively are the Yamabe type equations (see [9, 20, 26, 34]). In the context of smooth metric measure spaces these equations can be broadly studied as (see e.g., [17, 40, 41, 54])

$$\mathcal{L}_\phi[w] = \frac{\partial w}{\partial t} - \Delta_\phi w = A(x, t)w^p + B(x, t)w. \tag{1.7}$$

Incidentally, the case $A \equiv -1, B \equiv 1, p = 3$ [$\mathcal{G}(w) = w - w^3$] is the Allen–Cahn equation and the case $A \equiv -c, B \equiv c, p = 2$ [$\mathcal{G}(w) = cw(1 - w)$ with $c > 0$] is the Fisher-KKP equation (cf. [2, 18, 25]). Both these equations have been studied extensively in recent years due to the significance of the phenomenon they model and their huge applications in physics and other sciences (for various geometric estimates and their consequences, see [5, 14, 44] and the references therein). A far reaching generalization of (1.7) with a superposition of power-like nonlinearities consist of equations in the form

$$\mathcal{L}_\phi[w] = \frac{\partial w}{\partial t} - \Delta_\phi w = \sum_{j=1}^d A_j(x, t)w^{p_j} + \sum_{j=1}^d B_j(x, t)w^{q_j}. \tag{1.8}$$

Here A_j, B_j (with $1 \leq j \leq d$) are sufficiently smooth space–time dependent coefficients and $p_j \geq 0, q_j \leq 0$ real exponents (see [42, 43]). Other classes of equations

generalizing the above and close to (1.5) come in the form (see e.g., [1, 10, 19, 42, 43, 48, 53])

$$\mathcal{L}_\phi^a[w] = \left[\frac{\partial}{\partial t} - a(x, t) - \Delta_\phi \right] w = A(x, t)w^p \Phi(\log w) + B(x, t)w^q, \tag{1.9}$$

with p, q real exponents, A, B sufficiently smooth coefficients and $\Phi \in \mathcal{C}^1(\mathbb{R}, \mathbb{R})$. Some cases of interest for $\Phi = \Phi(s)$ include power function, e.g., s^α (integer $\alpha \geq 1$), $|s|^\alpha$ or $|s|^{\alpha-1}s$ (real $\alpha > 1$) with different sign-changing, growth and singular behaviour as $\log w \rightarrow \pm\infty$ or a superposition of such nonlinearities [43]. Furthermore, the case of iterated logarithms as introduced in [10] with $d, k_1, \dots, k_d \in \mathbb{N}$ and $\beta_1, \dots, \beta_d \in \mathbb{R}$,

$$\Phi_{k_1, \dots, k_d}^{\beta_1, \dots, \beta_d}(\log w) = |\log_{k_1} w|^{\beta_1} |\log_{k_2} w|^{\beta_2} \dots |\log_{k_d} w|^{\beta_d}, \tag{1.10}$$

where $\log_k w = \log \log_{k-1} w$ for $k \geq 2$ and $\log_1 w = \log w$ can also be considered but only with due care, e.g., subject to the assumption of w being sufficiently large, specifically, with respect to iterated exponentials of k_1, \dots, k_d (as otherwise the repeated logarithm is meaningless due to the possibility of $\log_{k-1} w$ being non-positive hence making $\log_k w$ undefined).¹ Naturally, one can also consider variations of the same theme, e.g., by replacing \log_k with either

$$\log_k^{\parallel} w = \log^{\parallel} \log_{k-1}^{\parallel} w \text{ for } k \geq 2, \quad \log_1^{\parallel} w = |\log w|, \tag{1.11}$$

$$\log_k^+ w = \log^+ \log_{k-1}^+ w \text{ for } k \geq 2, \quad \log_1^+ w = 1 + [\log w]_+. \tag{1.12}$$

However, one needs to observe that the function Φ thus obtained is only \mathcal{C}^1 outside a discrete set (the zero sets of the functions $\log_{k-1} w$) and hence does not lie in $\mathcal{C}^1(\mathbb{R}, \mathbb{R})$ as required.

Another related and yet more general form of Yamabe type equations is the Einstein-scalar field Lichnerowicz equation (see Choquet-Bruhat [15], Chow [16] and Zhang [55]). In the context of smooth metric measure spaces a generalization of the Einstein-scalar field Lichnerowicz equation with space-time dependent coefficients can be described as:

$$\mathcal{L}_\phi[w] = \frac{\partial w}{\partial t} - \Delta_\phi w = A(x, t)w^p + B(x, t)w^q + C(x, t)w \log w, \tag{1.13}$$

$$\mathcal{L}_\phi^a[w] = \left[\frac{\partial}{\partial t} - a(x, t) - \Delta_\phi \right] w = A(x, t)e^{2w} + B(x, t)e^{-2w} + C(x, t). \tag{1.14}$$

For gradient estimates, Harnack inequalities, Liouville type theorems and other related results in this direction see [17, 31, 42, 43, 54] and the references therein.

Moving on to the evolving case the time dependence of the metric-potential pair adds further complications and technical details as far as gradient estimates are concerned. Here the case of the weighted heat equation under the Perelman-Ricci

¹This point does not seem to have been taken into account before.

flow, generalizing in turn, the heat equation under the Ricci flow to the setting to smooth metric measure spaces, given by the system

$$\begin{cases} \mathcal{L}_\phi^a[w](x, t) = \left[\frac{\partial}{\partial t} - a(x, t) - \Delta_\phi \right] w(x, t) = 0, \\ \frac{1}{2} \frac{\partial g}{\partial t}(x, t) + Ric_\phi(g)(x, t) = 0, \end{cases} \tag{1.15}$$

has been considered by many authors (see, e.g., [4, 12, 16, 23, 30, 35, 36, 39, 47, 50, 55]).

System (1.1) can be seen as a generalization of (1.15) in two important ways. Firstly, the weighted linear heat equation is replaced by its nonlinear counterpart where the nonlinearity takes a considerably general formulation. Secondly, the Perelman–Ricci flow (with equality) is replaced by the (k, m) -super Perelman–Ricci flow (with inequality) which is equally a substantial and far reaching generalization (see [37, 38, 42, 43, 47]).

Let us end this introduction by describing the plan of the paper. In § 2 we fix notation and introduce some key quantities and bounds that will be used throughout the paper. In § 3 we gather some background on smooth metric measure spaces and prove some preliminary results on the evolution of geometric quantities of interest for time-dependent metrics and potentials. Section 4 which is the heart of the paper is devoted to the proof of the differential Harnack estimate in theorem 4.1 and its global version in theorem 4.5. Here as a consequence, we also prove a parabolic Harnack inequality for system (1.1). In § 5, we turn into the static case and present the counterparts of the Li–Yau estimates in the static context. We also prove a Liouville type constancy result and present some consequences of it. Finally, in § 6 we prove a Hamilton-type curvature free estimate and bounds using a different set of ideas. In this section, we assume the manifold is closed.

2. Local and global γ -quantities associated with the nonlinearity \mathcal{G}

To a given nonlinearity $\mathcal{G} = \mathcal{G}(t, x, w)$, we associate certain γ -quantities as defined below that will appear in different stages of the analysis and serve as bounds in various estimates. In order to describe these, for \mathcal{G} of class \mathcal{C}^2 and constant μ we set,

$$A_{\mathcal{G}}^\mu(t, x, w) = [-\mu w \mathcal{G}_{ww}(t, x, w) + \mathcal{G}_w(t, x, w) - w^{-1} \mathcal{G}(t, x, w)]_+, \tag{2.1}$$

$$B_{\mathcal{G}}^\mu(t, x, w) = |\mu \mathcal{G}_{xw}(t, x, w) - w^{-1} \mathcal{G}_x(t, x, w)|, \tag{2.2}$$

$$C_{\mathcal{G}}(t, x, w) = [\mathcal{G}_w(t, x, w) - w^{-1} \mathcal{G}(t, x, w)]_+, \tag{2.3}$$

$$D_{\mathcal{G}}(t, x, w) = [-w^{-1} \Delta_\phi \mathcal{G}^x(t, x, w)]_+. \tag{2.4}$$

Here and below $z = z_+ + z_-$ with $z_+ = \max(z, 0)$ and $z_- = \min(z, 0)$. The subscripts in \mathcal{G} stand for the partial derivatives in the respective arguments and $\mathcal{G}^x : x \mapsto \mathcal{G}(t, x, w)$ denotes the function obtained by freezing the variables t, w and viewing \mathcal{G} as a function of x only. (Thus in particular, we speak of $\nabla \mathcal{G}^x$ and $\Delta_\phi \mathcal{G}^x$.)

Fixing a reference point $x_0 \in M$, we denote by $d = d(x, x_0, t)$ the Riemannian distance between x and x_0 with respect to the evolving metric $g = g(t)$. We write $r = r(x, x_0, t)$ for the geodesic radial variable measuring the distance between x and x_0 at time $t > 0$. For $R > 0, T > 0$, we define $Q_{R,T}(x_0) \equiv \{(x, t) : d(x, x_0, t) \leq R, 0 \leq t \leq T\} \subset M \times [0, T]$ and for $0 < t \leq T$, we denote by $\mathcal{B}_r(x_0) \subset M$ the geodesic ball of radius $r > 0$ centred at x_0 . When the choice of the point x_0 is clear from the context, we often abbreviate and write $d(x, t), r(x, t)$ or $\mathcal{B}_r, Q_{R,T}$ respectively.

Having the above notation in place, we now define the four pairs of local and global γ -quantities associated with a given $\mathcal{G} = \mathcal{G}(t, x, w)$ and $w = w(x, t)$ ($x \in M, 0 \leq t \leq T$) by writing for fixed $x_0 \in M, R > 0$ and $T > 0$:

$$\gamma_A^{\mathcal{G},\mu}(R) = \sup_{\Theta_{R,T}} A_{\mathcal{G}}^{\mu}(t, x, w), \quad \gamma_A^{\mathcal{G},\mu} = \sup_{M \times [0,T]} A_{\mathcal{G}}^{\mu}(t, x, w), \tag{2.5}$$

$$\gamma_B^{\mathcal{G},\mu}(R) = \sup_{\Theta_{R,T}} B_{\mathcal{G}}^{\mu}(t, x, w), \quad \gamma_B^{\mathcal{G},\mu} = \sup_{M \times [0,T]} B_{\mathcal{G}}^{\mu}(t, x, w), \tag{2.6}$$

$$\gamma_C^{\mathcal{G}}(R) = \sup_{\Theta_{R,T}} C_{\mathcal{G}}(t, x, w), \quad \gamma_C^{\mathcal{G}} = \sup_{M \times [0,T]} C_{\mathcal{G}}(t, x, w), \tag{2.7}$$

$$\gamma_D^{\mathcal{G}}(R) = \sup_{\Theta_{R,T}} D_{\mathcal{G}}(t, x, w), \quad \gamma_D^{\mathcal{G}} = \sup_{M \times [0,T]} D_{\mathcal{G}}(t, x, w). \tag{2.8}$$

Here $\Theta_{R,T} = \{(t, x, w) : (x, t) \in Q_{R,T}, \underline{w} \leq w \leq \bar{w}\} \subset [0, T] \times M \times (0, \infty)$ where \bar{w}, \underline{w} denote the maximum and minimum of w on the compact space–time cylinder $Q_{R,T}$. Note that in the particular case $\mathcal{G}(t, x, w) = \mathbf{a}(x, t)w$ with \mathbf{a} of class \mathcal{C}^2 , we have:

$$A_{\mathbf{a}w}^{\mu}(t, x, w) \equiv 0, \quad B_{\mathbf{a}w}^{\mu}(t, x, w) = |(\mu - 1)\nabla \mathbf{a}(x, t)|, \tag{2.9}$$

$$C_{\mathbf{a}w}(t, x, w) \equiv 0, \quad D_{\mathbf{a}w}(x, t, w) = [-\Delta_{\phi} \mathbf{a}(x, t)]_+, \tag{2.10}$$

and subsequently

$$\gamma_B^{\mathbf{a}w,\mu}(R) = |\mu - 1| \sup_{Q_{R,T}} |\nabla \mathbf{a}(x, t)| \quad \gamma_B^{\mathbf{a}w,\mu} = |\mu - 1| \sup_{M \times [0,T]} |\nabla \mathbf{a}(x, t)|, \tag{2.11}$$

$$\gamma_D^{\mathbf{a}w}(R) = \sup_{Q_{R,T}} [-\Delta_{\phi} \mathbf{a}(x, t)]_+, \quad \gamma_D^{\mathbf{a}w} = \sup_{M \times [0,T]} [-\Delta_{\phi} \mathbf{a}(x, t)]_+, \tag{2.12}$$

with all the remaining γ -quantities being zero.

3. Bakry–Émery tensors and weighted Bochner–Weitzenböck formula

By a smooth metric measure space, we understand a triple $(M, g, d\sigma)$ in which (M, g) is a complete Riemannian manifold, $d\sigma = e^{-\phi} dv_g$ is a weighted measure associated with the potential ϕ , and dv_g is the standard Riemannian volume measure. Both the metric tensor g and the potential ϕ are assumed to be of class \mathcal{C}^2 . The ϕ -Laplacian (1.2) is a symmetric Markov diffusion operator with respect to the

invariant measure $d\sigma$ and

$$\mathcal{R}ic_\phi^m(g) = \mathcal{R}ic(g) + \nabla\nabla\phi - \frac{\nabla\phi \otimes \nabla\phi}{m-n}, \tag{3.1}$$

is the Bakry–Émery generalized Ricci curvature of the triple $(M, g, d\sigma)$. Here $\mathcal{R}ic(g)$ denotes the Riemannian Ricci curvature of g , $\nabla\nabla\phi = \text{Hess}(\phi)$ is the Hessian of ϕ , and $m \geq n$ is a constant (see [6–8]). For the sake of clarity, we point out that when $m = n$, by convention, ϕ is only allowed to be a constant, thus giving $\mathcal{R}ic_\phi^m(g) = \mathcal{R}ic(g)$, whereas, by formally passing to the limit $m \rightarrow \infty$ in (3.1) we can also set,

$$\mathcal{R}ic_\phi^\infty(g) = \mathcal{R}ic(g) + \nabla\nabla\phi := \mathcal{R}ic_\phi(g). \tag{3.2}$$

The following identity generalizing the classical Bochner–Weitzenböck formula in the Riemannian context to the smooth metric measure space context plays an important role throughout the paper (see [3, 28, 45, 46, 51]).

LEMMA 3.1 Weighted Bochner–Weitzenböck formula. *With Δ_ϕ and $\mathcal{R}ic_\phi(g)$ as above, for any function $h \in \mathcal{C}^3(M)$ we have,*

$$\frac{1}{2}\Delta_\phi|\nabla h|^2 = |\nabla\nabla h|^2 + \langle \nabla h, \nabla\Delta_\phi h \rangle + \mathcal{R}ic_\phi(\nabla h, \nabla h). \tag{3.3}$$

Next, making note of $(\Delta h)^2 \leq n|\nabla\nabla h|^2$ and recalling $\Delta_\phi h = \Delta h - \langle \nabla\phi, \nabla h \rangle$ it is easily seen that

$$|\nabla\nabla h|^2 + \frac{\langle \nabla\phi, \nabla h \rangle^2}{m-n} \geq \frac{(\Delta h)^2}{n} + \frac{\langle \nabla\phi, \nabla h \rangle^2}{m-n} \geq \frac{(\Delta h - \langle \nabla\phi, \nabla h \rangle)^2}{m} = \frac{(\Delta_\phi h)^2}{m}. \tag{3.4}$$

Therefore, from (3.1) and (3.3) it follows that

$$\frac{1}{2}\Delta_\phi|\nabla h|^2 \geq \frac{1}{m}(\Delta_\phi h)^2 + \langle \nabla h, \nabla\Delta_\phi h \rangle + \mathcal{R}ic_\phi^m(\nabla h, \nabla h). \tag{3.5}$$

Let us now present two useful identities on evolutionary metric-potential pairs that will be utilized later. For notational convenience we hereafter denote the metric time derivative tensor, i.e., the speed, as,

$$\frac{\partial g}{\partial t}(x, t) = 2\mathcal{S}(x, t), \quad (x, t) \in M \times (0, T), \tag{3.6}$$

(the factor 2 is only for notational convenience). In the following lemmas, we assume that the evolutionary metric g and potential ϕ are of class \mathcal{C}^2 in the space–time variables (x, t) .

LEMMA 3.2. *With notation (3.6) in place, for any pair of space–time functions $U = U(x, t)$ and $V = V(x, t)$ of class \mathcal{C}^1 , we have*

$$\partial_t \langle \nabla U, \nabla V \rangle = -2\mathcal{S}(\nabla U, \nabla V) + \langle \nabla\partial_t U, \nabla V \rangle + \langle \nabla U, \nabla\partial_t V \rangle. \tag{3.7}$$

In particular $\partial_t|\nabla U|^2 = -2\mathcal{S}(\nabla U, \nabla U) + 2\langle \nabla U, \nabla\partial_t U \rangle$.

Proof. This follows by first writing $\langle \nabla U, \nabla V \rangle = g^{ij} \nabla_i U \nabla_j V$ and then taking ∂_t making note of $\partial_t g^{ij} = -2g^{ik} g^{j\ell} \mathcal{S}_{k\ell} = -2\mathcal{S}^{ij}$. The second identity follows from the first one by setting $V = U$. \square

LEMMA 3.3. *With notation (3.6) in place, for any space–time function $U = U(x, t)$ of class \mathcal{C}^2 , we have*

$$\begin{aligned} \partial_t(\Delta_\phi U) &= \Delta_\phi(\partial_t U) - \langle 2\mathcal{S}, \nabla \nabla U \rangle - \langle 2\operatorname{div} \mathcal{S} - \nabla(\operatorname{Tr}_g \mathcal{S}), \nabla U \rangle \\ &\quad - \langle \nabla \partial_t \phi, \nabla U \rangle + 2\mathcal{S}(\nabla \phi, \nabla U). \end{aligned} \tag{3.8}$$

Proof. Let us first consider the case where the potential ϕ is a constant (thus $\Delta_\phi = \Delta$). Indeed here we have the identity

$$\partial_t \Delta U = \Delta \partial_t U - \langle 2\mathcal{S}, \nabla \nabla U \rangle - \langle 2\operatorname{div} \mathcal{S} - \nabla(\operatorname{Tr}_g \mathcal{S}), \nabla U \rangle. \tag{3.9}$$

Now to justify (3.8) we proceed by directly calculating $\partial_t \Delta_\phi U = \partial_t(\Delta U - \langle \nabla \phi, \nabla U \rangle)$ whilst making note of (3.7) and (3.9). Hence we can write

$$\begin{aligned} \partial_t \Delta_\phi U &= \Delta(\partial_t U) - \langle 2\mathcal{S}, \nabla \nabla U \rangle - \langle 2\operatorname{div} \mathcal{S} - \nabla(\operatorname{Tr}_g \mathcal{S}), \nabla U \rangle \\ &\quad - \langle \nabla \partial_t \phi, \nabla U \rangle - \langle \nabla \phi, \nabla \partial_t U \rangle + 2\mathcal{S}(\nabla \phi, \nabla U) \\ &= \Delta_\phi(\partial_t U) - \langle 2\mathcal{S}, \nabla \nabla U \rangle - \langle 2\operatorname{div} \mathcal{S} - \nabla(\operatorname{Tr}_g \mathcal{S}), \nabla U \rangle \\ &\quad - \langle \nabla \partial_t \phi, \nabla U \rangle + 2\mathcal{S}(\nabla \phi, \nabla U), \end{aligned} \tag{3.10}$$

which is the required conclusion. The proof is thus complete. \square

4. A differential Harnack estimate for system (1.1) under (k, m)-super Perelman–Ricci flow

In this section, we formulate and prove a differential Harnack estimate for positive smooth solutions to the nonlinear parabolic equation $(\partial_t - \mathfrak{a}(x, t) - \Delta_\phi)w = \mathcal{G}(t, x, w)$ where the metric and potential evolves under a (k, m)-super Perelman–Ricci flow. This means that the pair (g, ϕ) forms a complete smooth solution to the flow inequality (4.1) [with $n \leq m < \infty$, $(x, t) \in M \times [0, T]$ and the choice of constant $k = k_m = (m - 1)k + k^L$ (see (4.2) and (4.3) below)]

$$\begin{cases} \frac{1}{2} \frac{\partial g}{\partial t}(x, t) + \mathcal{R}ic_\phi^m(g)(x, t) \geq -kg(x, t), \\ \mathcal{R}ic_\phi^m(g)(x, t) = \mathcal{R}ic(g)(x, t) + \nabla_{g(t)} \nabla_{g(t)} \phi(x, t) - \frac{\nabla_{g(t)} \phi \otimes \nabla_{g(t)} \phi}{m - n}(x, t). \end{cases} \tag{4.1}$$

For future reference we also make note of the following bounds relating to the metric–potential pair (g, ϕ) [recall (3.6)]. For suitable constants $k^L, k^U, k^\nabla \geq 0, \ell_1, \ell_2 \geq 0$:

$$-k^L g \leq \mathcal{S} \leq k^U g, \quad |\nabla \mathcal{S}| \leq k^\nabla, \tag{4.2}$$

$$|\nabla \phi| \leq \ell_1, \quad |\nabla \partial_t \phi| \leq \ell_2. \tag{4.3}$$

THEOREM 4.1. *Let $(M, g, d\sigma)$ be a smooth metric measure space with $d\sigma = e^{-\phi}dv_g$ and time dependent metric-potential pair (g, ϕ) of class \mathcal{C}^2 . Assume $\text{Ric}_\phi^m(g) \geq -(m-1)kg$ in $Q_{2R,T}$ for some $m \geq n, k \geq 0$ and $R, T > 0$ and bounds (4.2)–(4.3) hold in $Q_{2R,T}$. If $w = w(x, t)$ is a positive solution to (1.1), then for every $\mu > 1, \varepsilon \in (0, 1)$ and for all (x, t) in $Q_{R,T}$ with $t > 0$ we have the gradient estimate*

$$\frac{|\nabla w|^2}{\mu w^2} - \frac{\partial_t w}{w} + \frac{\mathcal{G}(t, x, w)}{w} + \mathbf{a}(x, t) \leq \frac{m\mu}{t} + m\mu c_1 k^L + \mathbf{L} + m\mu\gamma_{\mathbf{C}}^{\mathcal{G}}(2R) + \mathbf{M}_a^{\mathcal{G}}. \tag{4.4}$$

The quantities appearing on the right-hand side of bound (4.4) are given respectively by

$$\mathbf{L} = \frac{m\mu}{R^2} \left[\frac{mc_1^2\mu^2}{2(\mu-1)} + c_2 + (m-1)c_1(1 + R\sqrt{k}) + 2c_1^2 \right] \tag{4.5}$$

and

$$\begin{aligned} \mathbf{M}_a^{\mathcal{G}} = \sqrt{m} & \left\{ n\mu^2(k^L + k^U)^2 + 2n\mu^2k^\nabla + \mu\gamma_{\mathbf{D}}^{\mathcal{G}}(2R) + \mu\gamma_{\mathbf{D}}^{\text{aw}}(2R) \right. \\ & + \frac{4m\mu^2[(m-1)k + (\mu-1)k^U + k^\nabla + \gamma_{\mathbf{A}}^{\mathcal{G},\mu}(2R)/2]^2}{4(1-\varepsilon)(\mu-1)^2} \\ & \left. + \frac{3}{4} \left[\frac{m\mu^2[\mu\ell_2 + 2\mu k^L\ell_1 + 2\gamma_{\mathbf{B}}^{\mathcal{G},\mu}(2R) + 2\gamma_{\mathbf{B}}^{\text{aw},\mu}(2R)]^4}{4\varepsilon(\mu-1)^2} \right]^{1/3} \right\}^{1/2}. \tag{4.6} \end{aligned}$$

The γ -quantities in (4.4) and (4.6) are as in (2.5)–(2.8) and (2.11)–(2.12) (with $2R$ replacing R) and the constants $c_1, c_2 > 0$ in (4.5) are as in (4.19) in lemma 4.4.

As the proof of the theorem is quite involved and requires several intermediate steps, for the sake of reader’s convenience we present this in several stages. The first task is to introduce a Harnack quantity built out of the solution w and consider its evolution under the weighted heat operator.

LEMMA 4.2. *Let w be a positive solution to the equation $\mathcal{L}_\phi^{\mathfrak{a}}[w] = (\partial_t - \mathbf{a}(x, t) - \Delta_\phi)w = \mathcal{G}(t, x, w)$ and let $F_a^{\mathcal{G}} = F_a^{\mathcal{G}}(x, t)$ be defined by*

$$F_a^{\mathcal{G}}(x, t) = t[|\nabla f|^2 - \mu\partial_t f + \mu\mathbf{a}(x, t) + \mu e^{-f}\mathcal{G}(t, x, e^f)], \quad t \geq 0, \tag{4.7}$$

where $f = \log w$ and $\mu > 1$ is a fixed constant. Suppose that the metric-potential pair (g, ϕ) is time dependent and of class \mathcal{C}^2 . Then $F_a^{\mathcal{G}}$ satisfies

$$\begin{aligned}
 (\Delta_\phi - \partial_t)[F_a^{\mathcal{G}}] &= 2t|\nabla\nabla f|^2 - 2\langle\nabla f, \nabla F_a^{\mathcal{G}}\rangle + 2t\mathcal{R}ic_\phi^m(\nabla f, \nabla f) \\
 &\quad - 2t(\mu - 1)\mathcal{S}(\nabla f, \nabla f) + 2t\langle\nabla\phi, \nabla f\rangle^2/(m - n) - F_a^{\mathcal{G}}/t \\
 &\quad - 2\mu t[\langle\mathcal{S}, \nabla\nabla 2f\rangle + \langle\text{div}\mathcal{S} - (1/2)\nabla(\text{Tr}_g\mathcal{S}), \nabla f\rangle] \\
 &\quad - \mu t[\langle\nabla\partial_t\phi, \nabla f\rangle - 2\mathcal{S}(\nabla\phi, \nabla f)] \\
 &\quad + 2t(\mu - 1)\langle\nabla f, \nabla\mathbf{a}(x, t)\rangle + \mu t\Delta_\phi[\mathbf{a}(x, t)] \\
 &\quad + 2t(\mu - 1)\langle\nabla f, \nabla[e^{-f}\mathcal{G}(t, x, e^f)]\rangle + \mu t\Delta_\phi[e^{-f}\mathcal{G}(t, x, e^f)]. \tag{4.8}
 \end{aligned}$$

Proof. Referring to the equation for w an easy calculation shows that f satisfies the equation

$$\mathcal{L}_\phi[f] = (\partial_t - \Delta_\phi)f = |\nabla f|^2 + e^{-f}\mathcal{G}(t, x, e^f) + \mathbf{a}(x, t). \tag{4.9}$$

Moreover, using (4.7) and (4.9) it is a straightforward matter to see that the following relation holds between $F_a^{\mathcal{G}}$ and $\Delta_\phi f$:

$$\begin{aligned}
 \Delta_\phi f &= -[\mu^{-1}|\nabla f|^2 - \partial_t f + \mathbf{a}(x, t) + e^{-f}\mathcal{G}(t, x, e^f)] - (1 - \mu^{-1})|\nabla f|^2 \\
 &= -F_a^{\mathcal{G}}/(\mu t) - (1 - \mu^{-1})|\nabla f|^2, \quad t > 0. \tag{4.10}
 \end{aligned}$$

We next calculate the different ingredients needed in the application of the weighted heat operator $\mathcal{L}_\phi = \partial_t - \Delta_\phi$ to the Harnack quantity in (4.7). To this end we first note that

$$\Delta_\phi F_a^{\mathcal{G}} = t(\Delta_\phi|\nabla f|^2 - \mu\Delta_\phi(\partial_t f) + \mu\Delta_\phi\mathbf{a}(x, t) + \mu\Delta_\phi[e^{-f}\mathcal{G}(t, x, e^f)]). \tag{4.11}$$

Recalling the weighted Bochner–Weitzenböck formula in lemma 3.1 as applied to f and making use of (3.8) in lemma 3.3 we then have

$$\begin{aligned}
 \Delta_\phi F_a^{\mathcal{G}} &= t[2|\nabla\nabla f|^2 + 2\langle\nabla f, \nabla\Delta_\phi f\rangle + 2\mathcal{R}ic_\phi^m(\nabla f, \nabla f) + 2\langle\nabla\phi, \nabla f\rangle^2/(m - n)] \\
 &\quad - \mu t\partial_t(\Delta_\phi f) - 2\mu t[\langle\mathcal{S}, \nabla\nabla f\rangle + \langle\text{div}\mathcal{S} - (1/2)\nabla(\text{Tr}_g\mathcal{S}), \nabla f\rangle] \tag{4.12} \\
 &\quad - \mu t[\langle\nabla\partial_t\phi, \nabla f\rangle - 2\mathcal{S}(\nabla\phi, \nabla f)] + \mu t\Delta_\phi[\mathbf{a}(x, t)] + \mu t\Delta_\phi[e^{-f}\mathcal{G}(t, x, e^f)].
 \end{aligned}$$

Now referring to the sum on the right, the contributions of the second and fifth terms modulo a factor t can be simplified and rewritten upon using (4.9) and (4.10)

as,

$$\begin{aligned}
 & 2\langle \nabla f, \nabla \Delta_\phi f \rangle - \mu \partial_t (\Delta_\phi f) \\
 &= 2\langle \nabla f, \nabla \Delta_\phi f \rangle + 2(\mu - 1)\langle \nabla f, \nabla [\Delta_\phi f + |\nabla f|^2 + e^{-f}\mathcal{G}(t, x, e^f) + \mathbf{a}(x, t)] \rangle \\
 &\quad + (t\partial_t F_a^\mathcal{G} - F_a^\mathcal{G})/t^2 - 2(\mu - 1)\mathcal{S}(\nabla f, \nabla f) \\
 &= 2\langle \nabla f, \nabla [-F_a^\mathcal{G}/t - (\mu - 1)|\nabla f|^2] \rangle + 2(\mu - 1)\langle \nabla f, \nabla |\nabla f|^2 \rangle + (t\partial_t F_a^\mathcal{G} - F_a^\mathcal{G})/t^2 \\
 &\quad + 2(\mu - 1)\langle \nabla f, \nabla \mathbf{a}(x, t) \rangle + 2(\mu - 1)\langle \nabla f, \nabla [e^{-f}\mathcal{G}(t, x, e^f)] \rangle \\
 &\quad - 2(\mu - 1)\mathcal{S}(\nabla f, \nabla f) \\
 &= 2(\mu - 1)[\langle \nabla f, \nabla \mathbf{a}(x, t) \rangle + \langle \nabla f, \nabla [e^{-f}\mathcal{G}(t, x, e^f)] \rangle - \mathcal{S}(\nabla f, \nabla f)] \\
 &\quad + (t\partial_t F_a^\mathcal{G} - F_a^\mathcal{G})/t^2 - 2\langle \nabla f, \nabla F_a^\mathcal{G} \rangle/t. \tag{4.13}
 \end{aligned}$$

Therefore, substituting this expression back into (4.12) and rearranging terms lead to

$$\begin{aligned}
 (\Delta_\phi - \partial_t)[F_a^\mathcal{G}] &= 2t|\nabla \nabla f|^2 + 2t(\mu - 1)[\langle \nabla f, \nabla \mathbf{a} \rangle \\
 &\quad + \langle \nabla f, \nabla [e^{-f}\mathcal{G}(t, x, e^f)] \rangle - \mathcal{S}(\nabla f, \nabla f)] \\
 &\quad - F_a^\mathcal{G}/t - 2\langle \nabla f, \nabla F_a^\mathcal{G} \rangle \\
 &\quad + 2t\mathcal{R}ic_\phi^m(\nabla f, \nabla f) + 2t\langle \nabla \phi, \nabla f \rangle^2/(m - n) \\
 &\quad - 2\mu t\langle \mathcal{S}, \nabla \nabla f \rangle - \mu t\langle 2\text{div} \mathcal{S} - \nabla(\text{Tr}_g \mathcal{S}), \nabla f \rangle - \mu t\langle \nabla \partial_t \phi, \nabla f \rangle \\
 &\quad + 2\mu t\mathcal{S}(\nabla \phi, \nabla f) + \mu t\Delta_\phi \mathbf{a} + \mu t\Delta_\phi [e^{-f}\mathcal{G}(t, x, e^f)] \tag{4.14}
 \end{aligned}$$

which is the desired conclusion. □

LEMMA 4.3. Let w be a positive solution to $\mathcal{L}_\phi^3[f] = (\partial_t - \mathbf{a}(x, t) - \Delta_\phi)w = \mathcal{G}(t, x, w)$ and let F be as in (4.7). Assume the metric-potential pair (g, ϕ) is time dependent and of class \mathcal{C}^2 . Moreover assume the bounds $\mathcal{R}ic_\phi^m(g) \geq -(m - 1)kg$ and

$$-k^L g \leq \mathcal{S} \leq k^U g, \quad |\nabla \mathcal{S}| \leq k^\nabla, \tag{4.15}$$

for suitable k, \underline{k}^L, k^U and $k^\nabla \geq 0$. Then

$$\begin{aligned}
 (\Delta_\phi - \partial_t)[F_a^\mathcal{G}] &\geq t(\Delta_\phi f)^2/m - F_a^\mathcal{G}/t - 2\langle \nabla f, \nabla F_a^\mathcal{G} \rangle \\
 &\quad - 2t[(m - 1)k + (\mu - 1)k^U]|\nabla f|^2 \\
 &\quad - \mu^2 n t(k^L + k^U)^2 - 3\mu t\sqrt{n}k^\nabla |\nabla f| \\
 &\quad - \mu t\langle \nabla \partial_t \phi, \nabla f \rangle + 2\mu t\mathcal{S}(\nabla \phi, \nabla f) \\
 &\quad + 2(\mu - 1)t\langle \nabla f, \nabla \mathbf{a}(x, t) \rangle + \mu t\Delta_\phi [\mathbf{a}(x, t)] \\
 &\quad + 2(\mu - 1)t\langle \nabla f, \nabla [e^{-f}\mathcal{G}(t, x, e^f)] \rangle + \mu t\Delta_\phi [e^{-f}\mathcal{G}(t, x, e^f)]. \tag{4.16}
 \end{aligned}$$

Proof. From the upper and lower bounds on \mathcal{S} in the sense of symmetric 2-tensors in (3.6)–(4.15) it follows that $|\mathcal{S}|^2 \leq (k^L + k^U)^2 |g|^2 = n(k^L + k^U)^2$ and hence

$$2|\mu \langle \mathcal{S}, \nabla \nabla f \rangle| \leq |\nabla \nabla f|^2 + \mu^2 |\mathcal{S}|^2 \leq |\nabla \nabla f|^2 + n\mu^2 (k^L + k^U)^2. \tag{4.17}$$

Moreover by virtue of the assumption $|\nabla \mathcal{S}| \leq k^\nabla$ we have

$$\begin{aligned} |2\operatorname{div} \mathcal{S} - \nabla(\operatorname{Tr}_g \mathcal{S})| &= |2g^{ij} \nabla_i \mathcal{S}_{j\ell} - g^{ij} \nabla_\ell \mathcal{S}_{ij}| = |g^{ij} (2\nabla_i \mathcal{S}_{j\ell} - \nabla_\ell \mathcal{S}_{ij})| \\ &\leq 3|g| |\nabla \mathcal{S}| \leq 3\sqrt{n} k^\nabla. \end{aligned} \tag{4.18}$$

Now the conclusion follows at once by referring (4.8) in lemma 4.2, making note of the bound $|\nabla \nabla f|^2 + \langle \nabla \phi, \nabla f \rangle^2 / (m - n) \geq (\Delta f)^2 / n + \langle \nabla \phi, \nabla f \rangle^2 / (m - n) \geq (\Delta_\phi f)^2 / m$, and the Bakry–Émery curvature lower bound $\mathcal{R}ic_\phi^m \geq -(m - 1)kg$ in the lemma. \square

For the purpose of localization we shall make use of standard spatial cut-off functions. To this end we note the following lemma gathering together some of the main properties of a profile function defined on the half-line $s \geq 0$ as needed later (see [4, 11, 28, 29, 54]).

LEMMA 4.4. *There exists a function $\bar{\psi} : [0, \infty) \rightarrow \mathbb{R}$ satisfying the following properties:*

- (i) $\bar{\psi}$ is of class $\mathcal{C}^2[0, \infty)$.
- (ii) $0 \leq \bar{\psi}(s) \leq 1$ for $0 \leq s < \infty$ and $\bar{\psi} \equiv 1$ on $[0, 1]$ and $\bar{\psi} \equiv 0$ on $[2, \infty)$.
- (iii) $\bar{\psi}' \leq 0$ and so $\bar{\psi}$ is non-increasing and for suitable constants $c_1, c_2 > 0$ we have the global bounds

$$-c_1 \bar{\psi}^{1/2} \leq \bar{\psi}' \leq 0, \text{ and } \bar{\psi}'' \geq -c_2. \tag{4.19}$$

Now, we pick a reference point $x_0 \in M$, fix $R, T > 0$ and $0 < \tau \leq T$ and then with $r(x, t)$ denoting the geodesic radial variable with respect to x_0 at time t , set

$$\psi(x, t) = \bar{\psi}(r(x, t)/R), \quad x \in M, 0 \leq t \leq T. \tag{4.20}$$

It is evident that the resulting function ψ satisfies $\psi \equiv 1$ for when $0 \leq r(x, t) \leq R$ and $\psi \equiv 0$ for when $r(x, t) \geq 2R$. Additionally from (4.20) we have $\nabla \psi = (\bar{\psi}'/R) \nabla r$ and $\Delta \psi = \bar{\psi}' |\nabla r|^2 / R^2 + \bar{\psi}'' \Delta r / R$ and so $\Delta_\phi \psi = \Delta \psi - \langle \nabla \phi, \nabla \psi \rangle = \bar{\psi}'' |\nabla r|^2 / R^2 + \bar{\psi}' \Delta_\phi r / R$. In particular $\nabla \psi, \Delta_\phi \psi$ vanish outside the space–time set $R \leq r(x, t) \leq 2R$.

Proof of theorem 4.1. Consider the localized function $\psi F_a^{\mathcal{G}}$ where $F_a^{\mathcal{G}}$ is the Harnack quantity in (4.7). Let (x_1, t_1) denote the point where this function attains its maximum over the compact cylinder $\{r(x, t) \leq 2R, 0 \leq t \leq \tau\}$. We assume $[\psi F_a^{\mathcal{G}}](x_1, t_1) > 0$ as otherwise the estimate follows from $F_a^{\mathcal{G}} \leq 0$. So in particular $t_1 > 0$ and $r(x_1, t_1) < 2R$ and therefore, at the maximum point (x_1, t_1) , we

have the relations

$$\begin{cases} \partial_t(\psi F_a^{\mathcal{G}}) \geq 0, \\ \nabla(\psi F_a^{\mathcal{G}}) = 0, \\ \Delta(\psi F_a^{\mathcal{G}}) \leq 0, \\ \Delta_\phi(\psi F_a^{\mathcal{G}}) \leq 0. \end{cases} \tag{4.21}$$

Utilizing the product identity for the ϕ -Laplacian as applied to the localized function $\psi F_a^{\mathcal{G}}$ we can write

$$\Delta_\phi(\psi F_a^{\mathcal{G}}) = F_a^{\mathcal{G}} \Delta_\phi \psi + 2\langle \nabla \psi, \nabla F_a^{\mathcal{G}} \rangle + \psi \Delta_\phi F_a^{\mathcal{G}}, \tag{4.22}$$

and so making note of relations (4.21) at the maximum point (x_1, t_1) we can further deduce

$$\begin{aligned} 0 &\geq F_a^{\mathcal{G}} \Delta_\phi \psi + 2\langle \nabla \psi, \nabla F_a^{\mathcal{G}} \rangle + \psi \Delta_\phi F_a^{\mathcal{G}} \\ &\geq F_a^{\mathcal{G}} \Delta_\phi \psi + (2/\psi)\langle \nabla \psi, \nabla(\psi F_a^{\mathcal{G}}) \rangle - 2(|\nabla \psi|^2/\psi)F_a^{\mathcal{G}} + \psi \Delta_\phi F_a^{\mathcal{G}} \\ &\geq F_a^{\mathcal{G}} \Delta_\phi \psi - 2(|\nabla \psi|^2/\psi)F_a^{\mathcal{G}} + \psi \Delta_\phi F_a^{\mathcal{G}}. \end{aligned} \tag{4.23}$$

Let us now proceed by bounding the sum on the right-hand side of (4.23) from below. To this end starting with the first term we have

$$\Delta_\phi \psi \geq -\frac{1}{R^2}[c_2 + (m-1)c_1(1 + R\sqrt{k})], \tag{4.24}$$

where $c_1, c_2 > 0$ are as in lemma 4.4. In fact, since $\mathcal{R}ic_\phi^m(g) \geq -(m-1)kg$ it follows from the Wei–Wylie weighted Laplacian comparison theorem, the ϕ -Laplacian relation $\Delta_\phi \psi = \bar{\psi}'|\nabla r|^2/R^2 + \bar{\psi}'\Delta_\phi r/R$ and the bounds in lemma 4.4 that,

$$\begin{aligned} \Delta_\phi \psi &\geq \frac{1}{R^2}\bar{\psi}' + \frac{(m-1)}{R}\bar{\psi}'\sqrt{k}\coth(\sqrt{k}r) \geq \frac{1}{R^2}\bar{\psi}' + \frac{(m-1)}{R}\bar{\psi}'\sqrt{k}\coth(\sqrt{k}R) \\ &\geq \frac{1}{R^2}\bar{\psi}' + (m-1)\frac{1+\sqrt{k}R}{R}\bar{\psi}' \geq -\frac{c_2}{R^2} - (m-1)\frac{c_1}{R}\left(\frac{1}{R} + \sqrt{k}\right), \end{aligned} \tag{4.25}$$

by virtue of $\Delta_\phi r \leq (m-1)\sqrt{k}\coth(\sqrt{k}r)$ and $\Delta_\phi \psi \equiv 0$ outside $R \leq r \leq 2R$. Note that here we have used the monotonicity of $s \mapsto \coth s$ and $s \coth s \leq (1+s)$ for $s > 0$. Onto the second term on the right in (4.23), again by (4.20) and lemma 4.4 we have,

$$\frac{|\nabla \psi|^2}{\psi} = \frac{\bar{\psi}'^2}{\bar{\psi}} \frac{|\nabla r|^2}{R^2} = \left(\frac{\bar{\psi}'}{\sqrt{\bar{\psi}}} \frac{|\nabla r|}{R}\right)^2 \leq \frac{c_1^2}{R^2}. \tag{4.26}$$

Now making use of (4.16), (4.24) and (4.26) and substituting in (4.23) it follows that at the maximum point (x_1, t_1) we have the inequality

$$\begin{aligned}
 0 &\geq F_a^{\mathcal{G}} \Delta_\phi \psi - 2(|\nabla \psi|^2 / \psi) F_a^{\mathcal{G}} + \psi \Delta_\phi F_a^{\mathcal{G}} \\
 &\geq -[c_2 + (m - 1)c_1(1 + R\sqrt{k})] F_a^{\mathcal{G}} / R^2 - 2(c_1^2 / R^2) F_a^{\mathcal{G}} + \psi \partial_t F_a^{\mathcal{G}} \\
 &\quad + \psi[(t_1/m)(\Delta_\phi f)^2 - F_a^{\mathcal{G}}/t_1 - 2\langle \nabla f, \nabla F_a^{\mathcal{G}} \rangle - 2[(m - 1)k + (\mu - 1)k^U] t_1 |\nabla f|^2 \\
 &\quad - \mu^2 n t_1 (k^L + k^U)^2 - 3\sqrt{n} k^\nabla \mu t_1 |\nabla f| - \mu t_1 \langle \nabla \partial_t \phi, \nabla f \rangle + 2\mu t_1 \mathcal{S}(\nabla \phi, \nabla f) \\
 &\quad + 2(\mu - 1)t_1[\langle \nabla f, \nabla \mathbf{a} \rangle + \langle \nabla f, \nabla(e^{-f} \mathcal{G}) \rangle] + \mu t_1 [\Delta_\phi \mathbf{a} + \Delta_\phi(e^{-f} \mathcal{G})]. \tag{4.27}
 \end{aligned}$$

For the sake of convenience in writing here and below we abbreviate the arguments of $\mathbf{a} = \mathbf{a}(x, t)$ and $\mathcal{G} = \mathcal{G}(t, x, e^f)$. The aim is now to bound each of the individual terms in the last inequality. Starting from the third term on the second line $\psi \partial_t F_a^{\mathcal{G}}$, upon recalling (4.20), we have,

$$\partial_t(\psi F_a^{\mathcal{G}}) = \psi \partial_t F_a^{\mathcal{G}} + F_a^{\mathcal{G}} \partial_t \psi = \bar{\psi}(r/R) \partial_t F_a^{\mathcal{G}} + \bar{\psi}'(r/R) F_a^{\mathcal{G}} \partial_t r / R. \tag{4.28}$$

Now as at the maximum point (x_1, t_1) we have $\partial_t(\psi F_a^{\mathcal{G}}) \geq 0$, by restricting to this point, utilizing lemma 4.4 [the left inequality in (4.19)] and the bound $\partial_t r(x, t) \geq -k^L r(x, t)$ [see (4.30)], we can write

$$\begin{aligned}
 \psi \partial_t F_a^{\mathcal{G}} &\geq -F_a^{\mathcal{G}} \partial_t \psi = -\bar{\psi}'(r/R) F_a^{\mathcal{G}} \partial_t r / R \\
 &\geq r k^L F_a^{\mathcal{G}} \bar{\psi}'(r/R) / R \geq -c_1 r k^L \sqrt{\bar{\psi}(r/R) F_a^{\mathcal{G}}} / R \geq -c_1 k^L F_a^{\mathcal{G}}. \tag{4.29}
 \end{aligned}$$

The justification for the lower bound on $\partial_t r(x_1, t_1)$ used above can be given as follows. Fix x and t such that $d(x, x_0, t) < 2R$. Let $X(x_0, x)$ be the set of all minimal geodesics $\zeta = \zeta(s) : [0, 1] \rightarrow M$ with respect to $g(t)$ connecting the reference point $x_0 = \zeta(0)$ to $x = \zeta(1)$ and let $\Gamma(x_0, x)$ be the set of all \mathcal{C}^1 curves connecting x_0 to x . Using the lower bound $\partial_t g = 2\mathcal{S} \geq -2k^L g$ in $Q_{2R,T}$ as given by (4.2) with $k^L \geq 0$ and Lemma B.40 p. 531 in [16] we can write:

$$\begin{aligned}
 \partial_t r(x, t) &= \frac{\partial}{\partial t} d(x, x_0; t) \\
 &= \frac{\partial}{\partial t} \left\{ \inf_{\gamma \in \Gamma(x_0, x)} \int_0^1 |\gamma'(s)|_{g(t)} ds \right\} \\
 &= \frac{\partial}{\partial t} \left\{ \inf_{\gamma \in \Gamma(x_0, x)} \int_0^1 \sqrt{[g(t)](\gamma'(s), \gamma'(s))} ds \right\} \\
 &= \inf_{\zeta \in X(x_0, x)} \int_0^1 \frac{[\partial_t g](\zeta'(s), \zeta'(s))}{2\sqrt{[g(t)](\zeta'(s), \zeta'(s))}} ds = \inf_{\zeta \in X(x_0, x)} \int_0^1 \frac{\mathcal{S}(\zeta'(s), \zeta'(s))}{|\zeta'(s)|_{g(t)}} ds \\
 &\geq \inf_{\zeta \in X(x_0, x)} \int_0^1 -k^L |\zeta'(s)|_{g(t)} ds = \inf_{\zeta \in X(x_0, x)} \left[-k^L \overbrace{\int_0^1 |\zeta'(s)|_{g(t)} ds}^{=r(x,t)} \right] \\
 &\geq -k^L r(x, t) \geq -k^L R, \tag{4.30}
 \end{aligned}$$

which is the desired bound. (We note that this can also be derived using the evolution formula of the geodesic length under geometric flow due to R. Hamilton [24].) Next, in view of $\nabla(\psi F_a^{\mathcal{G}}) = 0$ at (x_1, t_1) we can write,

$$\psi \langle \nabla f, \nabla F_a^{\mathcal{G}} \rangle = -F_a^{\mathcal{G}} \langle \nabla f, \nabla \psi \rangle \leq F_a^{\mathcal{G}} |\nabla f| |\nabla \psi| \leq c_1 (\sqrt{\psi}/R) F_a^{\mathcal{G}} |\nabla f|. \tag{4.31}$$

Likewise, we have $\mathcal{S}(\nabla \phi, \nabla f) \leq k^L |\nabla \phi| |\nabla f|$ and $\langle \nabla \partial_t \phi, \nabla f \rangle \leq |\nabla \partial_t \phi| |\nabla f|$ along with $3k^\nabla \sqrt{n} \mu |\nabla f| \leq 2nk^\nabla \mu^2 + 2k^\nabla |\nabla f|^2$. Hence substituting all the above back in (4.27) it follows that

$$\begin{aligned} 0 \geq & -[c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2] F_a^{\mathcal{G}} / R^2 \\ & - c_1 k^L F_a^{\mathcal{G}} - 2c_1 (\sqrt{\psi}/R) F_a^{\mathcal{G}} |\nabla f| + t_1 (\psi/m) (\Delta_\phi f)^2 \\ & - \psi F_a^{\mathcal{G}} / t_1 - 2t_1 [(m - 1)k + (\mu - 1)k^U] \psi |\nabla f|^2 \\ & - t_1 \psi [\mu^2 n (k^L + k^U)^2 + 2nk^\nabla \mu^2 + 2k^\nabla |\nabla f|^2] \\ & - \mu t_1 \psi [2k^L |\nabla \phi| |\nabla f| + |\nabla \partial_t \phi| |\nabla f|] \\ & + 2t_1 \psi (\mu - 1) [\langle \nabla f, \nabla a \rangle + \langle \nabla f, \nabla (e^{-f} \mathcal{G}) \rangle] \\ & + t_1 \psi \mu [\Delta_\phi a + \Delta_\phi (e^{-f} \mathcal{G})]. \end{aligned} \tag{4.32}$$

Next multiplying (4.32) through by the factor $t_1 \psi(x_1) = t_1 \psi$, making note of (4.9) and rearranging terms gives

$$\begin{aligned} 0 \geq & -t_1 \psi F_a^{\mathcal{G}} [(c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2) / R^2 - \psi^2 F_a^{\mathcal{G}} \\ & - c_1 k^L t_1 \psi F_a^{\mathcal{G}} + t_1^2 (\psi^2 / m) [|\nabla f|^2 + e^{-f} \mathcal{G} + a - \partial_t f]^2 \\ & - 2c_1 t_1 \psi (\sqrt{\psi}/R) F_a^{\mathcal{G}} |\nabla f| - 2t_1^2 [(m - 1)k + (\mu - 1)k^U + k^\nabla] \psi^2 |\nabla f|^2 \\ & - n \mu^2 t_1^2 \psi^2 [(k^L + k^U)^2 + 2k^\nabla] - \mu t_1^2 \psi^2 |\nabla \partial_t \phi| |\nabla f| - 2\mu t_1^2 \psi^2 k^L |\nabla \phi| |\nabla f| \\ & + t_1^2 \psi^2 \{2(\mu - 1) [\langle \nabla f, \nabla a \rangle + \langle \nabla f, \nabla (e^{-f} \mathcal{G}) \rangle] + \mu [\Delta_\phi a + \Delta_\phi (e^{-f} \mathcal{G})]\}. \end{aligned} \tag{4.33}$$

Let us now pause briefly to go through some calculations relating to $\mathcal{G} = \mathcal{G}(t, x, w)$ with $w = e^f$ and $f = f(x, t)$ needed below. Note that the arguments of the functions involved will be abbreviated for the sake of convenience in writing. Firstly,

$$\nabla \mathcal{G} = \mathcal{G}_x + e^f \mathcal{G}_w \nabla f, \quad \mathcal{G}_x = (\mathcal{G}_{x_1}, \dots, \mathcal{G}_{x_n}). \tag{4.34}$$

Upon introducing the notation $\mathcal{G}^x : x \mapsto \mathcal{G}(t, x, w)$ (i.e., viewing \mathcal{G} as a function of x whilst freezing the variable w) we can write

$$\begin{aligned} \Delta \mathcal{G} &= \Delta \mathcal{G}^x + e^f \langle \mathcal{G}_{xw} + e^f |\nabla f|^2 \mathcal{G}_w, \nabla f \rangle \\ &+ e^f \langle \mathcal{G}_{xw}, \nabla f \rangle + e^{2f} |\nabla f|^2 \mathcal{G}_{ww} + e^f \mathcal{G}_w \Delta f \\ &= \Delta \mathcal{G}^x + 2e^f \langle \mathcal{G}_{xw}, \nabla f \rangle + e^f |\nabla f|^2 (\mathcal{G}_w + e^f \mathcal{G}_{ww}) + e^f \mathcal{G}_w \Delta f. \end{aligned} \tag{4.35}$$

Subsequently calculating the ϕ -Laplacian, by utilizing the above fragments we have,

$$\begin{aligned} \Delta_\phi \mathcal{G} &= \Delta \mathcal{G} - \langle \nabla \phi, \nabla \mathcal{G} \rangle = \Delta \mathcal{G} - \langle \nabla \phi, (\mathcal{G}_x + e^f \mathcal{G}_w \nabla f) \rangle \\ &= \Delta \mathcal{G} - \langle \nabla \phi, \mathcal{G}_x \rangle - e^f \mathcal{G}_w \langle \nabla \phi, \nabla f \rangle \\ &= \Delta_\phi \mathcal{G}^x + 2e^f \langle \mathcal{G}_{xw}, \nabla f \rangle + e^f |\nabla f|^2 (\mathcal{G}_w + e^f \mathcal{G}_{ww}) + e^f \mathcal{G}_w \Delta_\phi f. \end{aligned} \tag{4.36}$$

For the sake of future reference we also note that

$$\begin{aligned} \Delta_\phi e^{-f} &= \Delta e^{-f} - \langle \nabla \phi, \nabla e^{-f} \rangle \\ &= -\operatorname{div}(e^{-f} \nabla f) + e^{-f} \langle \nabla \phi, \nabla f \rangle \\ &= -e^{-f} \Delta f + e^{-f} |\nabla f|^2 + e^{-f} \langle \nabla \phi, \nabla f \rangle = -e^{-f} (\Delta_\phi f - |\nabla f|^2). \end{aligned} \tag{4.37}$$

Returning to inequality (4.33) and picking up the estimate from where we left, for the last two terms about \mathcal{G} , we can write

$$\begin{aligned} &2(\mu - 1) \langle \nabla f, \nabla(e^{-f} \mathcal{G}) \rangle + \mu \Delta_\phi(e^{-f} \mathcal{G}) \\ &= 2(\mu - 1) [-e^{-f} \mathcal{G} |\nabla f|^2 + e^{-f} \langle \nabla f, \nabla \mathcal{G} \rangle] \\ &\quad + \mu [e^{-f} \Delta_\phi \mathcal{G} + \mathcal{G} \Delta_\phi e^{-f} - 2e^{-f} \langle \nabla f, \nabla \mathcal{G} \rangle] \\ &= -2(\mu - 1) e^{-f} \mathcal{G} |\nabla f|^2 + 2\mu e^{-f} \langle \nabla f, \nabla \mathcal{G} \rangle \\ &\quad - 2e^{-f} \langle \nabla f, (\mathcal{G}_x + e^f \mathcal{G}_w \nabla f) \rangle + \mu e^{-f} (\Delta_\phi \mathcal{G}^x + 2e^f \langle \mathcal{G}_{xw}, \nabla f \rangle) \\ &\quad + \mu e^{-f} (e^f |\nabla f|^2 (\mathcal{G}_w + e^f \mathcal{G}_{ww}) + e^f \mathcal{G}_w \Delta_\phi f) \\ &\quad + \mu \mathcal{G} e^{-f} (-\Delta_\phi f + |\nabla f|^2) - 2\mu e^{-f} \langle \nabla f, \nabla \mathcal{G} \rangle. \end{aligned} \tag{4.38}$$

As according to (4.10) we have

$$\begin{aligned} [\mu \mathcal{G}_w - \mu \mathcal{G} e^{-f}] \Delta_\phi f &= [-F_a^{\mathcal{G}} / (\mu t_1) - (\mu - 1) |\nabla f|^2 / \mu] [\mu (\mathcal{G}_w - \mathcal{G} e^{-f})] \\ &= -(F_a^{\mathcal{G}} / t_1) (\mathcal{G}_w - \mathcal{G} e^{-f}) - (\mu - 1) |\nabla f|^2 (\mathcal{G}_w - \mathcal{G} e^{-f}), \end{aligned} \tag{4.39}$$

upon substitution back in (4.38) this gives

$$\begin{aligned} &2(\mu - 1) \langle \nabla f, \nabla(e^{-f} \mathcal{G}) \rangle + \mu \Delta_\phi(e^{-f} \mathcal{G}) \\ &= -2(\mu - 1) e^{-f} \mathcal{G} |\nabla f|^2 - 2e^{-f} \langle \nabla f, \mathcal{G}_x \rangle - 2\mathcal{G}_w |\nabla f|^2 \\ &\quad + \mu e^{-f} \Delta_\phi \mathcal{G}^x + 2\mu \langle \mathcal{G}_{xw}, \nabla f \rangle + \mu |\nabla f|^2 \mathcal{G}_w + \mu |\nabla f|^2 e^f \mathcal{G}_{ww} \\ &\quad + \mu \mathcal{G} e^{-f} |\nabla f|^2 - (F_a^{\mathcal{G}} / t_1) (\mathcal{G}_w - \mathcal{G} e^{-f}) - (\mu - 1) |\nabla f|^2 (\mathcal{G}_w - \mathcal{G} e^{-f}) \\ &= |\nabla f|^2 [-2(\mu - 1) e^{-f} \mathcal{G} - 2\mathcal{G}_w + \mu \mathcal{G}_w + \mu e^f \mathcal{G}_{ww}] \\ &\quad + |\nabla f|^2 [\mu e^{-f} \mathcal{G} - (\mu - 1) \mathcal{G}_w + (\mu - 1) \mathcal{G} e^{-f}] \\ &\quad - [2 \langle \nabla f, (e^{-f} \mathcal{G}_x - \mu \mathcal{G}_{xw}) \rangle + (F_a^{\mathcal{G}} / t_1) (\mathcal{G}_w - \mathcal{G} e^{-f}) - \mu e^{-f} \Delta_\phi \mathcal{G}^x]. \end{aligned}$$

Therefore, by taking into account the relevant cancellations, after simplifying terms and using basic inequalities, we can write

$$\begin{aligned}
 & 2(\mu - 1)\langle \nabla f, \nabla(e^{-f}\mathcal{G}) \rangle + \mu\Delta_\phi(e^{-f}\mathcal{G}) \\
 & \geq |\nabla f|^2(e^{-f}\mathcal{G} - \mathcal{G}_w + \mu e^f \mathcal{G}_{ww}) - (F_a^\mathcal{G}/t_1)(\mathcal{G}_w - e^{-f}\mathcal{G}) \\
 & \quad - 2|\nabla f||e^{-f}\mathcal{G}_x - \mu\mathcal{G}_{xw}| + \mu e^{-f}\Delta_\phi \mathcal{G}^x.
 \end{aligned} \tag{4.40}$$

As a result, making use of relations (4.34)–(4.37) and inequality (4.40), and substituting all back into (4.33) whilst making note of $|\langle \nabla f, \nabla \mathbf{a} \rangle| \leq |\nabla f||\nabla \mathbf{a}|$ and the bound $0 \leq \psi \leq 1$ we obtain:

$$\begin{aligned}
 0 \geq & -\psi F_a^\mathcal{G}([c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2]t_1/R^2 + 1 + c_1k^L t_1) \\
 & - t_1\psi^2 F_a^\mathcal{G}(\mathcal{G}_w - e^{-f}\mathcal{G}) + t_1^2(\psi^2/m)[|\nabla f|^2 + e^{-f}\mathcal{G} + \mathbf{a} - \partial_t f]^2 \\
 & - 2c_1 t_1 \psi^{3/2} |\nabla f| F_a^\mathcal{G} / R \\
 & - 2t_1^2 \psi^2 |\nabla f|^2 [(m - 1)k + (\mu - 1)k^U + k^\nabla - (e^{-f}\mathcal{G} - \mathcal{G}_w + \mu e^f \mathcal{G}_{ww})/2] \\
 & - t_1^2 \psi^2 |\nabla f| (2|e^{-f}\mathcal{G}_x - \mu\mathcal{G}_{xw}| + \mu|\nabla \partial_t \phi| + 2\mu k^L |\nabla \phi| + 2(\mu - 1)|\nabla \mathbf{a}|) \\
 & + \mu t_1^2 \psi^2 (e^{-f}\Delta_\phi \mathcal{G}^x + \Delta_\phi \mathbf{a} - \mu n[(k^L + k^U)^2 + 2k^\nabla]).
 \end{aligned} \tag{4.41}$$

In order to derive the final bounds out of this inequality it is helpful to label by y, z and Y, Z the quantities defined by (4.42) as appearing on the right-hand side of (4.41) and identify the latter in terms of the γ -quantities introduced earlier:

$$\left\{ \begin{aligned}
 y &= \psi |\nabla f|^2, \\
 z &= \psi (\partial_t f - \mathbf{a} - e^{-f}\mathcal{G}), \\
 y - \mu z &= \psi F_a^\mathcal{G}/t_1 > 0, \\
 Y &= 2[(m - 1)k + (\mu - 1)k^U + k^\nabla] \\
 &\quad + \sup\{-e^{-f}\mathcal{G} + \mathcal{G}_w - \mu e^f \mathcal{G}_{ww}\}_+ : (t, x, w) \in Q_{2R,T} \} \\
 &= 2[(m - 1)k + (\mu - 1)k^U + k^\nabla] + \gamma_A^{\mathcal{G},\mu}(2R), \\
 Z &= \mu \ell_2 + 2\mu k^L \ell_1 + 2 \sup\{|e^{-f}\mathcal{G}_x - \mu\mathcal{G}_{xw}| : (t, x, w) \in Q_{2R,T} \} \\
 &\quad + 2(\mu - 1) \sup\{|\nabla \mathbf{a}| : (x, t) \in Q_{2R,T} \} \\
 &= \mu \ell_2 + 2\mu k^L \ell_1 + 2\gamma_B^{\mathcal{G},\mu}(2R) + 2\gamma_B^{aw,\mu}(2R).
 \end{aligned} \right. \tag{4.42}$$

By substituting these back into (4.41) it follows from basic considerations that,

$$\begin{aligned}
 0 \geq & -\psi F_a^\mathcal{G}([c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2]t_1/R^2 + 1 + c_1k^L t_1) \\
 & + (t_1^2/m)((y - z)^2 - (2mc_1/R)\sqrt{y}(y - \mu z) - mYy - mZ\sqrt{y}) \\
 & - t_1\psi F_a^\mathcal{G}[\mathcal{G}_w - e^{-f}\mathcal{G}]_+ + \mu t_1^2 \psi^2 [e^{-f}\Delta_\phi \mathcal{G}^x + \Delta_\phi \mathbf{a}]_- \\
 & - \mu t_1^2 \psi^2 [\mu n(k^L + k^U)^2 + 2\mu n k^\nabla].
 \end{aligned} \tag{4.43}$$

Moreover, by an application of the Cauchy–Schwartz and Young’s inequalities, it is seen that for any $\varepsilon \in (0, 1)$, we have

$$\begin{aligned} & (y - z)^2 - (2mc_1/R)\sqrt{y}(y - \mu z) - mYy - mZ\sqrt{y} \\ & \geq (y - \mu z)^2/\mu^2 - m^2c_1^2\mu^2(y - \mu z)/[2(\mu - 1)R^2] \\ & \quad - m^2\mu^2Y^2/[4(1 - \varepsilon)(\mu - 1)^2] - (3/4)(m^4\mu^2Z^4/[4\varepsilon(\mu - 1)^2])^{1/3}. \end{aligned} \tag{4.44}$$

Hence by taking advantage of (4.44) it follows from (4.43) that

$$\begin{aligned} 0 & \geq -\psi F_a^{\mathcal{G}} [(c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2)t_1/R^2 + 1 + c_1k^L t_1] \\ & \quad + (t_1^2/m)[(\psi F_a^{\mathcal{G}})^2/(t_1^2\mu^2) - m^2c_1^2\mu^2(\psi F_a^{\mathcal{G}})/(2(\mu - 1)R^2 t_1)] \\ & \quad - (mt_1^2\mu^2Y^2)/[4(1 - \varepsilon)(\mu - 1)^2] - t_1\psi F_a^{\mathcal{G}} [\mathcal{G}_w - e^{-f}\mathcal{G}]_+ \\ & \quad - [(3t_1^2)/(4m)](m^4\mu^2Z^4/[4\varepsilon(\mu - 1)^2])^{1/3} \\ & \quad + \mu t_1^2\psi^2 [e^{-f}\Delta_{\phi}\mathcal{G}^x + \Delta_{\phi}\mathbf{a}]_- - \mu t_1^2\psi^2 [\mu n(k^L + k^U)^2 + 2\mu nk^\nabla]. \end{aligned} \tag{4.45}$$

Subsequently and more shortly we can deduce and write the above inequality as a quadratic inequality

$$0 \geq (\psi F_a^{\mathcal{G}})^2/(m\mu^2) - (\psi F_a^{\mathcal{G}})\mathbf{X}_1 - t_1^2\mathbf{X}_2, \tag{4.46}$$

where we have set

$$\begin{aligned} \mathbf{X}_1 & = [c_2 + (m - 1)c_1(1 + R\sqrt{k}) + 2c_1^2]t_1/R^2 + 1 \\ & \quad + c_1k^L t_1 + mt_1c_1^2\mu^2/[2(\mu - 1)R^2] + t_1\gamma_C^{\mathcal{G}}(2R), \end{aligned} \tag{4.47}$$

and

$$\begin{aligned} \mathbf{X}_2 & = m\mu^2Y^2/[4(1 - \varepsilon)(\mu - 1)^2] \\ & \quad + (3/4)[m\mu^2Z^4/(4\varepsilon(\mu - 1)^2)]^{1/3} \\ & \quad + \mu^2n(k^L + k^U)^2 + 2\mu^2nk^\nabla + \mu(\gamma_D^{\mathcal{G}}(2R) + \gamma_D^{aw}(2R)), \end{aligned} \tag{4.48}$$

whilst making note of (2.7), (2.8) and (2.10) (with $2R$ replacing R). As a result it follows from (4.46) that

$$\begin{aligned} \psi F_a^{\mathcal{G}} & \leq (m\mu^2/2) \left(\mathbf{X}_1 + \sqrt{\mathbf{X}_1^2 + (4t_1^2\mathbf{X}_2)/(m\mu^2)} \right) \\ & \leq (m\mu^2/2) \left(2\mathbf{X}_1 + \sqrt{(4t_1^2\mathbf{X}_2)/(m\mu^2)} \right) = m\mu^2\mathbf{X}_1 + t_1\mu\sqrt{m\mathbf{X}_2}. \end{aligned} \tag{4.49}$$

Since $\psi \equiv 1$ for $r(x, \tau) \leq R$ and (x_1, t_1) is the point where $\psi F_a^{\mathcal{G}}$ attains its maximum on $\{r(x, t) \leq 2R, 0 \leq t \leq \tau\}$ we have from (4.49) the bound

$$F_a^{\mathcal{G}}(x, \tau) = [\psi F_a^{\mathcal{G}}](x, \tau) \leq [\psi F_a^{\mathcal{G}}](x_1, t_1) \leq m\mu^2\mathbf{X}_1 + t_1\mu\sqrt{m\mathbf{X}_2}. \tag{4.50}$$

Therefore, recalling (4.7), substituting for \mathbf{X}_1 and \mathbf{X}_2 from (4.47) and (4.48) respectively and making note of $t_1 \leq \tau$, we can write after dividing both sides by

$\mu\tau > 0$,

$$\begin{aligned}
 \mu^{-1}|\nabla f|^2 - \partial_t f + \mathbf{a}(x, t) + e^{-f}\mathcal{G} &\leq (m\mu/\tau)\mathsf{X}_1 + \sqrt{m}\mathsf{X}_2 \\
 &\leq (m\mu)[c_2 + (m-1)c_1(1 + R\sqrt{k}) + 2c_1^2]/R^2 \\
 &\quad + (m\mu/\tau) + (m\mu)(\gamma_{\mathbb{C}}^{\mathcal{G}}(2R) + c_1k^L \\
 &\quad + mc_1^2\mu^2/[2(\mu-1)R^2]) \\
 &\quad + \sqrt{m}\{m\mu^2\mathsf{Y}^2/[4(1-\varepsilon)(\mu-1)^2] \\
 &\quad + (3/4)[m\mu^2\mathsf{Z}^4/(4\varepsilon(\mu-1)^2)]^{1/3} \quad (4.51) \\
 &\quad + n\mu^2(k^L + k^U)^2 + 2n\mu^2k^{\nabla} \\
 &\quad + \mu(\gamma_{\mathbb{D}}^{\mathcal{G}}(2R) + \gamma_{\mathbb{D}}^{\mathbf{a}w}(2R))\}^{1/2}.
 \end{aligned}$$

Finally using the arbitrariness of $0 < \tau \leq T$ it follows after reverting back to w upon noting the relation $f = \log w$ and rearranging terms that

$$\begin{aligned}
 \frac{|\nabla w|^2}{\mu w^2} - \frac{\partial_t w}{w} + \frac{\mathcal{G}}{w} + \mathbf{a}(x, t) &\leq (m\mu)[1/t + \gamma_{\mathbb{C}}^{\mathcal{G}}(2R) + c_1k^L] \\
 &\quad + (m\mu)[mc_1^2\mu^2/[2(\mu-1)] + c_2 \\
 &\quad + (m-1)c_1(1 + R\sqrt{k}) + 2c_1^2]/R^2 \\
 &\quad + \sqrt{m}\{m\mu^2\mathsf{Y}^2/[4(1-\varepsilon)(\mu-1)^2] \\
 &\quad + (3/4)[m\mu^2\mathsf{Z}^4/(4\varepsilon(\mu-1)^2)]^{1/3} \\
 &\quad + n\mu^2(k^L + k^U)^2 + 2n\mu^2k^{\nabla} \\
 &\quad + \mu[\gamma_{\mathbb{D}}^{\mathcal{G}}(2R) + \gamma_{\mathbb{D}}^{\mathbf{a}w}(2R)]\}^{1/2}. \quad (4.52)
 \end{aligned}$$

A reference to (4.42) and substituting for Y , Z leads at once to the desired estimate as formulated in theorem 4.1. \square

The global version of the above estimate can now be obtained by imposing suitable global bounds and then passing to the limit $R \rightarrow \infty$ on the right-hand side.

THEOREM 4.5. *Let $(M, g, d\sigma)$ be a smooth metric measure space with $d\sigma = e^{-\phi}dv_g$ and assume that the metric-potential pair (g, ϕ) is time dependent and of class \mathcal{C}^2 . Assume $\mathcal{R}ic_{\phi}^m(g) \geq -(m-1)kg$ and (4.2)–(4.3) hold globally in $M \times [0, T]$. Let $w = w(x, t)$ be a positive solution to (1.1). Then for every $\mu > 1$, $\varepsilon \in (0, 1)$ and for all $x \in M$, $0 < t \leq T$ we have the global gradient estimate*

$$\frac{|\nabla w|^2}{\mu w^2} - \frac{\partial_t w}{w} + \frac{\mathcal{G}(t, x, w)}{w} + \mathbf{a}(x, t) \leq m\mu \left[\frac{1}{t} + c_1k^L + \gamma_{\mathbb{C}}^{\mathcal{G}} \right] + M_{\mathbf{a}}^{\mathcal{G}}, \quad (4.53)$$

where $\gamma_C^{\mathcal{G}}$ is as in (2.7) and

$$\begin{aligned}
 M_{\mathbf{a}}^{\mathcal{G}} &= \sqrt{m} \left\{ n\mu^2(k^L + k^U)^2 + 2n\mu^2k^\nabla + \mu\gamma_D^{\mathcal{G}} + \mu\gamma_D^{\mathbf{a}w} \right. \\
 &\quad + \frac{m\mu^2[(m-1)k + (\mu-1)k^U + k^\nabla + \gamma_A^{\mathcal{G},\mu}/2]^2}{(1-\varepsilon)(\mu-1)^2} \\
 &\quad \left. + \frac{3}{4} \left[\frac{m\mu^2[\mu\ell_2 + 2\mu k^L\ell_1 + 2\gamma_B^{\mathcal{G},\mu} + 2\gamma_B^{\mathbf{a}w,\mu}]^4}{4\varepsilon(\mu-1)^2} \right]^{1/3} \right\}^{1/2}. \tag{4.54}
 \end{aligned}$$

With the aid of the differential Harnack estimate established in theorem 4.1 we can now prove a parabolic Harnack inequality for positive solutions to the system (1.1).

THEOREM 4.6. *Under the assumptions of theorem 4.1 for all $(x_1, t_1), (x_2, t_2)$ in $Q_{R,T}$ with $t_2 > t_1$ and $\mu > 1$ we have*

$$w(x_2, t_2) \geq w(x_1, t_1) \left(\frac{t_2}{t_1} \right)^{-m\mu} \exp \left[\Sigma(t_2 - t_1) - \frac{\mu L(x_1, x_2)}{4(t_2 - t_1)} \right]. \tag{4.55}$$

Here $L(x_1, x_2) = \inf \| |\dot{\zeta}(t)|_{g(t)}^2 \|_{L^1(0,1;dt)}$ where the infimum is over all $\zeta \in \mathcal{C}^1([0, 1]; M)$ lying in $Q_{R,T}$ with $\zeta(0) = x_1, \zeta(1) = x_2$ and the constant Σ depends only on the bounds in theorem 4.1 [see (4.57)–(4.58)]. If the bounds as in theorem 4.5 are global then the estimate is global.

Proof. Here we shall focus only on the local Harnack inequality as the global one is very similar. The idea is to integrate estimate (4.4) along suitable space–time curves in $Q_{R,T} \subset M \times [0, T]$. To this end let us first rewrite the latter inequality (4.4) as

$$\frac{\partial_t w}{w} \geq \frac{|\nabla w|^2}{\mu w^2} - \frac{m\mu}{t} + \Sigma. \tag{4.56}$$

Here Σ is a constant containing all the terms (bounds) on the right-hand side of (4.4) except the first, and the terms $\underline{\mathbf{a}}, \underline{\mathcal{G}}$ resulting from the expression on the left, specifically,

$$\Sigma = \underline{\mathcal{G}} + \underline{\mathbf{a}} - (m\mu c_1 k^L + L + m\mu\gamma_C^{\mathcal{G}}(2R) + M_{\mathbf{a}}^{\mathcal{G}}), \tag{4.57}$$

$$\underline{\mathbf{a}} = \inf_{Q_{2R,T}} \mathbf{a}, \quad \underline{\mathcal{G}} = \inf_{\Theta_{2R,T}} \mathcal{G}(t, x, w)/w. \tag{4.58}$$

Suppose $\zeta \in \mathcal{C}^1([t_1, t_2]; M)$ is an arbitrary curve lying entirely in $Q_{R,T}$ with $\zeta(t_1) = x_1$ and $\zeta(t_2) = x_2$. Using the above inequality and writing $\dot{\zeta} = d\zeta/dt$ it

is seen that

$$\begin{aligned}
 d/dt[\log w(\zeta(t), t)] &= \langle \nabla w/w, \dot{\zeta}(t) \rangle + \partial_t w/w \\
 &\geq \langle \nabla w/w, \dot{\zeta}(t) \rangle + |\nabla w|^2/(\mu w^2) - (m\mu)/t + \Sigma \\
 &= \mu^{-1}|\nabla w/w + \mu\dot{\zeta}(t)/2|^2 - \mu|\dot{\zeta}(t)|^2/4 - (m\mu)/t + \Sigma \\
 &\geq -\mu|\dot{\zeta}(t)|^2/4 - (m\mu)/t + \Sigma,
 \end{aligned} \tag{4.59}$$

where the inner products are with respect to the metric $g(t)$. Integrating the above inequality thus gives

$$\begin{aligned}
 \log \frac{w(x_2, t_2)}{w(x_1, t_1)} &= \log w(\zeta(t), t) \Big|_{t_1}^{t_2} = \int_{t_1}^{t_2} \frac{d}{dt} \log w(\zeta(t), t) dt \\
 &\geq \int_{t_1}^{t_2} -\frac{\mu}{4} |\dot{\zeta}(t)|^2 dt - \int_{t_1}^{t_2} \frac{m\mu}{t} dt + \int_{t_1}^{t_2} \Sigma dt \\
 &= -m\mu \log(t_2/t_1) - (\mu/4) \int_{t_1}^{t_2} |\dot{\zeta}(t)|^2 dt + (t_2 - t_1)\Sigma.
 \end{aligned} \tag{4.60}$$

Reparametrizing the curve ζ and exponentiating both sides gives at once (4.55). \square

5. The case of time-independent metrics and potentials

In this section we discuss gradient estimates and the implications of what has been developed earlier in the so-called static case, that is, when the metric and potential are time independent ($\partial_t g \equiv 0$ and $\partial_t \phi \equiv 0$). The local and global differential Harnack estimates for positive smooth solutions to the equation

$$\mathcal{L}_\phi^a[w] = (\partial_t - a(x, t) - \Delta_\phi)w = \mathcal{G}(t, x, w), \tag{5.1}$$

can be formulated as below. Note that in this context $Q_{R,T} = \mathcal{B}_R \times [0, T]$ with $R, T > 0$ and the explicit bounds (4.2)–(4.3) are no longer needed (see the proof below).

THEOREM 5.1. *Let $(M, g, d\sigma)$ be a smooth metric measure space with $d\sigma = e^{-\phi} dv_g$ and let $\text{Ric}_\phi^m(g) \geq -(m-1)kg$ in \mathcal{B}_{2R} for some $m \geq n, k \geq 0$ and $R > 0$. If $w = w(x, t)$ is a positive solution to (5.1), then for every $\mu > 1, \varepsilon \in (0, 1)$ and for all (x, t) in $\mathcal{B}_R \times [0, T]$ with $t > 0$ we have the gradient estimate*

$$\frac{|\nabla w|^2}{\mu w^2} - \frac{\partial_t w}{w} + \frac{\mathcal{G}(t, x, w)}{w} + a(x, t) \leq \frac{m\mu}{t} + L + m\mu\gamma_C^\mathcal{G}(2R) + M_a^\mathcal{G}. \tag{5.2}$$

The quantities appearing on the right-hand side of the bound (5.2) are given respectively by

$$L = \frac{m\mu}{R^2} \left[\frac{mc_1^2\mu^2}{2(\mu-1)} + c_2 + (m-1)c_1(1 + R\sqrt{k}) + 2c_1^2 \right] \tag{5.3}$$

and

$$M_a^{\mathcal{G}} = \sqrt{m} \left\{ \mu[\gamma_D^{\mathcal{G}}(2R) + \gamma_D^{aw}(2R)] + \frac{m\mu^2[(m-1)k + \gamma_A^{\mathcal{G},\mu}(2R)/2]^2}{(1-\varepsilon)(\mu-1)^2} \right. \\ \left. + \frac{3}{4} \left[\frac{4m\mu^2[\gamma_B^{\mathcal{G},\mu}(2R) + \gamma_B^{aw,\mu}(2R)]^4}{\varepsilon(\mu-1)^2} \right]^{1/3} \right\}^{1/2}. \tag{5.4}$$

Proof. Referring to theorem 4.1 and bounds (4.2)–(4.3) we can set $k^L = 0, k^U = 0, k^\nabla = 0$ and $\ell_2 = 0$. Substituting these values in (4.5)–(4.6) leads to (5.3)–(5.4). Note that here in the formulation of $M_a^{\mathcal{G}}$ we have observed that in (4.42) in the expression for Y we have $(\mu-1)k^U + k^\nabla = 0$ and in the expression for Z we have $\mu\ell_2 + 2\mu k^L \ell_1 = 0$. The resulting cancellations lead to (5.4). \square

THEOREM 5.2. *Let $(M, g, d\sigma)$ be a smooth metric measure space with $d\sigma = e^{-\phi} dv_g$ and $\mathcal{R}ic_\phi^m(g) \geq -(m-1)kg$ in M for some $m \geq n$ and $k \geq 0$. If $w = w(x, t)$ is a positive solution to (5.1), then for every $\mu > 1, \varepsilon \in (0, 1)$ and for all (x, t) in $M \times [0, T]$ with $t > 0$ we have the global gradient estimate*

$$\frac{|\nabla w|^2}{\mu w^2} - \frac{\partial_t w}{w} + \frac{\mathcal{G}(t, x, w)}{w} + a(x, t) \leq \frac{m\mu}{t} + m\mu\gamma_C^{\mathcal{G}} + M_a^{\mathcal{G}}, \tag{5.5}$$

where

$$M_a^{\mathcal{G}} = \sqrt{m} \left\{ \mu(\gamma_D^{\mathcal{G}} + \gamma_D^{aw}) + \frac{m\mu^2[(m-1)k + \gamma_A^{\mathcal{G},\mu}/2]^2}{(1-\varepsilon)(\mu-1)^2} \right. \\ \left. + \frac{3}{4} \left[\frac{4m\mu^2[\gamma_B^{\mathcal{G},\mu} + \gamma_B^{aw,\mu}]^4}{\varepsilon(\mu-1)^2} \right]^{1/3} \right\}^{1/2} \tag{5.6}$$

and the γ -quantities used are as in (2.5)–(2.8).

Proof. Starting from (5.2) and the terms on the right, an inspection of (5.3) shows that $L \rightarrow 0$ as $R \rightarrow \infty$. A similar consideration for expression (5.6) shows that it suffices to replace the local constants $\gamma_A^{\mathcal{G},\mu}(2R), \gamma_B^{\mathcal{G},\mu}(2R), \gamma_C^{\mathcal{G}}(2R), \gamma_D^{\mathcal{G}}(2R)$ and $\gamma_D^{aw}(2R), \gamma_B^{aw,\mu}(2R)$ with their global counterparts. The conclusion now follows by passing to the limit $R \rightarrow \infty$ in (5.2) and taking the above into account. \square

In the static case and for the elliptic counterpart of (5.1) ($\partial_t g \equiv 0, \partial_t \phi \equiv 0, \partial_t w \equiv 0, \partial_t a \equiv 0$) we can deduce from the above global estimate by passing to the limit $t \rightarrow \infty$ the following global elliptic estimate

$$\frac{|\nabla w|^2}{\mu w^2} + \frac{\mathcal{G}(x, w)}{w} + a(x) \leq m\mu\gamma_C^{\mathcal{G}} + M_a^{\mathcal{G}}. \tag{5.7}$$

As a consequence of the global elliptic estimate (5.7) we can now prove the following Liouville type result for positive solutions to the equation $\Delta_\phi w + \mathcal{G}(w) = 0$ [i.e., $a \equiv 0$ and $\mathcal{G} = \mathcal{G}(w)$]. We later present some applications to specific nonlinearities.

THEOREM 5.3. *Let $(M, g, d\sigma)$ be a smooth metric measure space with $d\sigma = e^{-\phi} dv_g$ and $\mathcal{R}ic_\phi^m(g) \geq 0$ everywhere in M . Let w be a positive solution to $\Delta_\phi w + \mathcal{G}(w) = 0$. Then for every $\mu > 1, \varepsilon \in (0, 1)$ and all $x \in M$ we have the global gradient estimate*

$$\frac{|\nabla w|^2}{\mu w^2} + \frac{\mathcal{G}(w)}{w} \leq \frac{m\mu\gamma_A^{\mathcal{G},\mu}}{2(\mu - 1)\sqrt{1 - \varepsilon}} + m\mu\gamma_C^{\mathcal{G}}. \tag{5.8}$$

In particular, if along the solution w , we have the following

- $\mathcal{G}(w) \geq 0$,
- $\mathcal{G}(w) - w\mathcal{G}'(w) \geq 0$,
- $\mathcal{G}(w) - w\mathcal{G}'(w) + \mu w^2\mathcal{G}''(w) \geq 0$ for some $\mu > 1$,

everywhere on M , then w is constant and as a result $\mathcal{G}(w) = 0$.

Proof. Since $\mathcal{G} = \mathcal{G}(w)$ and $\mathbf{a} \equiv 0$ we obtain from (2.4), (2.8) the identities $\gamma_D^{\mathcal{G}} = 0, \gamma_D^{aw} = 0$ and from (2.2), (2.6) the identities $\gamma_B^{\mathcal{G},\mu} = 0, \gamma_B^{aw,\mu} = 0$. Hence substituting these together with $k = 0$ in (5.7) we arrive at once at (5.8). Next, from the prescribed inequalities satisfied by \mathcal{G} we have

$$\gamma_A^{\mathcal{G},\mu} = \sup_M A_{\mathcal{G}}^\mu(w) = \sup_M \left\{ \frac{1}{w} [-\mathcal{G}(w) + w\mathcal{G}'(w) - \mu w^2\mathcal{G}''(w)]_+ \right\} = 0 \tag{5.9}$$

and in a similar way

$$\gamma_C^{\mathcal{G}} = \sup_M C_{\mathcal{G}}(w) = \sup_M \left\{ \frac{1}{w} [w\mathcal{G}'(w) - \mathcal{G}(w)]_+ \right\} = 0. \tag{5.10}$$

Thus substituting back into (5.8) it follows that

$$\frac{|\nabla w|^2}{\mu w^2} \leq \frac{|\nabla w|^2}{\mu w^2} + \frac{\mathcal{G}(w)}{w} \equiv 0 \tag{5.11}$$

and so $|\nabla w| \equiv 0$. Therefore, w is a constant and so a further reference to the equation $\Delta_\phi w + \mathcal{G}(w) = 0$ gives $\mathcal{G}(w) = 0$. □

In order to illustrate the strength of the Liouville result in theorem 5.3 let us turn to considering some specific cases. As a first application consider the nonlinearity

$$\mathcal{G}(w) = \sum_{j=1}^d A_j w^{p_j}, \tag{5.12}$$

with real coefficients A_j and real exponents p_j ($1 \leq j \leq d$). Then by a direct calculation $\mathcal{G} - w\mathcal{G}' = \sum A_j(1 - p_j)w^{p_j}$ and $\mathcal{G} - w\mathcal{G}' + \mu w^2\mathcal{G}'' = \sum [A_j(p_j - 1)(\mu p_j - 1)]w^{p_j}$. If $A_j \geq 0$ we have $\mathcal{G}(w) \geq 0$ whilst if $p_j \leq 1$ we have $\mathcal{G} - w\mathcal{G}' \geq 0$ and so $\mathcal{G} - w\mathcal{G}' + \mu w^2\mathcal{G}'' \geq 0$ (by choosing $\mu > 1$ suitably). Hence theorem 5.3 now leads to the following conclusion extending earlier results on Yamabe type problems to more general nonlinearities.

THEOREM 5.4. *Let $(M, g, d\sigma)$ be a complete smooth metric measure space with $d\sigma = e^{-\phi} dv_g$ and $\mathcal{R}ic_\phi^m(g) \geq 0$. Let w be a positive smooth solution to the equation*

$$\Delta_\phi w + \sum_{j=1}^d A_j w^{p_j} = 0. \tag{5.13}$$

If $A_j \geq 0$ and $p_j \leq 1$ for $1 \leq j \leq d$ then w is a constant.

As another application relating to the discussions in § 1 consider a nonlinearity $\mathcal{G} = \mathcal{G}(w)$ in the form of a superposition of logarithmic and power-like nonlinearities, with real coefficients A, B , real exponents p, q , and a function $\Phi \in \mathcal{C}^2(\mathbb{R})$, specifically, in the form

$$\mathcal{G}(w) = Aw^p\Phi(\log w) + Bw^q. \tag{5.14}$$

A straightforward calculation pertaining to the quantities formulated in theorem 5.3 now leads to

$$\mathcal{G} - w\mathcal{G}' = Aw^p[(1 - p)\Phi - \Phi'] + B(1 - q)w^q, \tag{5.15}$$

and subsequently

$$\begin{aligned} \mathcal{G} - w\mathcal{G}' + \mu w^2\mathcal{G}' &= A(1 - p)(1 - \mu p)w^p\Phi + A[\mu(2p - 1) - 1]w^p\Phi' \\ &\quad + A\mu w^p\Phi' + B(q - 1)(\mu q - 1)w^q, \end{aligned} \tag{5.16}$$

where the argument $s = \log w$ of Φ and its derivatives have been abbreviated. Next let us formally fix an upper and a lower bound on the solutions w , say, $0 < \underline{w} \leq w \leq \bar{w}$ and put $\alpha = \log \underline{w}$, $\beta = \log \bar{w}$. Then evidently $s = \log w \in [\alpha, \beta] \subset (0, \infty)$. Furthermore, suppose $\Phi > 0$ and let c^L, c^U denote the infimum and supremum of Φ'/Φ and d the infimum of Φ''/Φ over $[\alpha, \beta]$ respectively. Then it is clear that we have the inequalities

$$c^L \leq \Phi'/\Phi \leq c^U, \quad \Phi''/\Phi \geq d. \tag{5.17}$$

Next, regarding the condition $\mathcal{G} - w\mathcal{G}' \geq 0$ in theorem 5.3, we have upon noting (5.15) the implications $\mathcal{G} - w\mathcal{G}' \geq 0 \iff (1 - p)\Phi \geq \Phi' \iff 1 - p \geq \Phi'/\Phi$ which then holds when

$$1 - p \geq c^U. \tag{5.18}$$

Likewise regarding the condition $\mathcal{G} - w\mathcal{G}' + \mu w^2\mathcal{G}' \geq 0$ in theorem 5.3 we have upon referring to (5.16) the relation $(1 - p)(1 - \mu p) + [\mu(2p - 1) - 1]\Phi'/\Phi + \mu\Phi''/\Phi \geq 0$ which by taking advantage of (5.17) holds when

$$(1 - p)(1 - \mu p) + [\mu(2p - 1) - 1]c^L + \mu d \geq 0 \text{ if } (2p - 1)\mu - 1 \geq 0, \tag{5.19}$$

$$(1 - p)(1 - \mu p) + [\mu(2p - 1) - 1]c^U + \mu d \geq 0 \text{ if } (2p - 1)\mu - 1 \leq 0. \tag{5.20}$$

The following theorem now directly results from theorem 5.3.

THEOREM 5.5. *Let $(M, g, d\sigma)$ be a complete smooth metric measure space with $d\sigma = e^{-\phi} dv_g$ and $\mathcal{R}ic_\phi^m(g) \geq 0$. Let w be a positive smooth solution to the equation*

$$\Delta_\phi w + Aw^p \Phi(\log w) + Bw^q = 0. \tag{5.21}$$

where $\Phi \in \mathcal{C}^2(\mathbb{R})$ is as described above. Assume $A, B \geq 0, q \leq 1$ and $p \in \mathbb{R}$ satisfies (5.18) and (5.19)–(5.20) for some $\mu > 1$ (with $1 < \mu < 1/q$ when $B \neq 0$ and $0 < q < 1$). Then w is a constant. Moreover $Aw^p \Phi(\log w) + Bw^q = 0$.

It is interesting to note that in the case $p = 1$ the above follows from $\Phi \geq 0, \Phi' \leq 0$ and $\mu\Phi'' + (\mu - 1)\Phi' \geq 0$ for some $\mu > 1$ (or $1 < \mu < 1/q$ if $B \neq 0$ and $0 < q < 1$).

6. Further global bounds under super Perelman–Ricci flow

In this section we establish a different type of gradient estimate for positive smooth solutions to system (1.1). In fact, here, we establish the estimate under a weaker form of the flow inequality (i.e., corresponding to the case $m = \infty$) by assuming that the metric and potential evolve under a k -super Perelman–Ricci flow:

$$\begin{cases} \frac{1}{2} \frac{\partial g}{\partial t}(x, t) + \mathcal{R}ic_\phi(g)(x, t) \geq -kg(x, t), & k \geq 0, \\ \mathcal{R}ic_\phi(g)(x, t) = \mathcal{R}ic(g)(x, t) + \nabla_{g(t)} \nabla_{g(t)} \phi(x, t). \end{cases} \tag{6.1}$$

To this end suppose that $w = w(x, t)$ is a positive smooth solution to the equation $\mathcal{L}_\phi^a(w) = [\partial_t - a(x, t) - \Delta_\phi]w = \mathcal{G}(t, x, w)$. For $p \geq 2, q \in \mathbb{R}$ a pair of exponents and $\zeta = \zeta(t)$ a non-negative, smooth but otherwise arbitrary function set

$$\mathbf{X}_\zeta^{p,q}[w] = \zeta(t) \frac{|\nabla w|^p}{w^q} + \Gamma(w), \tag{6.2}$$

where $\Gamma = \Gamma(w)$ with $w > 0$ is some given function of class \mathcal{C}^2 .

LEMMA 6.1. *For $p \geq 2, q \in \mathbb{R}, \zeta = \zeta(t)$ as above and \mathcal{L}_ϕ^a as in (1.3), $\mathbf{X}_\zeta^{p,q}[w]$ satisfies the evolution identity*

$$\begin{aligned} \mathcal{L}_\phi^a(\mathbf{X}_\zeta^{p,q}[w]) &= \zeta' \frac{|\nabla w|^p}{w^q} - p\zeta \frac{|\nabla w|^{p-2}}{w^q} \left[\frac{1}{2} \partial_t g + \mathcal{R}ic_\phi \right] (\nabla w, \nabla w) \\ &\quad + p\zeta \frac{|\nabla w|^{p-2}}{w^q} \left[a|\nabla w|^2 + w \langle \nabla w, \nabla a \rangle + \mathcal{G}_w |\nabla w|^2 + \langle \nabla w, \mathcal{G}_x \rangle \right] \\ &\quad - q\zeta [a w + \mathcal{G}] \frac{|\nabla w|^p}{w^{q+1}} - \zeta a \frac{|\nabla w|^p}{w^q} - \frac{p}{2} \frac{\zeta}{w^q} \langle \nabla |\nabla w|^{p-2}, \nabla |\nabla w|^2 \rangle \\ &\quad - p\zeta \frac{|\nabla w|^{p-2}}{w^q} \left| \nabla \nabla w - \frac{q}{w} [\nabla w \otimes \nabla w] \right|^2 - q[q + 1 - pq] \zeta \frac{|\nabla w|^{p+2}}{w^{q+2}} \\ &\quad - a\Gamma(w) + (a w + \mathcal{G})\Gamma'(w) - \Gamma''(w) |\nabla w|^2. \end{aligned} \tag{6.3}$$

In particular if the metric-potential pair evolves under the super Perelman–Ricci flow inequality (6.1) then

$$\begin{aligned} \mathcal{L}_\phi^a(\mathbf{X}_\zeta^{p,q}[w]) &\leq \frac{|\nabla w|^p}{w^q} \left[\zeta' + p\zeta(\mathbf{k} + \mathbf{a} + \mathcal{G}_w) - (q + 1)\zeta\mathbf{a} - q\zeta \frac{\mathcal{G}}{w} \right] \\ &\quad + p\zeta \frac{|\nabla w|^{p-1}}{w^q} (|\nabla \mathbf{a}|w + |\mathcal{G}_x|) - q[q + 1 - pq]\zeta \frac{|\nabla w|^{p+2}}{w^{q+2}} \\ &\quad - \mathbf{a}\Gamma(w) + (\mathbf{a}w + \mathcal{G})\Gamma'(w) - \Gamma''(w)|\nabla w|^2. \end{aligned} \tag{6.4}$$

Proof. The application of the operator $\mathcal{L}_\phi^a = \partial_t - \mathbf{a}(x, t) - \Delta_\phi$ on (6.2) is easily seen to split into the sum

$$\mathcal{L}_\phi^a(\mathbf{X}_\zeta^{p,q}[w]) = \zeta'|\nabla w|^p/w^q + \zeta\mathcal{L}_\phi^a(|\nabla w|^p/w^q) + \mathcal{L}_\phi^a(\Gamma(w)). \tag{6.5}$$

Starting from the last term on the right-hand side it is not difficult to see that

$$\mathcal{L}_\phi^a(\Gamma(w)) = \Gamma'(w)\mathcal{L}_\phi^a(w) - \mathbf{a}[\Gamma(w) - w\Gamma'(w)] - \Gamma''(w)|\nabla w|^2. \tag{6.6}$$

Moving onto the second term which is the more involved one we proceed by calculating the first order space–time derivatives of $|\nabla w|^p/w^q$ as

$$\left[\frac{\partial_t}{\nabla} \right] \frac{|\nabla w|^p}{w^q} = p \frac{|\nabla w|^{p-2}}{2w^q} \left[\frac{\partial_t}{\nabla} \right] |\nabla w|^2 - q \frac{|\nabla w|^p}{w^{q+1}} \left[\frac{\partial_t}{\nabla} \right] w. \tag{6.7}$$

Now an application of the divergence operator to the second line gives

$$\begin{aligned} \Delta \frac{|\nabla w|^p}{w^q} &= \frac{p}{2} \frac{|\nabla w|^{p-2}}{w^q} \Delta |\nabla w|^2 + \frac{p}{2w^q} \langle \nabla |\nabla w|^{p-2}, \nabla |\nabla w|^2 \rangle \\ &\quad - pq \frac{|\nabla w|^{p-2}}{w^{q+1}} \langle \nabla |\nabla w|^2, \nabla w \rangle - q \frac{|\nabla w|^p}{w^{q+1}} \Delta w + q(q + 1) \frac{|\nabla w|^{p+2}}{w^{q+2}}, \end{aligned} \tag{6.8}$$

and thus after using the relation $\Delta_\phi(|\nabla w|^p/w^q) = \Delta(|\nabla w|^p/w^q) - \langle \nabla \phi, |\nabla w|^p/w^q \rangle$ it follows that

$$\begin{aligned} \mathcal{L}_\phi^a \left(\frac{|\nabla w|^p}{w^q} \right) &= \frac{p}{2} \frac{|\nabla w|^{p-2}}{w^q} \mathcal{L}_\phi^a(|\nabla w|^2) - \frac{p}{2w^q} \langle \nabla |\nabla w|^{p-2}, \nabla |\nabla w|^2 \rangle \\ &\quad + pq \frac{|\nabla w|^{p-2}}{w^{q+1}} \langle \nabla |\nabla w|^2, \nabla w \rangle - q \frac{|\nabla w|^p}{w^{q+1}} \mathcal{L}_\phi^a(w) \\ &\quad + \frac{1}{2}(p + 2q - 2) \frac{|\nabla w|^p}{w^q} - q(q + 1) \frac{|\nabla w|^{p+2}}{w^{q+2}}. \end{aligned} \tag{6.9}$$

Next by making note of the weighted Bochner–Weitzenböck formula in lemma 3.1 and the differentiation identity in lemma 3.2 it is seen that

$$\left[\frac{\partial_t}{\Delta_\phi} \right] |\nabla w|^2 = 2 \left[\langle \nabla w, \nabla \partial_t w \rangle - (1/2)[\partial_t g](\nabla w, \nabla w) \right] + \left[|\nabla \nabla w|^2 + \langle \nabla w, \nabla \Delta_\phi w \rangle + \mathcal{R}ic_\phi(g) \right]. \tag{6.10}$$

Therefore, $(\partial_t - \Delta_\phi)|\nabla w|^2 = -\partial_t g + 2\langle \nabla w, \nabla \partial_t w \rangle - 2[\mathcal{R}ic_\phi(g) + |\nabla \nabla w|^2 + \langle \nabla w, \nabla \Delta_\phi w \rangle]$ and so together with the equation $\partial_t w = \Delta_\phi w + \mathbf{a}(x, t)w +$

$\mathcal{G}(t, x, w)$ this gives

$$\begin{aligned} \mathcal{L}_\phi^a |\nabla w|^2 &= (\partial_t - \mathbf{a} - \Delta_\phi) |\nabla w|^2 = -[\partial_t g + 2\mathcal{R}ic_\phi(g)] + \mathbf{a} |\nabla w|^2 \\ &\quad + 2w \langle \nabla w, \nabla \mathbf{a} \rangle + 2 \langle \nabla w, \nabla \mathcal{G} \rangle - 2 |\nabla \nabla w|^2. \end{aligned} \tag{6.11}$$

Therefore, upon substitution back in (6.9) it follows that

$$\begin{aligned} \mathcal{L}_\phi^a \left(\frac{|\nabla w|^p}{w^q} \right) &= -\frac{p}{2} \frac{|\nabla w|^{p-2}}{w^q} [\partial_t g + 2\mathcal{R}ic_\phi] + p \frac{|\nabla w|^{p-2}}{w^q} \langle \nabla w, \nabla \mathcal{G} \rangle \\ &\quad + \frac{|\nabla w|^{p-2}}{w^q} [(p-1) |\nabla w|^2 \mathbf{a} + pw \langle \nabla w, \nabla \mathbf{a} \rangle] \\ &\quad - p \frac{|\nabla w|^{p-2}}{w^q} \left(|\nabla \nabla w|^2 - \frac{q}{w} \langle \nabla |\nabla w|^2, \nabla w \rangle + \frac{q}{p} (q+1) \frac{|\nabla w|^4}{w^2} \right) \\ &\quad - \frac{p}{2w^q} \langle \nabla |\nabla w|^{p-2}, \nabla |\nabla w|^2 \rangle - q \frac{|\nabla w|^p}{w^{q+1}} [\mathbf{a}w + \mathcal{G}]. \end{aligned} \tag{6.12}$$

Now for the term inside the brackets on the third line of the right note that

$$\begin{aligned} |\nabla \nabla w|^2 - \frac{q}{w} \langle \nabla |\nabla w|^2, \nabla w \rangle + \frac{q}{p} (q+1) \frac{|\nabla w|^4}{w^2} \\ = \left| \nabla \nabla w - \frac{q}{w} [\nabla w \otimes \nabla w] \right|^2 + \left[\frac{q}{p} (q+1) - q^2 \right] \frac{|\nabla w|^4}{w^2}, \end{aligned} \tag{6.13}$$

and therefore substituting back results in the formulation

$$\begin{aligned} \mathcal{L}_\phi^a \left(\frac{|\nabla w|^p}{w^q} \right) &= -\frac{p}{2} \frac{|\nabla w|^{p-2}}{w^q} [\partial_t g + 2\mathcal{R}ic_\phi] + p \frac{|\nabla w|^{p-2}}{w^q} \langle \nabla w, \nabla \mathcal{G} \rangle \\ &\quad + \frac{|\nabla w|^{p-2}}{w^q} [(p-1) |\nabla w|^2 \mathbf{a} + pw \langle \nabla w, \nabla \mathbf{a} \rangle] \\ &\quad - \frac{p}{2w^q} \langle \nabla |\nabla w|^{p-2}, \nabla |\nabla w|^2 \rangle - q \frac{|\nabla w|^p}{w^{q+1}} [\mathbf{a}w + \mathcal{G}] \\ &\quad - p \frac{|\nabla w|^{p-2}}{w^q} \left| \nabla \nabla w - \frac{q}{w} [\nabla w \otimes \nabla w] \right|^2 - q[q+1-pq] \frac{|\nabla w|^{p+2}}{w^{q+2}}. \end{aligned}$$

Now returning to (6.5) and making note of the above fragments and the basic identity $\nabla \mathcal{G}(t, x, w) = \mathcal{G}_x(t, x, w) + \mathcal{G}_w(t, x, w) \nabla w$ it follows that

$$\begin{aligned} \mathcal{L}_\phi^a (\mathbf{X}_\zeta^{p,q}[w]) &= \zeta' \frac{|\nabla w|^p}{w^q} + \zeta \mathcal{L}_\phi^a \left(\frac{|\nabla w|^p}{w^q} \right) + \mathcal{L}_\phi^a (\Gamma(w)) \\ &= \zeta' \frac{|\nabla w|^p}{w^q} - p\zeta \frac{|\nabla w|^{p-2}}{w^q} \left[\frac{1}{2} \partial_t g + \mathcal{R}ic_\phi \right] - q\zeta \frac{|\nabla w|^p}{w^{q+1}} [\mathbf{a}w + \mathcal{G}] \end{aligned}$$

$$\begin{aligned}
 & + \zeta \frac{|\nabla w|^{p-2}}{w^q} \left\{ p|\nabla w|^2 \mathcal{G}_w + p\langle \nabla w, \mathcal{G}_x \rangle + p|\nabla w|^2 \mathbf{a} + pw\langle \nabla w, \nabla \mathbf{a} \rangle \right\} \\
 & - \frac{p\zeta}{2w^q} \langle \nabla|\nabla w|^{p-2}, \nabla|\nabla w|^2 \rangle - q[q+1-pq]\zeta \frac{|\nabla w|^{p+2}}{w^{q+2}} \\
 & - p\zeta \frac{|\nabla w|^{p-2}}{w^q} \left| \nabla \nabla w - \frac{q}{w} [\nabla w \otimes \nabla w] \right|^2 - \left[\zeta \frac{|\nabla w|^p}{w^q} + \Gamma(w) \right] \mathbf{a} \\
 & + \Gamma'(w)[\mathbf{a}w + \mathcal{G}] - \Gamma''(w)|\nabla w|^2
 \end{aligned} \tag{6.14}$$

which is the desired conclusion. The final assertion follows by using the flow inequality $\partial_t g + 2\text{Ric}_\phi(g) \geq -2kg$ and $\langle \nabla|\nabla w|^{p-2}, \nabla|\nabla w|^2 \rangle \geq 0$ when $p \geq 2$. The last inequality follows by writing $W = |\nabla w|^2$ and setting $\alpha = (p - 2)/2$ and noting that the expression is $\langle \nabla W^\alpha, \nabla W \rangle = \alpha W^{\alpha-1} |\nabla W|^2 \geq 0$. \square

We now give two consequences of this evolution identity to global gradient estimates. Here w is taken a positive smooth solution to $\mathcal{L}_\phi[w] = (\partial_t - \Delta_\phi)w = \mathcal{G}(t, x, w)$ [i.e., with $\mathbf{a}(x, t) = 0$] whilst the pair (g, ϕ) is assumed to evolve under a k -super Perelman–Ricci flow. Furthermore, as we will be applying the maximum principle for the sake of convenience M is taken to be closed.

COROLLARY 6.2. *Under the assumptions of lemma 6.1, if $q[q + 1 - pq] \geq 0$, $\Gamma''(w) \geq 0$, $\Gamma'(w)\mathcal{G}(w) \leq 0$ and $q\mathcal{G}(w)/(pw) - \mathcal{G}'(w) \geq a$ along w then for $x \in M$ and $0 < t \leq T$*

$$\frac{|\nabla w|^p}{w^q}(x, t) \leq e^{p(k-a)t} \left\{ \max_M \left[\frac{|\nabla w|^p}{w^q} + \Gamma(w) \right](x, 0) - \Gamma(w)(x, t) \right\}. \tag{6.15}$$

Proof. Utilizing (6.4), $\Gamma''(w) \geq 0$, $\Gamma'(w)\mathcal{G}(w) \leq 0$ and $p\mathcal{G}'(w) - q\mathcal{G}(w)/w \leq -ap$ for some a we can write²

$$\begin{aligned}
 \mathcal{L}_\phi(\mathbf{X}_\zeta^{p,q}[w]) & \leq \frac{|\nabla w|^p}{w^q} \left[\zeta' + p\zeta k + p\zeta \mathcal{G}'(w) - q\zeta \frac{\mathcal{G}(w)}{w} \right] \\
 & \quad - q[q+1-pq]\zeta \frac{|\nabla w|^{p+2}}{w^{q+2}} + \Gamma'(w)\mathcal{G}(w) - \Gamma''(w)|\nabla w|^2 \\
 & \leq \frac{|\nabla w|^p}{w^q} [\zeta' + p(k-a)]\zeta - q[q+1-pq]\zeta \frac{|\nabla w|^{p+2}}{w^{q+2}}.
 \end{aligned} \tag{6.16}$$

The function $\zeta(t) = X e^{-p(k-a)t}$ is non-negative, smooth and satisfies $\zeta' + p(k-a)\zeta = 0$ (for $t \geq 0$ and fixed $X > 0$). Thus by substituting in (6.16) and noting $q(q + 1 - pq) \geq 0$ we have $\mathcal{L}_\phi(\mathbf{X}_\zeta^{p,q}[w]) \leq 0$. The assertion is now a consequence of the weak maximum principle giving

$$\mathbf{X}_\zeta^{p,q}[w](x, t) \leq \max_M \mathbf{X}_\zeta^{p,q}[w]|_{t=0}. \tag{6.17}$$

The proof is thus complete. \square

²Note that since $p \geq 2$ the condition $q[q + 1 - pq] \geq 0$ is equivalent to $0 \leq q \leq 1/(p - 1)$.

COROLLARY 6.3. Under the assumptions of lemma 6.1 if $\mathcal{G}, \mathcal{G}' \leq 0$ along w and (4.1) holds with $k \geq 0$ then for $x \in M$ and $0 < t \leq T$

$$|\nabla w|^2(x, t) \leq \frac{1 + 2kt}{2t} \left[\max_M w^2(x, 0) - w^2(x, t) \right]. \tag{6.18}$$

Proof. Setting $p = 2, q = 0$ and $\Gamma(w) = w^2/2$ in (6.2) gives $\mathbf{X} = \mathbf{X}_\zeta^{2,0}[w] = \zeta|\nabla w|^2 + w^2/2$ and so from (6.4) it follows that

$$\begin{aligned} \mathcal{L}_\phi(\mathbf{X}[w]) &= \mathcal{L}_\phi(\zeta|\nabla w|^2 + w^2/2) \\ &\leq [\zeta' + 2k\zeta + 2\zeta\mathcal{G}'(w)]|\nabla w|^2 + w\mathcal{G}(w) - |\nabla w|^2 \\ &\leq (\zeta' + 2k\zeta - 1)|\nabla w|^2 + 2\zeta\mathcal{G}'(w)|\nabla w|^2 + w\mathcal{G}(w). \end{aligned} \tag{6.19}$$

Now when $k \geq 0$ by taking $\zeta(t) = t/(1 + 2kt)$ we have $(\zeta' + 2k\zeta - 1) \leq 0$ and so subject to $\mathcal{G}, \mathcal{G}' \leq 0$ we have $\mathcal{L}_\phi(\mathbf{X}[w]) \leq 0$. The conclusion now follows by an application of the weak maximum principle. \square

Acknowledgements

The authors gratefully acknowledge support from the Engineering and Physical Sciences Research Council (EPSRC) through the grant EP/V027115/1. They also wish to thank the anonymous reviewer for a careful reading of the paper. All data is provided in full in the results section. Additional data is in the public domain at locations cited in the reference section.

References

- 1 E. Acerbi and R. Mingione. Gradient estimates for a class of parabolic systems. *Duke Math. J.* **136** (2007), 285–320.
- 2 S. M. Allen and J. W. Cahn. A microscopic theory for antiphase boundary motion and its applications to antiphase domain coarsening. *Acta Metall.* **27** (1979), 1085–1095.
- 3 T. Aubin. *Nonlinear analysis on manifolds* (Springer, New York, 1982).
- 4 M. Baileşteanu, X. Cao and A. Pulemotov. Gradient estimates for the heat equation under the Ricci flow. *J. Funct. Anal.* **258** (2010), 3517–3542.
- 5 M. Bateanu. A Harnack inequality for the parabolic Allen–Cahn equation. *Ann. Glob. Anal. Geom.* **51** (2017), 367–378.
- 6 D. Bakry, L’hypercontractivité et son utilisation en théorie des semigroupes, in: *Lecture Notes in Math.*, Vol. 1581 (Springer-Verlag, Berlin/New York, 1994), pp. 1–114.
- 7 D. Bakry and M. Emery, Diffusions hypercontractives In: Azma J., Yor M. (eds) *Séminaire de Probabilités XIX 1983/84. Lecture Notes in Mathematics*, Vol. 1123 (Springer, Berlin, Heidelberg).
- 8 D. Bakry, I. Gentil and M. Ledoux, *Analysis and Geometry of Markov Diffusion Operators*, A Series of Comprehensive Studies in Mathematics, Vol. 348 (Springer, New York, 2012).
- 9 M. F. Biduat-Véron and L. Veròn. Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of the Emden equations. *Invent. Math.* **106** (1991), 489–539.
- 10 L. A. Cafarelli, B. Gidas and J. Spruck. Asymptotic symmetry and local behavior of semi-linear elliptic equations with critical Sobolev growth. *Comm. Pure Appl. Math.* **42** (1989), 271–297.
- 11 E. Calabi. An extension of Hopf maximum principle with application to Riemannian geometry. *Duke Math. J.* **25** (1958), 45–56.
- 12 H. D. Cao, B. Chow, S. C. Chu and S. T. Yau. *Collected Papers on Ricci Flow, Surveys in Geometry and Topology*, Vol. 37 (International Press, Boston, 2003).

- 13 H. D. Cao, Recent progress on Ricci solitons, In: Recent Advances in Geometric Analysis, Advanced Lectures in Mathematics (ALM), Vol. 11 (International Press, Boston, 2010), pp. 1–38.
- 14 X. D. Cao, B. W. Liu, I. Pendleton and A. Ward. Differential Harnack estimates for Fisher’s equation. *Pac. J. Math.* **290** (2017), 273–300.
- 15 Y. Choquet-Bruhat, General Relativity and the Einstein Equations, Oxford Mathematical Monographs (OUP, Oxford, 2009).
- 16 B. Chow, P. Lu and L. Ni, Hamilton’s Ricci Flow, Graduate Studies in Mathematics Vol. 77 (AMS, USA, 2006).
- 17 N. T. Dung, N. N. Khanh and Q. A. Ngô. Gradient estimates for f -heat equations driven by Lichnerowicz’s equation on complete smooth metric measure spaces. *Manuscripta Math.* **155** (2018), 471–501.
- 18 R. A. Fisher. The wave of advance of advantageous genes. *Ann Eugen.* **7** (1937), 355–369.
- 19 M. Ghergu, S. Kim and H. Shahgholian. Exact behaviour around isolated singularity for semilinear elliptic equations with a log-type nonlinearity. *Adv. Nonlinear Anal.* **8** (2019), 995–1003.
- 20 B. Gidas and J. Spruck. Global and local behaviour of positive solutions of nonlinear elliptic equations. *Comm. Pure Appl. Math.* **34** (1981), 525–598.
- 21 A. Grigor’yan, Heat kernel analysis on manifolds, Studies in Advanced Mathematics (AMS, USA, 2013).
- 22 L. Gross. Logarithmic Sobolev inequalities. *Am. J. Math.* **97** (1976), 1061–1083.
- 23 R. Hamilton. A matrix Harnack estimate for heat equation. *Comm. Anal. Geom.* **1** (1993), 113–126.
- 24 R. Hamilton. The formation of singularities in the Ricci flow. *Surv. Diff. Geom.* **2** (1995), 7–136.
- 25 A. N. Kolmogorov, I. G. Petrovskii and N. S. Piskunov. Etude de l’équation de la diffusion avec croissance de la quantité de matière et son application a un problème biologique: Bull. Univ. d’Etatá. *Moscou, Ser. Int. A* **1** (1937), 1–26.
- 26 J. M. Lee and T. H. Parker. The Yamabe problem. *Bull. Am. Math. Soc.* **17** (1987), 37–91.
- 27 J. Li. Gradient estimates and Harnack inequalities for nonlinear parabolic and nonlinear elliptic equations on Riemannian manifolds. *J. Funct. Anal.* **100** (1991), 233–256.
- 28 P. Li, Geometric Analysis, Cambridge Studies in Advanced Mathematics, Vol. 134 (CUP, Cambridge, 2012).
- 29 P. Li and S. T. Yau. On the parabolic kernel of Schrödinger operator. *Acta Math.* **156** (1986), 153–201.
- 30 S. Li and X. D. Li. Harnack inequalities for Witten Laplacian on Riemannian manifolds with super Ricci flows. *Asian J. Math.* **22** (2018), 577–598.
- 31 X. D. Li. Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds. *J. Math. Pures Appl.* **84** (2005), 1295–1361.
- 32 J. Lott. Some geometric properties of the Bakry–Émery Ricci tensor. *Comment. Math. Helv.* **78** (2003), 865–883.
- 33 L. Ma. Gradient estimates for a simple elliptic equation on complete noncompact Riemannian manifolds. *J. Funct. Anal.* **241** (2006), 374–382.
- 34 P. Mastroli, M. Rigoli and A. G. Setti. *Yamabe Type Equations on Complete Non-compact Manifolds* (Springer, Basel, 2012).
- 35 R. Müller, Differential Harnack Inequalities and the Ricci Flow, EMS Series of Lectures in Mathematics (EMS, Zurich, 2006).
- 36 G. Perelman, The entropy formula for the Ricci Flow and its geometric application [arXiv: math.DG/0211159v1](https://arxiv.org/abs/math/0211159v1) (2002).
- 37 P. Souplet and Q. S. Zhang. Sharp gradient estimate and Yau’s Liouville theorem for the heat equation on noncompact manifolds. *Bull. Lond. Math. Soc.* **38** (2006), 1045–1053.
- 38 K. T. Sturm. Super-Ricci flows for metric measure spaces. *J. Funct. Anal.* **275** (2018), 3504–3569.
- 39 J. Sun. Gradient estimates for positive solutions of the heat equation under geometric flow. *Pac. J. Math.* **253** (2011), 489–510.

- 40 A. Taheri, *Function Spaces and Partial Differential Equations*, Vol. I, Oxford Lecture Series in Mathematics and its Applications, Vol. 40 (OUP, Oxford, 2015).
- 41 A. Taheri, *Function Spaces and Partial Differential Equations*, Vol. II, Oxford Lecture Series in Mathematics and its Applications, Vol. 41 (OUP, Oxford, 2015).
- 42 A. Taheri. Liouville theorems and elliptic gradient estimates for a nonlinear parabolic equation involving the Witten Laplacian. *Adv. Calc. Var.* **16** (2023), 425–441.
- 43 A. Taheri. Gradient estimates for a weighted Γ -nonlinear parabolic equation coupled with a super Perelman–Ricci flow and implications. *Potential Anal.* **59** (2023), 311–335.
- 44 A. Taheri and V. Vahidifar. On multiple solutions to a family of nonlinear elliptic systems in divergence form combined with an incompressibility constraint. *Nonlinear Anal.* **221** (2022), 112889.
- 45 A. Taheri and V. Vahidifar. Gradient estimates for nonlinear elliptic equations involving the Witten Laplacian on smooth metric measure spaces. *Adv. Nonlinear Anal.* **12** (2023), 20220288.
- 46 A. Taheri and V. Vahidifar. Souplet–Zhang and Hamilton type gradient estimates for nonlinear elliptic equations on smooth metric measure spaces. *Mathematika* **69** (2023), 751–779.
- 47 A. Taheri and V. Vahidifar. Gradient estimates for a nonlinear parabolic equation on smooth metric measure spaces with evolving metrics and potentials. *Nonlinear Anal.* **232** (2023), 113255.
- 48 A. Taheri and V. Vahidifar, Curvature conditions, Liouville-type theorems and Harnack inequalities for a nonlinear parabolic equation on smooth metric measure spaces, To appear in *Adv. Nonlinear Studies*, De Gruyter, 2024.
- 49 C. Villani, *Optimal transport: Old and New*, A Series of Comprehensive Studies in Mathematics, Vol. 338 (Springer, New York, 2008).
- 50 W. Wang. Harnack inequality, heat kernel bounds and eigenvalue estimate under integral Ricci curvature bounds. *J. Differ. Eqn.* **269** (2020), 1243–1277.
- 51 G. Wei and W. Wylie. Comparison geometry for the Bakry–Émery Ricci tensor. *J. Diff. Geom.* **83** (2009), 377–405.
- 52 J. Y. Wu. Li-Yau type estimates for a nonlinear parabolic equation on complete manifolds. *J. Math. Anal. Appl.* **369** (2010), 400–407.
- 53 J. Y. Wu. Elliptic gradient estimates for a weighted heat equation and applications. *Math. Z.* **280** (2015), 451–468.
- 54 J. Y. Wu. Gradient estimates for a nonlinear parabolic equation and Liouville theorems. *Manuscript Math.* **159** (2019), 511–547.
- 55 Q. S. Zhang, *Sobolev inequalities, heat kernels under Ricci flow and the Poincaré conjecture* (CRC Press, 2011).