

PAH anions as carriers of the mid-IR emission bands in planetary nebulae

Ryszard Szczerba¹, Christine Joblin², Olivier Berné², and Cezary Szyszka³

¹N. Copernicus Astronomical Center,
Rabiańska 8, 87-100 Toruń, Poland
email: szczerba@ncac.torun.pl

²Centre d'Etude Spatiale des Rayonnements, Université Toulouse 3 et CNRS,
9 Av. du Colonel Roche, 31028 Toulouse cedex 04, France

³Nicolaus Copernicus University,
Gagarina 11, 87-100 Toruń, Poland

Abstract. We present results of the mid-IR spectra decomposition for planetary nebulae and compact H II regions in our Galaxy and Magellanic Clouds. The striking correlation between the required PAH component with “7.7” μm band shifted to about 7.8 μm and electron densities of the modeled sources allows us to argue that this PAH component may be in fact PAH anions (PAH⁻).

Keywords. (ISM:) dust, extinction, (ISM:) planetary nebulae: general, infrared: ISM

1. Introduction

Among the different dust populations, the carriers of the so-called aromatic infrared bands (AIBs) are the subject of a long debate. The main AIBs are located at 3.3, 6.2, 7.7, 8.6, 11.3, 12.7 μm and are attributed to very small aromatic dust particles amongst which there are large polycyclic aromatic hydrocarbons (PAHs). However, these band positions are varying. In particular, the dominant band in the “7.7” μm complex arises at 7.8 μm in planetary nebulae (PNe) compared to 7.6 μm in compact-H II (c-H II) regions (Peeters *et al.* 2002) or in photo dissociation regions (PDRs). These shifts in the AIB positions may be related to chemical or physical changes of the carriers in different environments. To investigate possible evolution of the AIB carriers, Joblin *et al.* (2008) have analyzed spectra of several PNe and c-H II regions in the Milky Way (MW) and in the Large and Small Magellanic Clouds (LMC & SMC). Their analysis is based on the results of Berné *et al.* (2007) and Rapacioli *et al.* (2005) and uses a decomposition of the mid-IR spectra into a set of template spectra (see Sect.2).

In this paper we discuss results of the above analysis for selected PNe and c-H II regions in MW and LMC and show that the obtained decomposition may be understood by the presence of different amount of PAH anions in environments with different ionization parameter (U_{ion}).

2. Templates and the spectra decomposition

Rapacioli *et al.* (2005) and Berné *et al.* (2007) were able to extract the spectra of PAH neutrals (PAH⁰), PAH cations (PAH⁺) and very small grains (VSGs) from the mid-IR spectra of the analyzed PDRs. Joblin *et al.* (2008) have built three template spectra

Table 1. Relative amount of PAHs in [%] ($\text{PAH}^0 + \text{PAH}^+ + \text{PAH}^x = 100\%$) required for the fits presented in Figure 1.

Object	PAH^0 [%]	PAH^+ [%]	PAH^x [%]	N_{el} [cm^{-3}]	Ref
Vo 1	4	0	96	10^5	1:
SMP LMC 99	69	0	31	3600	2:
IRAS 22308+5812	48	39	13	770	3:
MSX LMC 1798	60	40	0	?	

Ref – 1: Menzies & Wolstencroft (1990); 2: Leisy & Dennefeld (2006); 3: Martín-Hernández *et al.* (2002)

from these results. However, the basis made of the three PDR components was found to be insufficient to obtain a satisfactory fit to the observations of PNe. Broad emission features around 8 and 12 μm , which are known to be prominent in post-Asymptotic Giant Branch (post-AGB) star spectra (see, e.g., Kwok *et al.* 2001), appear to be still present in the spectra of PNe. As template for these features we chose a simplified gaussian profiles which provide a reasonable fit to the bands observed in the post-AGB star IRAS 13416–6243, a prototype of C-type object from Peeters *et al.* (2002). In addition, the “7.7” μm band is shifted to near 7.8 μm in PNe and it was necessary to introduce an additional PAH-type (hereafter PAH^x) template with the peak position shifted to the red. Altogether, we have used a set of six template spectra to perform decomposition of the mid-IR spectra of PNe and c-H II regions (see Joblin *et al.* 2008 for details).

The obtained decompositions for two PNe (Vo 1 from MW and SMP LMC 99 from LMC) and two c-H II regions (IRAS 22308+5812 from MW and MSX LMC 1798 from LMC) are shown in Figure 1. Only PAH components are displayed (see figure caption). In Table 1 the relative amount of the different PAH components required for the fits is collected. A striking correlation is found between the required amount of PAH^x and electron densities (N_{el}) of the modeled sources. Joblin *et al.* (2008) have discussed the possibility that PAH^x may be in fact PAH anions, on the basis of trends in bands shift observed in theoretical data from Mallocci *et al.* (2007). To further progress on this suggestion we have computed the charge of PAH molecules in environments characterized by a different ionization parameter ($U_{\text{ion}} = Q_H / (4\pi R_S^2 N_{\text{el}} c)$; where Q_H is the number of H-ionizing photons, R_S is the Strömrgren radius and c is the light speed).

3. PAH charge

The results of the PAH charge computations are shown in Figure 3. The method used to compute the PAH charge is the same as described in Stasińska & Szerba (2001). However, the physics of the dust–gas–radiation interaction was updated according to Weingartner & Draine (2001) and the neutral PAH properties from Li & Draine (2001) were implemented. Our method used to compute the charge is applicable to large dust particles, where the dust charge may be treated as a continuous quantity and is probably not correct for so small dust particles as PAHs. However, we have checked that the computed PAH charge when rounded to the greatest integer charge less than the computed one (for instance -2 instead of -1.3) was never smaller than the most negative allowed charge (Eq. 24 of Weingartner & Draine 2001). In addition, our method reproduces well the known result that dust charge is approaching -2.16 eV for low U_{ion} (see PAH charge for $U_{\text{ion}}=1.e-5$ in Figure 3). Therefore, we think that our results give a representative value of PAH charge in different environments.




Figure 1. Mid-IR spectra after continuum subtraction and smoothing to the resolution of $\lambda/\Delta\lambda = 45$ (dotted line) and fit obtained using a set of six template spectra (solid line). Below each fit, only PAH components are displayed: PAH⁰ (dashed line), PAH⁺ (dash-dot line) and PAH^x (thin solid line). On each panel, PAH components are listed (from top to bottom) according to their contribution to the fit.

At least three of the four selected objects (the two PNe and IRAS 22308+5812) show (roughly) similar Q_H (they have similar effective temperatures and – roughly – luminosities) and R_S . The only striking difference between the parameters characterizing these objects, which enter into the ionization parameter, is the electron density (see Table 1). We may expect that the object with the highest electron density (Vo1) will have the smallest ionization parameter and therefore a lot of PAH anions, while IRAS 22308+5812 which has the lowest N_{el} will have the largest U_{ion} and therefore the lowest amount of PAH⁻. For LMC c-H II region (MSX LMC 1798) we do not know N_{el} , however, in general, the density of H II regions in galaxies with lower metallicity will be lower so their ionization parameters will be larger and therefore the percentage of PAH⁻ may be negligible. Note that Cox & Spaans (2006) have shown that the fraction of PAH anions is much lower in low metallicity environments, so our results are in line with their findings.




Figure 2. Calculated charge of PAH molecules with different size in environments characterized by a different ionization parameter U_{ion} (pluses – 1.e-5; circles – 1.e-4, squares – 1.e-3, diamonds – 1.e-2; triangles – 1.e-1.) and a gas electron temperature of 10^4 K.

Acknowledgements

R.Sz. acknowledges support from grant N203 019 31/2874 of the Science and High Education Ministry of Poland. Support from the French national program, Physique et Chimie du Milieu Interstellaire, is also acknowledged.

References

- Berné, O., Joblin, C., Deville, Y., Smith, J. D., Rapacioli, M., Bernard, J. P., Thomas, J., Reach, W., & Abergel, A. 2007, *A&A*, 469, 575
- Cox, N. & Spaans, M. 2006, *A&A*, 451, 973
- Joblin, C., Szczerba, R., Berné, O., & Szyszka, C. 2008, *A&A* submitted
- Kwok, S., Volk, K., & Bernath, P. 2001, *ApJ* (Letter), 554, L87
- Leisy, P. & Dennefeld, M. 2006, *A&A*, 456, 451
- Li, A., & Draine, B. T. 2001, *ApJ*, 554, 778
- Mallici, G., Joblin, C., & Mulas, G. 2007, *Chem. Phys.*, 332, 353
- Martín-Hernández, N. L., Peeters, E., Morisset, C., Tielens, A. G. G. M., Cox, P., Roelfsema, P. R., Baluteau, J.-P., Schaerer, D., Mathis, J. S., Damour, F., Churchwell, E., Kessler, M. F. 2002, *A&A*, 381, 606
- Menzies, J. W. & Wolstencroft, R. D. 1990, *MNRAS*, 247, 177
- Peeters, E., Hony, S., Van Kerckhoven, C., Tielens, A. G. G. M., Allamandola, L.J., Hudgins, D. M., & Bauschlicher, C. W. 2002, *A&A*, 390, 1089
- Rapacioli, M., Joblin, C., & Boissel, P. 2005, *A&A*, 429, 193
- Stasińska, G. & Szczerba, R. 2001, *A&A*, 379, 1024
- Weingartner, J. C. & Draine, B. 2001, *ApJS*, 134, 263

Discussion

SARRE: I would like to ask what molecular form PAH anions have? Are they stable closed shell systems, small example would be coronene with electrons attached, or are they radicals with electrons added to make them closed shell of anions?

SZCZERBA: It is difficult to make a definite statement. However, the spectral agreement is based on the theoretical spectral database of PAHs <http://astrochemistry.ca.astro.it/database/> in which most of the PAH species considered are closed shell as neutrals and open shell as singly ionized (cations or anions).



Maja Kazmierczak, Ryszard Szczerba, and Vera K. Rosenbush in front of the Wan Chai harbour (photo by Yuriy Serozhkin).



Philippe Bréchnignac and Rosario Brunetto.