

On closure operations in the space of subgroups and applications

DOMINIK FRANCOEUR[†] and ADRIEN LE BOUDEC[‡]

[†]*Universidad Autónoma de Madrid, Departamento de Matemáticas,
28049 Madrid, Spain*

(e-mail: dominik.francoeur@uam.es)

[‡]*CNRS, UMPA - ENS Lyon, 69364 Lyon, France*

(e-mail: adrien.le-boudec@ens-lyon.fr)

(Received 10 September 2024 and accepted in revised form 20 May 2025)

Abstract. We establish some interactions between uniformly recurrent subgroups (URSs) of a group G and coset topologies $\tau_{\mathcal{N}}$ on G associated to a family \mathcal{N} of normal subgroups of G . We show that when \mathcal{N} consists of finite index subgroups of G , there is a natural closure operation $\mathcal{H} \mapsto \text{cl}_{\mathcal{N}}(\mathcal{H})$ that associates to a URS \mathcal{H} another URS $\text{cl}_{\mathcal{N}}(\mathcal{H})$, called the $\tau_{\mathcal{N}}$ -closure of \mathcal{H} . We give a characterization of the URSs \mathcal{H} that are $\tau_{\mathcal{N}}$ -closed in terms of stabilizer URSs. This has consequences on arbitrary URSs when G belongs to the class of groups for which every faithful minimal profinite action is topologically free. We also consider the largest amenable URS \mathcal{A}_G and prove that for certain coset topologies on G , almost all subgroups $H \in \mathcal{A}_G$ have the same closure. For groups in which amenability is detected by a set of laws (a property that is variant of the Tits alternative), we deduce a criterion for \mathcal{A}_G to be a singleton based on residual properties of G .

Key words: profinite topology and other coset topologies, space of subgroups and uniformly recurrent subgroups, minimal actions on compact spaces, proximal and strongly proximal actions, C*-simplicity

2020 Mathematics Subject Classification: 37B05 (Primary); 20E26, 37C85, 37A55 (Secondary)

1. Introduction

Let G be a group[†]. We denote by \mathcal{N}_G the set of normal subgroups of G . Let $\mathcal{N} \subseteq \mathcal{N}_G$ be a family of normal subgroups of G that is filtering: for every $N_1, N_2 \in \mathcal{N}$, there exists

[†] The main situation we have in mind is when G is countable. Some of our results will require this countability assumption.

$N_3 \in \mathcal{N}$ such that $N_3 \leq N_1 \cap N_2$. There is a group topology $\tau_{\mathcal{N}}$ on G associated to \mathcal{N} , defined by declaring that the family of cosets gN , $g \in G$, $N \in \mathcal{N}$, forms a basis for $\tau_{\mathcal{N}}$. When \mathcal{N} is the family of all finite index normal subgroups of G , $\tau_{\mathcal{N}}$ is the profinite topology on G . If p is a prime and \mathcal{N} is the family of finite index normal subgroups N of G such that G/N is a p -group, $\tau_{\mathcal{N}}$ is the pro- p topology.

If H is a subgroup of G , the closure of H with respect to $\tau_{\mathcal{N}}$ is denoted by $\text{cl}_{\mathcal{N}}(H)$. In the case of the profinite topology, we use the shorter notation $\text{cl}(H)$. The closure operation defines a map

$$\text{cl}_{\mathcal{N}} : \text{Sub}(G) \rightarrow \text{Sub}(G), \quad H \mapsto \text{cl}_{\mathcal{N}}(H).$$

Here, $\text{Sub}(G)$ is the set of subgroups of G . That set is equipped with the topology inherited from the set $\{0, 1\}^G$ of all subsets of G , equipped with the product topology. The space $\text{Sub}(G)$ is a compact space. The group G acts on $\text{Sub}(G)$ by conjugation and this action is by homeomorphisms. The first object of study of this article is the behaviour of the map $\text{cl}_{\mathcal{N}}$ with respect to the dynamical system $G \curvearrowright \text{Sub}(G)$.

It follows from the definitions that the map $\text{cl}_{\mathcal{N}}$ is always increasing, idempotent and G -equivariant. In general, $\text{cl}_{\mathcal{N}}$ is far from being continuous. This failure of continuity already happens in the most classical case where $\tau_{\mathcal{N}}$ is the profinite topology. An elementary example illustrating this is the group $G = \mathbb{Z}[1/p]$ of p -adic rational numbers, for which the map cl is not upper semi-continuous on $\text{Sub}(G)$ (see Remark 3.6). Another example is $G = F_k$ (a finitely generated non-abelian free group of rank k). M. Hall showed that every finitely generated subgroup H of F_k verifies $\text{cl}(H) = H$ [Hal49] (that is, F_k is a LERF group). Since finitely generated subgroups always form a dense subset in the space of subgroups, it follows that cl is the identity on a dense set of points. However, cl is not the identity everywhere, for instance, because F_k admits infinite index subgroups H such that $\text{cl}(H) = F_k$ (e.g. any infinite index maximal subgroup). So cl is not lower semi-continuous on $\text{Sub}(F_k)$.

The starting result of this article is that if we restrict to minimal subsystems of $\text{Sub}(G)$ (that is, non-empty closed minimal G -invariant subsets of $\text{Sub}(G)$), the situation is better behaved. Recall that a minimal subsystem $\mathcal{H} \subset \text{Sub}(G)$ is called a URS (uniformly recurrent subgroup) [GW15].

PROPOSITION 1.1. *Let $\mathcal{N} \subseteq \mathcal{N}_G$ be a family of finite index normal subgroups of G and let \mathcal{H} be a URS of G . Then, the following hold:*

- (1) *the restriction $\text{cl}_{\mathcal{N}|_{\mathcal{H}}} : \mathcal{H} \rightarrow \text{Sub}(G)$ is upper semi-continuous;*
- (2) *there exists a unique URS contained in $\overline{\{\text{cl}_{\mathcal{N}}(H) : H \in \mathcal{H}\}}$, denoted $\text{cl}_{\mathcal{N}}(\mathcal{H})$ and called the $\tau_{\mathcal{N}}$ -closure of \mathcal{H} .*

The proposition also holds in a more general situation not necessarily requiring that \mathcal{N} consists of finite index subgroups of G (see Proposition 3.4).

Statement (2) says that there is a natural closure operation

$$\text{URS}(G) \rightarrow \text{URS}(G), \quad \mathcal{H} \mapsto \text{cl}_{\mathcal{N}}(\mathcal{H}),$$

where $\text{URS}(G)$ is the set of URSs of the group G . We say that a URS \mathcal{H} is closed for the topology $\tau_{\mathcal{N}}$ if $\text{cl}_{\mathcal{N}}(\mathcal{H}) = \mathcal{H}$. When G is a countable group, this happens if and only if there is a dense G_{δ} -set of points $H \in \mathcal{H}$ such that H is closed for the topology $\tau_{\mathcal{N}}$.

Recently, URSs were studied and appeared in a large number of works, including [BH21, FG23, LBMB18, LBMB22]. We refer notably to the introduction of [LBMB22] for more references. A common theme is to establish rigidity results saying that the set of URSs of certain groups is restricted, or to establish connections between certain group theoretic properties of the ambient group and properties of its URSs. We believe that, in certain situations, the above process $\mathcal{H} \mapsto \text{cl}_{\mathcal{N}}(\mathcal{H})$ and, more generally, the consideration of coset topologies on the ambient group can be profitably used to study properties of URSs. In §§4 and 5, we exhibit situations where it is indeed the case. In the remainder of this introduction, we shall describe these results.

When \mathcal{N} consists of finite index subgroups, the property that a URS \mathcal{H} is closed for the topology $\tau_{\mathcal{N}}$ admits the following natural characterization. Glasner and Weiss showed that to every minimal action of G on a compact space X , there is a naturally associated URS of G , called the stabilizer URS of X and denoted $S_G(X)$ [GW15]. We say that the action of G on a compact space X is *pro- \mathcal{N}* if $G \times X \rightarrow X$ is continuous, where G is equipped with the topology $\tau_{\mathcal{N}}$ (see Proposition 4.2 for characterizations of this property).

PROPOSITION 1.2. *Suppose that G is a countable group and that \mathcal{N} consists of finite index subgroups of G . For a URS \mathcal{H} of G , the following are equivalent:*

- (1) \mathcal{H} is closed for the topology $\tau_{\mathcal{N}}$;
- (2) there exists a *pro- \mathcal{N}* compact minimal G -space X such that $S_G(X) = \mathcal{H}$.

In the case of the profinite topology, the notion of *pro- \mathcal{N}* G -space coincides with the classical notion of profinite G -space. So, in that situation, the above proposition says that a URS \mathcal{H} is closed for the profinite topology if and only if \mathcal{H} is the stabilizer URS associated to a minimal profinite action of G . Consequences on all URSs can be drawn out of this when G belongs to the class of groups for which, for a faithful minimal compact G -space, profinite implies topologically free. See Proposition 4.17. This class of groups includes non-abelian free groups and, more generally, any group G admitting an isometric action on a hyperbolic space with unbounded orbits such that the G -action on its limit set is faithful. It also includes hereditarily just-infinite groups. Recall that a group G is just-infinite if G is infinite and G/N is finite for every non-trivial normal subgroup N , and G is hereditarily just-infinite if every finite index subgroup of G is just-infinite. We call a subgroup H of G *co-finitely dense* in G for the profinite topology if the profinite closure of H has finite index in G . For hereditarily just-infinite groups, we obtain the following proposition.

PROPOSITION 1.3. *Let G be a hereditarily just-infinite group and let \mathcal{H} be a non-trivial URS of G . Then, for every $H \in \mathcal{H}$, H is co-finitely dense in G for the profinite topology.*

In cases where we know *a priori* that the group G has the property that the only subgroups that are co-finitely dense are the finite index subgroups, we deduce that such a group G admits no continuous URS (a URS is continuous if it is not a finite set). See Corollary 4.20, and the surrounding discussion for context and examples.

Another setting in which we show that the consideration of a coset topology $\tau_{\mathcal{N}}$ is fruitful with respect to the study of URSs is the case of amenable URSs. A URS \mathcal{H} is amenable if it consists of amenable subgroups. Every group G admits a largest amenable URS (with respect to a natural partial order), which is the stabilizer URS associated to the action of G on its Furstenberg boundary (the largest minimal and strongly proximal compact G -space). This URS is denoted \mathcal{A}_G and is called the Furstenberg URS of G . The action of G on \mathcal{A}_G is minimal and strongly proximal. Here, \mathcal{A}_G is either a singleton, in which case we have $\mathcal{A}_G = \{\text{Rad}(G)\}$, where $\text{Rad}(G)$ is the amenable radical of G , or \mathcal{A}_G is continuous. We refer to [LBMB18] for a more detailed discussion.

Let \mathcal{F} denote the class of groups G such \mathcal{A}_G is a singleton. Equivalently, G belongs to \mathcal{F} if and only if every amenable URS of G lives inside the amenable radical of G . The class \mathcal{F} is known to be very large. It plainly contains amenable groups. It also contains all linear groups, all groups with non-vanishing ℓ^2 -Betti numbers, all hyperbolic groups and, more generally, all acylindrically hyperbolic groups. We refer to [BKKO17] for references and details. Examples of groups outside the class \mathcal{F} have been given in [LB17].

The following result provides a criterion for a group to be in \mathcal{F} that is based on residual properties of the group. If \mathcal{C} is a class of groups, a group G is residually- \mathcal{C} if the intersection of all normal subgroups N such that $G/N \in \mathcal{C}$ is trivial.

THEOREM 1.4. *Let G be a group such that every amenable subgroup of G is virtually solvable. If G is residually- \mathcal{F} , then G is in \mathcal{F} .*

We point out that this theorem is applicable without necessarily relying on other methods related to \mathcal{F} to verify the assumption that the group is residually- \mathcal{F} . The point is that the statement applies provided that G is residually- \mathcal{C} for some subclass \mathcal{C} of \mathcal{F} that is potentially much smaller. For instance, the theorem applies and is already interesting if G is residually finite.

Every group satisfying the Tits alternative has the property that every amenable subgroup is virtually solvable. We refer to §5.1 for examples of groups that are known to satisfy the Tits alternative.

One interest of such a statement is that it is based on intrinsic algebraic properties of the group. It does not require the group G to admit a rich action of geometric flavour, or to have an explicit minimal and strongly proximal compact G -space at our disposal. The residual properties are used as a tool in Theorem 1.4, but the confrontation of residual properties and the class \mathcal{F} is also motivated by the fact that it is not known whether there exist residually finite groups G with trivial amenable radical such that G does not belong to \mathcal{F} . The groups from [LB17] are never residually finite (and some of them are virtually simple).

As an application, Theorem 1.4 allows to recover the following result from [BKKO17].

COROLLARY 1.5. (Breuillard–Kalantar–Kennedy–Ozawa) *If G is a linear group, then G is in \mathcal{F} .*

The proof from [BKKO17] relies on linear group technology. Here, the argument to deduce Corollary 1.5 from Theorem 1.4 uses a reduction to the case of finitely generated groups, and then only appeals to Malcev’s theorem that finitely generated linear groups are residually finite, and the Tits alternative [Tit72].

The consideration of the class \mathcal{F} is also motivated by the result of Kalantar and Kennedy that a group G belongs to \mathcal{F} if and only if the quotient of G by its amenable radical is a C^* -simple group (that is, its reduced C^* -algebra is simple) [KK17]. We refer to the survey of de la Harpe [dlH07] for an introduction and historical developments on C^* -simple groups, and to the Bourbaki seminar of Raum for recent developments [Rau20]. Hence, using the result of Kalantar and Kennedy, Theorem 1.4 can be reinterpreted as a criterion to obtain C^* -simplicity (under the assumption on amenable subgroups) based on residual properties of the group. See Corollary 5.15. We are not aware of other results of this kind.

The proof of Theorem 1.4 is based on the following proposition, of independent interest. Given a group G , we denote by $\mathcal{N}_G(\mathcal{F})$ the set of normal subgroups of G such that $G/N \in \mathcal{F}$. The set $\mathcal{N}_G(\mathcal{F})$ is stable under taking finite intersections (Lemma 5.7), and we can consider the coset topology on G associated to $\mathcal{N}_G(\mathcal{F})$ (and more generally to a subset $\mathcal{N} \subseteq \mathcal{N}_G(\mathcal{F})$). The following result says that within the Furstenberg URS \mathcal{A}_G , almost all points have the same closure for such a topology (for technical reasons, we are led to make some countability assumptions).

PROPOSITION 1.6. *Let G be a countable group and let \mathcal{N} be a countable subset of $\mathcal{N}_G(\mathcal{F})$. Then, there exists a normal subgroup M of G and a comeagre subset $\mathcal{H}_0 \subseteq \mathcal{A}_G$ such that $\text{cl}_{\mathcal{N}}(H) = M$ for every $H \in \mathcal{H}_0$.*

The proof of the proposition makes crucial use of the strong proximality of the action of G on \mathcal{A}_G . The proof of Theorem 1.4 is easily deduced from the proposition. The additional point is to ensure that the closed normal subgroup M appearing in the conclusion of the proposition remains amenable, and this is where the two assumptions in the theorem are used. We refer to §5 for details. Here, we only mention that the actual setting in which we prove Theorem 1.4 does not necessarily require amenable subgroups to be virtually solvable. The assumption that we need is that amenability within subgroups of G can be detected by a set of laws (Definition 5.10), a property that can be thought of as a version of the Tits alternative (Proposition 5.11). See Theorem 5.13 for the more general formulation of the theorem.

2. Preliminaries

A space X is a G -space if G admits a continuous action $G \times X \rightarrow X$. Throughout the paper, we make the standing assumption that G -spaces are non-empty. The action (or the G -space X) is *minimal* if all orbits are dense. For $x \in X$, we write G_x for the stabilizer of x in G , and G_x^0 for the set of $g \in G$ such that g acts trivially on a neighbourhood of x . The action of G on X is free if $G_x = \{1\}$ for every $x \in X$, and *topologically free* if $G_x^0 = \{1\}$ for every $x \in X$.

Let X, Y be compact spaces. A continuous surjective map $\pi : Y \rightarrow X$ is called *irreducible* if every proper closed subset of Y has a proper image in X . If X, Y are compact G -spaces and $\pi : Y \rightarrow X$ is a continuous surjective G -equivariant map, we say that X is a factor of Y and that Y is an extension of X . When $\pi : Y \rightarrow X$ is irreducible, we also say that Y is an irreducible extension of X . If $\pi : Y \rightarrow X$ is irreducible, then X is minimal if and only if Y is minimal. Also for X, Y minimal, $\pi : Y \rightarrow X$ is irreducible if and only if it is *highly proximal*: for every $x \in X$, the fibre $\pi^{-1}(x)$ is compressible [AG77].

2.1. Semi-continuous maps. If Y is a locally compact space, we denote by 2^Y the space of closed subsets of Y , endowed with the Chabauty topology. The space 2^Y is compact.

Let X be a compact G -space. A map $\varphi: X \rightarrow 2^Y$ is *upper semi-continuous* if for every compact subset K of Y , $\{x \in X : \varphi(x) \cap K = \emptyset\}$ is open in X . It is *lower semi-continuous* if for every open subset U of Y , $\{x \in X : \varphi(x) \cap U \neq \emptyset\}$ is open in X . We say that φ is semi-continuous if it is either upper or lower semi-continuous.

Let $\varphi: X \rightarrow 2^Y$ be a semi-continuous map and $X_\varphi \subseteq X$ be the set of points where φ is continuous. Let

$$F_\varphi := \overline{\{(x, \varphi(x)) : x \in X\}} \subseteq X \times 2^Y,$$

$$E_\varphi := \overline{\{(x, \varphi(x)) : x \in X_\varphi\}} \subseteq F_\varphi(X),$$

$$T_\varphi := \overline{\{\varphi(x) : x \in X\}},$$

$$S_\varphi := \overline{\{\varphi(x) : x \in X_\varphi\}}.$$

We denote by $\eta: X \times 2^Y \rightarrow X$ and $p: X \times 2^Y \rightarrow 2^Y$ the projections to the first and second coordinate. If Y is second-countable, semi-continuity of φ implies that X_φ is a comeagre subset of X [Kur28, Theorem VII].

PROPOSITION 2.1. *Suppose X is a minimal compact G -space, Y is a locally compact G -space, and $\varphi: X \rightarrow 2^Y$ is G -equivariant and semi-continuous. Then, the following hold.*

- (i) F_φ has a unique non-empty minimal closed G -invariant subset E'_φ , and T_φ has a unique minimal closed G -invariant subset S'_φ , and $p(E'_\varphi) = S'_\varphi$.
- (ii) The extension $\eta: E'_\varphi \rightarrow X$ is highly proximal.

If moreover Y is second-countable, then $E'_\varphi = E_\varphi$ and $S'_\varphi = S_\varphi$.

Proof. See [Gla75, Theorem 2.3] and [AG77, Lemma I.1]. □

2.2. The space of subgroups and URSs. We denote by $\text{Sub}(G)$ the space of subgroups of G , equipped with the product topology from $\{0, 1\}^G$. It is a compact G -space, where G acts by conjugation. When G is countable, the space $\text{Sub}(G)$ is second-countable. If $H \in \text{Sub}(G)$, we denote by H^G the G -conjugates of H , that is, the G -orbit of H in $\text{Sub}(G)$.

A URS of G is a (non-empty) minimal closed G -invariant subset of $\text{Sub}(G)$. By Zorn's lemma, every (non-empty) closed G -invariant subset of $\text{Sub}(G)$ contains a URS. A URS is *finite* if it is a finite G -orbit. A URS that is not finite is called *continuous*. By minimality and compactness, a continuous URS has no isolated points. The singleton $\{\{1\}\}$ is called the trivial URS. If \mathcal{P} is a property of groups, we say that a URS \mathcal{H} has \mathcal{P} if H has \mathcal{P} for every $H \in \mathcal{H}$.

Definition 2.2. If \mathcal{H} is a URS of G , we denote by $\text{Env}(\mathcal{H})$ the subgroup generated by all subgroups H in \mathcal{H} , called the envelope of \mathcal{H} . The subgroup $\text{Env}(\mathcal{H})$ is normal in G and it is the smallest normal subgroup of G containing some subgroup $H \in \mathcal{H}$.

Every minimal compact G -space naturally gives rise to a URS [GW15].

PROPOSITION 2.3. *If X is a compact G -space, then the stabilizer map $S : X \rightarrow \text{Sub}(G)$, $x \mapsto G_x$, is G -equivariant and upper semi-continuous. In particular, if X is minimal, then Proposition 2.1 applies.*

Definition 2.4. If X is a minimal compact G -space, the unique URS contained in $\overline{\{G_x : x \in X\}}$ is denoted $S_G(X)$ and is called the *stabilizer URS* associated to the G -space X .

One verifies that the G -action on X is topologically free if and only if the URS $S_G(X)$ is trivial.

LEMMA 2.5. *Let H, K, L be subgroups of G such that $H \leq L$. If K belongs to the closure of the L -orbit of H in $\text{Sub}(G)$, then $K \leq L$.*

Proof. The subset $\text{Sub}(L)$ is a closed subset of $\text{Sub}(G)$ and contains the L -orbit of H since $H \leq L$. □

LEMMA 2.6. *Let N be a normal subgroup of G . Then, the map $\text{Sub}(G) \rightarrow \text{Sub}(G)$, $H \mapsto HN$, is G -equivariant and lower semi-continuous.*

Proof. It is G -equivariant because N is normal in G . Since G is discrete, lower semi-continuity means that for every $g \in G$ and every $H \in \text{Sub}(G)$ such that $g \in HN$, there is a neighbourhood of H in which $g \in H'N$ remains true. If $h \in H$ is such that $g \in hN$, then the set of subgroups of G containing h is such a neighbourhood. □

2.3. *Coset topologies on groups.* Let G be a group. We denote by \mathcal{N}_G the set of normal subgroups of G . We make the standing convention that when considering a family $\mathcal{N} \subseteq \mathcal{N}_G$ of normal subgroups of G , we always assume that \mathcal{N} is non-empty.

Definition 2.7. Let $\mathcal{N} \subseteq \mathcal{N}_G$. If E is a subset of G , we denote

$$\text{cl}_{\mathcal{N}}(E) = \bigcap_{N \in \mathcal{N}} EN.$$

We say that \mathcal{N} is filtering if for every $N_1, N_2 \in \mathcal{N}$, there exists $N_3 \in \mathcal{N}$ such that $N_3 \leq N_1 \cap N_2$. We record the following proposition [Bou71, Ch. III].

PROPOSITION 2.8. *Fix $\mathcal{N} \subseteq \mathcal{N}_G$. Then:*

- (1) *the family of cosets gN , $g \in G$, $N \in \mathcal{N}$, forms a subbasis for a group topology $\tau_{\mathcal{N}}$ on G ;*
- (2) *the topology $\tau_{\mathcal{N}}$ is Hausdorff if and only if $\bigcap_{N \in \mathcal{N}} N = \{1\}$;*
- (3) *suppose that \mathcal{N} is filtering. Then, for every subset E of G , the closure of E with respect to $\tau_{\mathcal{N}}$ is equal to $\text{cl}_{\mathcal{N}}(E)$.*

If \mathcal{C} is a class of groups, we denote by $\mathcal{N}_G(\mathcal{C})$ the normal subgroups of G such that $G/N \in \mathcal{C}$. Note that a group G is residually- \mathcal{C} if and only if $\bigcap_{N \in \mathcal{N}_G(\mathcal{C})} N = \{1\}$.

When \mathcal{C} is the class of all finite groups and $\mathcal{N} = \mathcal{N}_G(\mathcal{C})$, $\tau_{\mathcal{N}}$ is the profinite topology on G . For simplicity, we write $\text{cl}(H)$ for the closure in the profinite topology. When \mathcal{C} is

the class of all finite p -groups (p is a prime number) and $\mathcal{N} = \mathcal{N}_G(\mathcal{C})$, $\tau_{\mathcal{N}}$ is the pro- p topology. In that case, we write $\text{cl}_p(H)$ for the closure in the pro- p topology.

2.4. Laws. Let $w = w(x_1, \dots, x_k)$ be a word in k letters x_1, \dots, x_k , meaning that w is an element of the free group F_k freely generated by x_1, \dots, x_k . We assume w is non-trivial. Given a group G , the word w naturally defines a map $G^k \rightarrow G$, a k -tuple (g_1, \dots, g_k) being mapped to the element $w(g_1, \dots, g_k)$ of G that is obtained by replacing each x_i by g_i . We denote by $\Sigma_w(G) \subseteq G^k$ the set of (g_1, \dots, g_k) such that $w(g_1, \dots, g_k) = 1$. We say that G satisfies the law w if $\Sigma_w(G) = G^k$.

Example 2.9. Abelian groups are the groups that satisfy the law $w_1 = [x_1, x_2] = x_1 x_2 x_1^{-1} x_2^{-1}$. More generally, let $(w_\ell)_{\ell \geq 1}$ be the sequence of laws defined inductively by the rule: if w_ℓ is already defined and involves 2^ℓ variables, then

$$w_{\ell+1}(x_1, \dots, x_{2^{\ell+1}}) = [w_\ell(x_1, \dots, x_{2^\ell}), w_\ell(x_{2^\ell+1}, \dots, x_{2^{\ell+1}})].$$

Then, a group G is solvable of length at most ℓ if and only if G satisfies the law w_ℓ .

LEMMA 2.10. Suppose G is a Hausdorff topological group and let $w \in F_k$. Then, $\Sigma_w(G)$ is a closed subset of G^k . In particular, if a subgroup H of G satisfies the law w , then so does its closure.

Proof. Since G is Hausdorff, $\{1\}$ is closed in G . The map $G^k \rightarrow G$ associated to w being continuous, the preimage $\Sigma_w(G)$ of $\{1\}$ is a closed subset of G^k . \square

3. The $\tau_{\mathcal{N}}$ -closure of a URS

Let \mathcal{H} be a closed subset of $\text{Sub}(G)$, and L a subgroup of G . Recall that L^G is the set of G -conjugates of L . For every $\Sigma \subseteq L^G$, we write

$$\mathcal{H}_\Sigma = \{H \in \mathcal{H} : \text{for all } K \in L^G, H \subset K \Leftrightarrow K \in \Sigma\}.$$

The relevance of this definition is indicated by the following two lemmas.

LEMMA 3.1. If $\mathcal{H}_\Sigma \cap \mathcal{H}_{\Sigma'} \neq \emptyset$, then $\Sigma = \Sigma'$, and \mathcal{H} is the disjoint union of the \mathcal{H}_Σ when Σ ranges over subsets of L^G .

Proof. The first assertion is a consequence of the definitions. The second assertion is also clear since for every $H \in \mathcal{H}$, one has $H \in \mathcal{H}_\Sigma$ with $\Sigma = \{K \in L^G : H \subset K\}$. \square

LEMMA 3.2. Let L be a subgroup of G such that L^G is finite. Suppose \mathcal{H} is a URS of G . Then, \mathcal{H}_Σ is a clopen subset of \mathcal{H} for every $\Sigma \subseteq L^G$.

Proof. For H in \mathcal{H} , we let $n(H)$ be the number of conjugates of L containing H . By our assumption, the number $n(H)$ is finite. We claim that $n(H)$ is constant on \mathcal{H} . To see this, take $H \in \mathcal{H}$ such that $n(H) = r$ is minimal. Since not being contained in a subgroup is an open condition, one can find a neighbourhood V of H such that $n(H') \leq r$ for every $H' \in V$. Hence, by minimality of r , we have $n(H') = r$ for every $H' \in V$. Now, for every

$K \in \mathcal{H}$, by minimality of the G -action on \mathcal{H} , the subset V contains a conjugate of K . Since $n(K)$ is invariant under conjugation, we deduce $n(K) = r$.

Now, fix $\Sigma \subseteq L^G$ such that \mathcal{H}_Σ is non-empty and let $H \in \mathcal{H}_\Sigma$. Again, there is a neighbourhood V of H in \mathcal{H} such that for every H' in V , we have $H' \not\subseteq J$ for every $J \in L^G \setminus \Sigma$. Moreover, by the previous paragraph, we have $n(H') = n(H)$. Hence, by the pigeonhole principle, we deduce that $H' \subset J$ for every $J \in \Sigma$. This shows that \mathcal{H}_Σ is open. Since the family (\mathcal{H}_Σ) forms a partition of \mathcal{H} by Lemma 3.1, it follows that \mathcal{H}_Σ is also closed. \square

Definition 3.3. Let $\mathcal{N} \subseteq \mathcal{N}_G$. We say that a URS \mathcal{H} of G is \mathcal{N} -finitary if $(HN)^G$ is finite for every $H \in \mathcal{H}$, $N \in \mathcal{N}$.

PROPOSITION 3.4. Let $\mathcal{N} \subseteq \mathcal{N}_G$ and let \mathcal{H} be a URS of G that is \mathcal{N} -finitary. Then, the map $\mathcal{H} \rightarrow \text{Sub}(G)$, $H \mapsto \text{cl}_{\mathcal{N}}(H)$, is upper semi-continuous.

Proof. Let K be a finite subset of G and let $H \in \mathcal{H}$ such that $\text{cl}_{\mathcal{N}}(H) \cap K = \emptyset$. One shall prove that $\text{cl}_{\mathcal{N}}(H') \cap K = \emptyset$ remains true for every H' inside a neighbourhood of H in \mathcal{H} . Let $g \in K$. By definition of $\text{cl}_{\mathcal{N}}(H)$, there exists $N_g \in \mathcal{N}$ such that $g \notin HN_g$. Since $L = HN_g$ verifies that $(HN_g)^G$ is finite, according to Lemma 3.2, one can find a neighbourhood V_g of H in \mathcal{H} such that $H' \leq HN_g$ for every $H' \in V_g$. *A fortiori* we have $H'N_g \leq HN_g$ and hence $\text{cl}_{\mathcal{N}}(H') \leq HN_g$. Since K is finite, taking the intersection over all $g \in K$, we obtain a neighbourhood V of H in \mathcal{H} such that $\text{cl}_{\mathcal{N}}(H') \leq \bigcap_{g \in K} HN_g$ for every $H' \in V$. Since K does not intersect $\bigcap_{g \in K} HN_g$, the neighbourhood V satisfies the required property. \square

Remark 3.5. If \mathcal{N} consists of finite index normal subgroups of G , then trivially, every URS of G is \mathcal{N} -finitary. Hence, the previous proposition applies.

Remark 3.6. Here, we still consider the case where \mathcal{N} consists of finite index normal subgroups of G and we point out that, in general, the map $\text{Sub}(G) \rightarrow \text{Sub}(G)$, $H \mapsto \text{cl}_{\mathcal{N}}(H)$, is *not* upper semi-continuous. Therefore, it is necessary to restrict to a URS in Proposition 3.4 to obtain upper semi-continuity. As an illustration, consider the group $G = \mathbb{Z}[1/p]$ of p -adic rational numbers. For $n \geq 1$, let $H_n = p^n\mathbb{Z}$. Then, G/H_n is a Prüfer p -group and, hence, has no proper finite index subgroup. So G has no proper finite index subgroup containing H_n or, equivalently, $\text{cl}(H_n) = G$. However, (H_n) converges to the trivial subgroup in $\text{Sub}(G)$, which is closed for the profinite topology since G is residually finite. Hence, $H \mapsto \text{cl}(H)$ is not upper semi-continuous.

COROLLARY 3.7. Let $\mathcal{N} \subseteq \mathcal{N}_G$, and \mathcal{H} be a URS of G that is \mathcal{N} -finitary. Then, the set

$$\overline{\{\text{cl}_{\mathcal{N}}(H) : H \in \mathcal{H}\}}$$

contains a unique URS of G that will be denoted $\text{cl}_{\mathcal{N}}(\mathcal{H})$. Moreover, when G is countable, there is a dense G_δ subset $\mathcal{H}_0 \subseteq \mathcal{H}$ such that

$$\text{cl}_{\mathcal{N}}(\mathcal{H}) = \overline{\{\text{cl}_{\mathcal{N}}(H) : H \in \mathcal{H}_0\}}.$$

Proof. Proposition 3.4 asserts that $\mathcal{H} \rightarrow \text{Sub}(G)$, $H \mapsto \text{cl}_{\mathcal{N}}(H)$, is upper semi-continuous. This allows to invoke Proposition 2.1, from which the statement follows. \square

COROLLARY 3.8. Let $\mathcal{N} \subseteq \mathcal{N}_G$, and \mathcal{H} be a URS of G that is \mathcal{N} -finitary.

- (1) If there exists $H \in \mathcal{H}$ such that $H = \text{cl}_{\mathcal{N}}(H)$, then $\mathcal{H} = \text{cl}_{\mathcal{N}}(\mathcal{H})$.
- (2) If G is countable, then $\text{cl}_{\mathcal{N}}(\text{cl}_{\mathcal{N}}(\mathcal{H})) = \text{cl}_{\mathcal{N}}(\mathcal{H})$.

Proof. The assumption in condition (1) implies that

$$\mathcal{H} \cap \overline{\{\text{cl}_{\mathcal{N}}(H) : H \in \mathcal{H}\}} \neq \emptyset.$$

So by minimality, \mathcal{H} is contained in $\overline{\{\text{cl}_{\mathcal{N}}(H) : H \in \mathcal{H}\}}$. Corollary 3.7 then implies $\mathcal{H} = \text{cl}_{\mathcal{N}}(\mathcal{H})$. Condition (2) follows from the second statement in Corollary 3.7 and condition (1). \square

Definition 3.9. Suppose that \mathcal{N} is filtering and let \mathcal{H} be a URS of G that is \mathcal{N} -finitary. We say that a URS \mathcal{H} is closed for the topology $\tau_{\mathcal{N}}$ if $\text{cl}_{\mathcal{N}}(\mathcal{H}) = \mathcal{H}$.

4. On profinite closures of a URS

4.1. *Profinutely closed URSs and profinite actions.* In all this section, we assume that $\mathcal{N} \subseteq \mathcal{N}_G$ is filtering and that \mathcal{N} consists of finite index subgroups of G . Let $\widehat{G}^{\mathcal{N}}$ be the inverse limit of the inverse system of finite groups G/N , $N \in \mathcal{N}$, and $\psi : G \rightarrow \widehat{G}^{\mathcal{N}}$ be the associated canonical group homomorphism (for simplicity, we omit \mathcal{N} in the notation of ψ). The group $\widehat{G}^{\mathcal{N}}$ is profinite and $\psi : G \rightarrow \widehat{G}^{\mathcal{N}}$ is continuous, where G is equipped with the topology $\tau_{\mathcal{N}}$. Recall that if H is a subgroup of G , one has $\psi^{-1}(\overline{\psi(H)}) = \text{cl}_{\mathcal{N}}(H)$.

PROPOSITION 4.1. Let \mathcal{N} be as above. Then, the following hold.

- (1) Let H be a subgroup of G such that $H = \text{cl}_{\mathcal{N}}(H)$. Then, the closure of the conjugacy class of H contains a unique URS.
- (2) Let \mathcal{H} be a URS of G . Let $H \in \mathcal{H}$ and $L = \overline{\psi(H)}$. Then, the stabilizer URS associated to the left translation action of G on $\widehat{G}^{\mathcal{N}}/L$ is equal to $\text{cl}_{\mathcal{N}}(\mathcal{H})$.

Proof. Write $L = \overline{\psi(H)}$ and $X = \widehat{G}^{\mathcal{N}}/L$, which is a minimal compact G -space since G has dense image in \widehat{G} . The stabilizer of the coset $L \in X$ in G is $\psi^{-1}(L) = \text{cl}_{\mathcal{N}}(H)$. So in the case where $H = \text{cl}_{\mathcal{N}}(H)$, one has

$$\overline{H^G} \subseteq \overline{\{G_x : x \in X\}}.$$

By Zorn's lemma, $\overline{H^G}$ contains at least one URS, and it follows that it contains exactly one because $\{G_x : x \in X\}$ has this property by Proposition 2.3. Hence, part (1) holds.

For part (2), we have

$$\overline{\{G_x : x \in X\}} \cap \overline{\{\text{cl}_{\mathcal{N}}(K) : K \in \mathcal{H}\}} \neq \emptyset.$$

Each one of these two sets contains a unique URS, namely $S_G(X)$ and $\text{cl}_{\mathcal{N}}(\mathcal{H})$. Hence, equality $S_G(X) = \text{cl}_{\mathcal{N}}(\mathcal{H})$ follows. \square

PROPOSITION 4.2. *Let \mathcal{N} be as above and let X be a compact totally disconnected G -space. The following are equivalent:*

- (1) $G \times X \rightarrow X$ is continuous, where G is equipped with the topology $\tau_{\mathcal{N}}$;
- (2) for every clopen subset U of X , the stabilizer of U in G is open for the topology $\tau_{\mathcal{N}}$;
- (3) $G \times X \rightarrow X$ extends to a continuous action of $\widehat{G}^{\mathcal{N}}$ on X .

If X is a minimal G -space, these are also equivalent to:

- (4) there exists a closed subgroup L of $\widehat{G}^{\mathcal{N}}$ such that X is isomorphic to $\widehat{G}^{\mathcal{N}}/L$ as a G -space (where G acts on $\widehat{G}^{\mathcal{N}}/L$ by left translations).

Proof. Since X is totally disconnected, clopen subsets form a basis of the topology on X . The equivalence between parts (1) and (2) is therefore a consequence of the definitions. Suppose these conditions hold and let L be the closure of the image of G in the group $\text{Homeo}(X)$. Since it follows in particular from part (2) that every clopen subset of X has a finite G -orbit, the group L is a profinite group. Since $G \rightarrow L$ is continuous, by the universal property of $\widehat{G}^{\mathcal{N}}$ [Wil98, Propositions 1.4.1–1.4.2], $G \rightarrow L$ extends to a continuous homomorphism $\widehat{G}^{\mathcal{N}} \rightarrow L$. So part (3) holds. Finally, part (3) implies part (1) because $\psi : G \rightarrow \widehat{G}^{\mathcal{N}}$ is continuous.

The last statement is clear since a minimal continuous action of a compact group on a compact space is necessarily transitive. \square

Definition 4.3. The G -action on X is called *pro- \mathcal{N}* if it satisfies the equivalent conditions (1)–(2)–(3). We also say that the G -space X is *pro- \mathcal{N}* .

In the case where \mathcal{N} consists of all finite index normal subgroups of G , this corresponds to the common notion of profinite G -space.

PROPOSITION 4.4. *Let \mathcal{H} be a URS of a countable group G . Then, the following are equivalent:*

- (1) \mathcal{H} is closed for the topology $\tau_{\mathcal{N}}$;
- (2) for every $H \in \mathcal{H}$, the stabilizer URS associated to the left translation action of G on $\widehat{G}^{\mathcal{N}}/\overline{\psi(H)}$ is equal to \mathcal{H} ;
- (3) there exists a minimal G -space X that is *pro- \mathcal{N}* such that $S_G(X) = \mathcal{H}$.

Proof. Proposition 4.1 implies that parts (1) and (2) are equivalent. Part (2) clearly implies part (3), so we only have to see that part (3) implies part (1). Let X be a minimal G -space that is *pro- \mathcal{N}* that admits \mathcal{H} as a stabilizer URS. By Proposition 4.2, there exists a closed subgroup L of $\widehat{G}^{\mathcal{N}}$ such that X is isomorphic to $\widehat{G}^{\mathcal{N}}/L$ as a G -space. The stabilizers in G for the action on $\widehat{G}^{\mathcal{N}}/L$ are closed for the topology $\tau_{\mathcal{N}}$ on G . Since G is countable, there is a dense set of points x in $\widehat{G}^{\mathcal{N}}/L$ such that $G_x \in S_G(\widehat{G}^{\mathcal{N}}/L)$ (Proposition 2.3). Since $S_G(\widehat{G}^{\mathcal{N}}/L)$ is equal to \mathcal{H} by assumption, it follows that \mathcal{H} contains some elements that are closed for the topology $\tau_{\mathcal{N}}$. By Corollary 3.8, this implies $\mathcal{H} = \text{cl}(\mathcal{H})$. \square

Remark 4.5. Matte Bon, Tsankov and Elek showed that every URS \mathcal{H} is equal to the stabilizer URS associated to some compact G -space [Ele18, MBT20], and among the compact G -spaces associated to \mathcal{H} , there is a unique one that is universal in a certain

sense [MBT20]. We point out that this G -space is very different from the G -space $\widehat{G}^N/\overline{\psi(H)}$ associated to the specific setting considered in Proposition 4.4.

Remark 4.6. In the case of the profinite topology, Proposition 4.4 says that a URS \mathcal{H} is closed for the profinite topology if and only if \mathcal{H} is the stabilizer URS associated to a minimal profinite G -space. It is worth noting that if \mathcal{H} is such a URS, then the G -action on \mathcal{H} need *not* be profinite. Such a phenomenon has been exploited by Matte Bon [MB17] and Nekrashevych [Nek20].

We end this section by showing that, in general, the restriction of cl to a URS is not continuous. Recall that a closed subset F of a space X is regular if F equals the closure of its interior.

LEMMA 4.7. *Let X be a compact minimal G -space such that $\text{Fix}_X(g)$ is a regular closed set of X for every $g \in G$. Then, for every $x \in X$, we have $G_x \leq \text{cl}(G_x^0)$.*

Proof. Fix $x \in X$, $g \in G_x$, and a finite index normal subgroup N of G . We want to see that $g \in G_x^0 N$. Since N has finite index in G and G acts minimally on X , each minimal closed N -invariant subset of X is clopen, and the minimal closed N -invariant subsets form a finite partition $\{U_1, \dots, U_n\}$ of X . Let U_i be the one containing x . Since U_i is a neighbourhood of x and $g \in G_x$, the assumption that $\text{Fix}_X(g)$ is regular implies that there exists a non-empty open subset $V \subseteq U_i$ on which g acts trivially. Since N acts minimally on U_i , one can find $h \in N$ such that $y = hx \in V$. It follows that $g \in G_y^0 = hG_x^0 h^{-1}$, and since $h \in N$, we deduce that $g \in G_x^0 N$. \square

PROPOSITION 4.8. *Suppose that G is countable. Let X be a minimal profinite compact G -space and $\mathcal{H} = S_G(X)$. Suppose that $\text{Fix}_X(g)$ is a regular closed set of X for every $g \in G$ and the stabilizer map $X \rightarrow \text{Sub}(G)$ is not continuous on X . Then, $\text{cl} : \mathcal{H} \rightarrow \text{Sub}(G)$ is the identity on a dense set of points, but is not the identity everywhere on \mathcal{H} . In particular, it is not continuous.*

Proof. By Proposition 4.4, the URS \mathcal{H} is closed for the profinite topology. Since G is countable, this means that there is a dense set of $H \in \mathcal{H}$ such that $\text{cl}(H) = H$. Since $x \mapsto G_x$ is not continuous on X , one easily verifies that one can find $x \in X$ and $H \in \mathcal{H}$ such that $H \not\leq G_x$ and $G_x^0 \leq H$ (see [LBMB18, Lemma 2.8]). It follows from Lemma 4.7 that $G_x \leq \text{cl}(G_x^0) \leq \text{cl}(H)$ and, hence, H is properly contained in $\text{cl}(H)$ since it is properly contained in G_x . \square

Remark 4.9. An example of the above situation is provided by G the Grigorchuk group and X the boundary of the defining rooted tree of G [Gri11, §7].

4.2. Hereditarily minimal actions.

Definition 4.10. A compact G -space X is *hereditarily minimal* if every finite index subgroup of G acts minimally on X .

Recall that every minimal and proximal G -space is hereditarily minimal [Gla76, Lemma 3.2].

PROPOSITION 4.11. *Let \mathcal{H} be a URS of G that is hereditarily minimal. Then, for every $H, K \in \mathcal{H}$, we have $\text{cl}(H) = \text{cl}(K) = \text{cl}(\text{Env}(\mathcal{H}))$.*

This holds in particular if $\mathcal{H} = S_G(X)$ with X a hereditarily minimal compact G -space.

Proof. Let L be a finite index subgroup of G such that $H \leq L$. Since L acts minimally on \mathcal{H} , Lemma 2.5 says that $K \leq L$. Consequently, $\text{cl}(H) = \text{cl}(K)$. Since \mathcal{H} is G -invariant, it follows that this common subgroup is normal in G . Call it N . We shall see that $N = \text{cl}(\text{Env}(\mathcal{H}))$. Since $H \leq \text{Env}(\mathcal{H})$ for every $H \in \mathcal{H}$, the inclusion $N \leq \text{cl}(\text{Env}(\mathcal{H}))$ is clear. However, N contains $\text{Env}(\mathcal{H})$ since N contains all elements of \mathcal{H} . Since N is closed in the profinite topology, N contains $\text{cl}(\text{Env}(\mathcal{H}))$. Hence, equality holds.

As for the last claim, it follows from the fact that Propositions 2.1 and 2.3 ensure that the stabilizer URS associated to a hereditarily minimal compact G -space is itself hereditarily minimal. \square

We refer to §2.4 for the definition of a law.

THEOREM 4.12. *Suppose G is a residually finite group. Let \mathcal{H} be a URS of G that is hereditarily minimal and suppose that $H \in \mathcal{H}$ satisfies the law w . Then, $\text{Env}(\mathcal{H})$ also satisfies the law w .*

Proof. G is residually finite, so the profinite topology on G is Hausdorff. Hence, Lemma 2.10 says that $\text{cl}(H)$ still satisfies w . Since $\text{cl}(H)$ contains $\text{Env}(\mathcal{H})$ by Proposition 4.11, $\text{Env}(\mathcal{H})$ also satisfies w . \square

Without the hereditarily minimal assumption, it does not hold in general that a URS satisfying a law w lives inside a normal subgroup of G satisfying w , as the following example shows.

Example 4.13. Let (F_n) be a sequence of non-abelian finite groups. Suppose that for every n , there is an abelian subgroup E_n of F_n such that the only normal subgroup N of F_n containing E_n is $N = F_n$. Let $\mathbb{G} = \prod_n F_n$ and let G be a countable dense subgroup of \mathbb{G} containing $\bigoplus_n F_n$. Consider the G -action on $X = \prod_n F_n/E_n$. This action is minimal and G_x is abelian for every $x \in X$. In particular, every $H \in \mathcal{H} := S_G(X)$ is abelian. However, $\text{Env}(\mathcal{H})$ contains the normal closure in G of $\bigoplus_n E_n$. In particular, $\text{Env}(\mathcal{H})$ contains $\bigoplus_n F_n$ and, hence, $\text{Env}(\mathcal{H})$ is not abelian.

Examples as above can be found among finitely generated groups. For instance, the groups constructed by B.H. Neumann in [Neu37, Ch. III] satisfy these properties. Given a sequence $\mathbf{u} = (u_n)_n$ of odd integers greater than 5, B.H. Neumann constructs a subgroup $G_{\mathbf{u}}$ of the product $\prod_n \text{Alt}(u_n)$. The subgroup $G_{\mathbf{u}}$ is generated by two elements and $G_{\mathbf{u}}$ contains $\bigoplus_n \text{Alt}(u_n)$ [Neu37, Ch. III(16)]. Since $\text{Alt}(u_n)$ is simple, $G_{\mathbf{u}}$ therefore falls into the above setting with $F_n = \text{Alt}(u_n)$ and E_n any non-trivial abelian subgroup of $\text{Alt}(u_n)$.

Remark 4.14. We note that in the above examples $G_{\mathbf{u}}$, we actually have an abelian URS \mathcal{H} for which $\text{Env}(\mathcal{H})$ satisfies no law at all (since a given non-trivial law cannot be satisfied by all finite groups, and hence neither by arbitrary large finite alternating groups).

4.3. PIF groups. In this subsection, we focus on the class of groups for which, for a faithful minimal compact G -space, profinite implies topologically free.

Definition 4.15. We say that a group G is *PIF* if for every faithful minimal compact G -space X , if the G -action on X is profinite, then it is topologically free.

This notion was studied notably by Grigorchuk. We will use the following proposition from [Gri11]. Recall that a group G is *just-infinite* (JI) if G is infinite and G/N is finite for every non-trivial normal subgroup N . Also, G is *hereditarily just-infinite* (HJI) if every finite index subgroup of G is JI.

PROPOSITION 4.16. *Each one of the following conditions implies that G is PIF:*

- (1) *for every non-trivial subgroup $H_1, H_2 \leq G$ such that the normalizer $N_G(H_i)$ of H_i has finite index in G for $i = 1, 2$, we have that $H_1 \cap H_2$ is non-trivial;*
- (2) *G is hereditarily just-infinite.*

Proof. The first assertion is [Gri11, Proposition 4.11] (the formulation there is not quite the same, but the argument is the same). The second assertion follows from the first one because every HJI group satisfies assertion (1). \square

The first condition of the proposition is satisfied, for instance, by non-abelian free groups and also by all Gromov-hyperbolic groups with no non-trivial finite normal subgroup, and more generally by any group G admitting an isometric action on a hyperbolic space X with unbounded orbits such that the G -action on its limit set $\partial_X G$ is faithful.

PROPOSITION 4.17. *Suppose G is PIF and let \mathcal{H} be a non-trivial URS of G . Then, there exists a non-trivial normal subgroup N of G such that $N \leq \text{cl}(H)$ for every $H \in \mathcal{H}$.*

Proof. Note that since \mathcal{H} is not the trivial URS, $\text{cl}(\mathcal{H})$ is not the trivial URS either. Let $H \in \mathcal{H}$ and $L = \overline{\psi(H)}$, where $\psi : G \rightarrow \widehat{G}$ is the canonical map from G to its profinite completion. By Proposition 4.1, the stabilizer URS associated to the left translation action of G on \widehat{G}/L is equal to $\text{cl}(\mathcal{H})$ and, hence, is not trivial. This means that the action of G on \widehat{G}/L is not topologically free. Since this action is profinite and G is PIF, the action cannot be faithful. So there is a non-trivial normal subgroup N of G that is contained in the stabilizer in G of the coset L , which is $\psi^{-1}(L) = \text{cl}(H)$. Upper semi-continuity of cl on \mathcal{H} then implies that $N \leq \text{cl}(H')$ for every $H' \in \mathcal{H}$. \square

Definition 4.18. We say that a subgroup H of a group G is *co-finitely dense for the profinite topology* if $\text{cl}(H)$ is a finite index subgroup of G .

COROLLARY 4.19. *Suppose G is hereditarily just-infinite and let \mathcal{H} be a non-trivial URS of G . Then, for every $H \in \mathcal{H}$, H is co-finitely dense in G for the profinite topology.*

Proof. Proposition 4.16 says that G is PIF. So Proposition 4.17 applies and gives a non-trivial normal subgroup N such that $N \leq \text{cl}(H)$ for every $H \in \mathcal{H}$. By the assumption, N must have finite index and the conclusion follows. \square

Recall that every HJI-group is either virtually simple or residually finite. Corollary 4.19 is void for virtually simple groups, so the focus here is on residually finite HJI-groups. By Margulis normal subgroup theorem, every irreducible lattice Γ in a connected semisimple Lie group \mathbf{G} (with trivial centre and no compact factor) of rank ≥ 2 is HJI. Under the assumption that every simple factor of the ambient Lie group \mathbf{G} has rank ≥ 2 , it is known that every non-trivial URS of Γ is just the conjugacy class of a finite index subgroup [BH21, Corollary F]. The normal subgroup theorem of Bader and Shalom asserts that any irreducible cocompact lattice Γ in a product $\mathbf{G}_1 \times \mathbf{G}_2$, where $\mathbf{G}_1, \mathbf{G}_2$ are compactly generated topologically simple locally compact groups, is HJI [BS06]. In this setting, the URSs of Γ are not understood.

Following [Cor06], we shall say that a group G has *property (PF)* if for every subgroup H of G , H is co-finitely dense in G for the profinite topology only if H has finite index. Let p be a prime number. Following [EJZ13], we say that a group G is *weakly p -LERF* if for every subgroup H of G , the closure $\text{cl}_p(H)$ of H for the pro- p topology has finite index in G only if H has finite index. Note that applying the definition of weakly p -LERF to the trivial subgroup, we see that a just-infinite group that is weakly p -LERF is necessarily residually- p . Every group that is weakly p -LERF has property (PF). Among their striking properties, the finitely generated groups obtained as the outputs of the process carried out in [EJZ13] are HJI and weakly p -LERF.

COROLLARY 4.20. *Suppose G is hereditarily just-infinite and has property (PF). Then, G has no continuous URS.*

Proof. Let \mathcal{H} be a URS of G . If \mathcal{H} is trivial, then there is nothing to show. Otherwise, for every $H \in \mathcal{H}$, $\text{cl}(H)$ has finite index in G by Corollary 4.19. Hence, so does H since G has property (PF). In particular, H has only finitely many conjugates, that is, \mathcal{H} is finite. \square

5. The Furstenberg URS

Definition 5.1. Given two closed subsets $\mathcal{X}_1, \mathcal{X}_2 \subset \text{Sub}(G)$, we write $\mathcal{X}_1 \preccurlyeq \mathcal{X}_2$ if there exist $H_1 \in \mathcal{X}_1$ and $H_2 \in \mathcal{X}_2$ such that $H_1 \leq H_2$.

One verifies that, when restricted to the set $\text{URS}(G)$, the relation \preccurlyeq is a partial order [LMB18, Corollary 2.15].

Recall that a compact G -space X is strongly proximal if the orbit closure of every probability measure on X in the space $\text{Prob}(X)$ contains a Dirac measure. The Furstenberg boundary $\partial_F G$ of G is the universal minimal and strongly proximal G -space [Fur73, Gla76]. We denote by $\text{Rad}(G)$ the amenable radical of the group G . It coincides with the kernel of the action of G on $\partial_F G$ (a result that holds more generally for arbitrary locally compact groups [Fur03]).

Definition 5.2. The stabilizer URS associated to the G -action on $\partial_F G$ is denoted \mathcal{A}_G and is called the Furstenberg URS of G .

A result of Frolík implies that the map $x \mapsto G_x$ is continuous on $\partial_F G$, so that \mathcal{A}_G is exactly the collection of point stabilizers for the action of G on $\partial_F G$ (see [Ken20] and references therein).

PROPOSITION 5.3. *The following hold:*

- (1) \mathcal{A}_G is amenable and $\mathcal{X} \preccurlyeq \mathcal{A}_G$ for every non-empty closed G -invariant subset \mathcal{X} of $\text{Sub}(G)$ consisting of amenable subgroups;
- (2) \mathcal{A}_G is invariant under the action of $\text{Aut}(G)$ on $\text{Sub}(G)$;
- (3) $\text{Rad}(G) \leq H$ for every $H \in \mathcal{A}_G$;
- (4) if N is an amenable normal subgroup of G and if $\text{Sub}_{\geq N}(G)$ is the set of subgroups of G containing N , then the natural map $\varphi : \text{Sub}(G/N) \rightarrow \text{Sub}_{\geq N}(G)$ induces a G -equivariant homeomorphism between $\mathcal{A}_{G/N}$ and \mathcal{A}_G ;
- (5) \mathcal{A}_G is a singleton if and only if $\mathcal{A}_G = \{\text{Rad}(G)\}$. When this does not hold, \mathcal{A}_G is continuous;
- (6) $\mathcal{A}_G = \{\text{Rad}(G)\}$ if and only if every amenable URS of G lives inside the amenable radical: $H \leq \text{Rad}(G)$ for every amenable URS \mathcal{H} and every $H \in \mathcal{H}$.

Proof. See [LMB18] and references therein. □

LEMMA 5.4. *Let N be a normal subgroup of G such that $H \leq N$ for every $H \in \mathcal{A}_G$. Then, $\mathcal{A}_N = \mathcal{A}_G$. In particular, N acts minimally on \mathcal{A}_G .*

Proof. \mathcal{A}_N being $\text{Aut}(N)$ -invariant, it is G -invariant. Hence, \mathcal{A}_N is an amenable URS of G . So $\mathcal{A}_N \preccurlyeq \mathcal{A}_G$. However, \mathcal{A}_G is a closed N -invariant subset of $\text{Sub}(N)$ consisting of amenable subgroups, so $\mathcal{A}_G \preccurlyeq \mathcal{A}_N$. Since \preccurlyeq is an order in restriction to URSs, $\mathcal{A}_G = \mathcal{A}_N$. □

Definition 5.5. We denote by \mathcal{F} the class of groups whose Furstenberg URS is a singleton.

LEMMA 5.6. *Suppose $G = \bigcup_I G_i$ is the directed union of subgroups G_i such that eventually, G_i is in \mathcal{F} (respectively \mathcal{A}_{G_i} is trivial). Then, G is in \mathcal{F} (respectively \mathcal{A}_G is trivial).*

Proof. Write R_i for the amenable radical of G_i , so that eventually, $\mathcal{A}_{G_i} = \{R_i\}$. Consider $\varphi_i : \text{Sub}(G) \rightarrow \text{Sub}(G_i)$, $H \mapsto H \cap G_i$. This map is continuous and G_i -equivariant. Take $H \in \mathcal{A}_G$. The subset $\mathcal{X}_i := \overline{(H \cap G_i)^{G_i}}$ is a closed G_i -invariant subset of $\text{Sub}(G_i)$ consisting of amenable subgroups, so $\mathcal{X}_i \preccurlyeq \mathcal{A}_{G_i} = \{R_i\}$ by Proposition 5.3. Since $\varphi_i(\mathcal{A}_G)$ is closed and G_i -invariant, $\mathcal{X}_i \subseteq \varphi_i(\mathcal{A}_G)$. We infer that there exists $K_i \in \mathcal{A}_G$ such that $K_i \cap G_i \leq R_i$. Upon passing to a subnet, we may assume that (K_i) converges to some $K \in \mathcal{A}_G$ and (R_i) converges to some R . The subgroup R is normal and amenable, so $R \leq \text{Rad}(G)$. Since $G = \bigcup_I G_i$, $(K_i \cap G_i)$ also converges to K , and the inclusion $K_i \cap G_i \leq R_i$ then implies $K \leq R$. So, $\mathcal{A}_G \preccurlyeq \{\text{Rad}(G)\}$, which means that $\mathcal{A}_G = \{\text{Rad}(G)\}$ by Proposition 5.3. We also immediately obtain that in the case where R_i is trivial eventually, then \mathcal{A}_G is trivial. □

5.1. *Proofs of Proposition 1.6 and Theorem 1.4.* Recall that if \mathcal{C} is a class of groups, we denote by $\mathcal{N}_G(\mathcal{C})$ the normal subgroups of G such that $G/N \in \mathcal{C}$. In the following, we mainly use this notation with $\mathcal{C} = \mathcal{F}$. We have the following lemma.

LEMMA 5.7. $\mathcal{N}_G(\mathcal{F})$ is stable under taking finite intersections.

Proof. Let $N_1, N_2 \in \mathcal{N}_G(\mathcal{F})$. Let $Q_i = G/N_i$ and let $R_i = \text{Rad}(Q_i)$ be the amenable radical of Q_i . Let $\pi_i : G \rightarrow Q_i$ be the canonical projection and $M_i := \pi_i^{-1}(R_i)$. Let also $X_i = \partial_F Q_i$. The subgroup R_i acts trivially on X_i and the assumption that Q_i belongs to \mathcal{F} means that the Q_i/R_i -action on X_i is free.

We consider the G -action on the product $X_1 \times X_2$. This action remains strongly proximal [Gla76, III.1]. It follows that there exists a unique minimal closed G -invariant subset $X \subseteq X_1 \times X_2$ and the G -action on X is strongly proximal [Gla76, III.1]. The subgroup $M_1 \cap M_2$ of G acts trivially on $X_1 \times X_2$ and, hence, on X . Moreover, since the Q_i/R_i -action on X_i is free, it follows that for the G -action on X , every point stabilizer is equal to $M_1 \cap M_2$. Equivalently, the G -action on X factors through a free action of $G/M_1 \cap M_2$. In particular, $G/M_1 \cap M_2$ is in \mathcal{F} . Since the group $M_1 \cap M_2/N_1 \cap N_2$ is amenable (as it embeds in the amenable group $R_1 \times R_2$) and since being in \mathcal{F} is invariant under forming an extension with amenable normal subgroup, it follows that $G/N_1 \cap N_2$ is in \mathcal{F} . \square

As a consequence of the lemma, it follows that the family of cosets gN , $g \in G$, $N \in \mathcal{N}_G(\mathcal{F})$, forms a basis for a group topology $\tau_{\mathcal{N}_G(\mathcal{F})}$ on G , and that the closure of H with respect to this topology is equal to $\text{cl}_{\mathcal{N}_G(\mathcal{F})}(H)$ (Proposition 2.8).

The proof of the following is the technical part of this section.

PROPOSITION 5.8. Let G be a countable group. Then, for every $N \in \mathcal{N}_G(\mathcal{F})$, there exists a normal subgroup M of G such that $N \leq M$ and $M/N \leq \text{Rad}(G/N)$, and a comeagre subset $\mathcal{H}_0 \subseteq \mathcal{A}_G$ such that $NH = M$ for every $H \in \mathcal{H}_0$.

Proof. The map $\varphi_N : \mathcal{A}_G \rightarrow \text{Sub}(G)$, $H \mapsto NH$, is lower semi-continuous by Lemma 2.6. Hence, the set $\mathcal{H}_0 \subseteq \mathcal{A}_G$ of points where φ_N is continuous is a comeagre subset of \mathcal{A}_G , and

$$E_{\varphi_N} = \overline{\{(H, NH) : H \in \mathcal{H}_0\}} \quad \text{and} \quad S_{\varphi_N} = \overline{\{NH : H \in \mathcal{H}_0\}}$$

satisfy the conclusion of Proposition 2.1. Note that S_{φ_N} is contained in the closed subset $\text{Sub}_{\geq N}(G)$ of $\text{Sub}(G)$ consisting of subgroups of G containing N .

Since the URS \mathcal{A}_G is strongly proximal and strong proximality passes to highly proximal extensions and factors [Gla75, Lemma 5.2], we deduce that the G -action on S_{φ_N} is minimal and strongly proximal. The map $\pi_N : S_{\varphi_N} \rightarrow \text{Sub}(G/N)$, $K \mapsto K/N$, is a G -equivariant homeomorphism onto its image (indeed, one easily verifies that modding out by N defines a homeomorphism from $\text{Sub}_{\geq N}(G)$ onto $\text{Sub}(G/N)$). Hence, $\pi_N(S_{\varphi_N})$ is a strongly proximal URS of G/N . Moreover, $\pi_N(S_{\varphi_N})$ consists of amenable subgroups. If we let R be the amenable radical of G/N , it follows from the assumption that G/N belongs to \mathcal{F} that $\pi_N(S_{\varphi_N})$ is contained in $\text{Sub}(R)$. However, R must act trivially on $\pi_N(S_{\varphi_N})$ by minimality and strong proximality.

Consider the envelope $E = \text{Env}(S_{\varphi_N})$. Since $\pi_N(S_{\varphi_N}) \subseteq \text{Sub}(R)$, we have $E/N \leq R$. So by the previous paragraph and the fact that $\pi_N : S_{\varphi_N} \rightarrow \pi_N(S_{\varphi_N})$ is a G -equivariant homeomorphism, it follows that E acts trivially on S_{φ_N} . However, E contains H for every $H \in \mathcal{H}_0$ and since \mathcal{H}_0 is dense in \mathcal{A}_G , this easily implies that E contains H for every $H \in \mathcal{A}_G$. Hence, Lemma 5.4 can be applied to E and we infer that E acts minimally on \mathcal{A}_G . Using Proposition 2.1 and the fact that minimality passes to irreducible extensions, we deduce that E acts minimally on S_{φ_N} . All together, this shows that S_{φ_N} is a one-point space. The corresponding normal subgroup M of G verifies the conclusion. \square

We deduce Proposition 1.6 from the introduction, which we state again for the reader's convenience.

PROPOSITION 5.9. *Let G be a countable group and let \mathcal{N} be a countable subset of $\mathcal{N}_G(\mathcal{F})$. Then, there exists a normal subgroup M of G and a comeagre subset $\mathcal{H}_0 \subseteq \mathcal{A}_G$ such that $\text{cl}_{\mathcal{N}}(H) = M$ for every $H \in \mathcal{H}_0$.*

Proof. We apply Proposition 5.8 for every $N \in \mathcal{N}$. We obtain a normal subgroup M_N of G and a comeagre subset \mathcal{H}_N of \mathcal{A}_G . Set $\mathcal{H}_0 = \bigcap_{\mathcal{N}} \mathcal{H}_N$ and $M = \bigcap_{\mathcal{N}} M_N$. Since \mathcal{N} is countable, \mathcal{H}_0 is a comeagre subset of \mathcal{H} . By construction, for every $H \in \mathcal{H}_0$,

$$\text{cl}_{\mathcal{N}}(H) = \bigcap_{\mathcal{N}} NH = \bigcap_{\mathcal{N}} M_N = M. \quad \square$$

Definition 5.10. We say that a set of laws \mathbb{W} detects amenability in a group G if for every subgroup H of G , one has that H is amenable if and only if there exists $w \in \mathbb{W}$ such that H virtually satisfies w .

Note that if a set of laws detects amenability in a group G , it also detects amenability in any subgroup of G .

Recall that a group G satisfies the Tits alternative if every subgroup of G is either virtually solvable or contains a non-abelian free subgroup. Beyond the original result of Tits about linear groups [Tit72], many classes of groups are known to satisfy the Tits alternative. Examples include hyperbolic groups [Gro87], the outer-automorphism group of a finitely generated free group [BFH00, BFH05] and of many free products [Hor14], groups acting properly on a finite-dimensional CAT(0) cube complex and with finite subgroups of bounded order [SW05], the group of polynomial automorphisms of \mathbb{C}^2 [Lam01], and the Cremona group of birational transformations of the projective space $\mathbb{P}^2(\mathbb{C})$ [Can11, Ure21].

PROPOSITION 5.11. *If a group G satisfies the Tits alternative, then there is a set of laws that detects amenability in G .*

Proof. For $\ell \geq 1$, let w_ℓ be the law from Example 2.9. So a group is solvable of length at most ℓ if and only if it satisfies w_ℓ . Let H be a subgroup of G . Since G satisfies the Tits alternative, H is amenable if and only if H is virtually solvable, if and only if there is ℓ such that H virtually satisfies w_ℓ . So, $\mathbb{W} = \{w_\ell\}_{\ell \geq 1}$ detects amenability in G . \square

The following is an immediate consequence of the definition and Lemma 2.10.

LEMMA 5.12. *Let G be a group such that there is a set of laws that detects amenability in G and let (G, τ) be a group topology on G that is Hausdorff. Then, for every amenable subgroup H of G , the τ -closure of H remains amenable.*

We note that in view of Proposition 5.11, the following result covers Theorem 1.4 from the introduction. Additionally, this result applies to all the groups (and to all their subgroups) satisfying the Tits alternative mentioned above.

THEOREM 5.13. *Let G be a group such that there is a set of laws that detects amenability in G . Then, G is in \mathcal{F} if and only if G is residually- \mathcal{F} .*

Proof. Only one direction is non-trivial. Writing G as the directed limit of its countable subgroups and invoking Lemma 5.6, one sees that it suffices to prove the result when G is countable. Under this assumption, since G is residually- \mathcal{F} , one can find $\mathcal{N} \subseteq \mathcal{N}_G(\mathcal{F})$ such that \mathcal{N} is countable and $\bigcap_{N \in \mathcal{N}} N = \{1\}$. Since $\mathcal{N}_G(\mathcal{F})$ is stable under taking finite intersections by Lemma 5.7, we can replace \mathcal{N} by the collection of finite intersections of elements of \mathcal{N} , so that we may assume that \mathcal{N} is filtering. So for a subgroup H of G , $\text{cl}_{\mathcal{N}}(H)$ equals the closure of H in the topology $\tau_{\mathcal{N}}$ (Proposition 2.8).

Proposition 5.9 provides a normal subgroup M of G such that $\text{cl}_{\mathcal{N}}(H) = M$ for every H in a comeagre subset of \mathcal{A}_G . The topology $\tau_{\mathcal{N}}$ is Hausdorff since $\bigcap_{N \in \mathcal{N}} N = \{1\}$, so it follows from Lemma 5.12 that the closure of an amenable subgroup of G remains amenable. This shows M is amenable and it follows that $M \leq \text{Rad}(G)$. By Proposition 5.3, this means that $\mathcal{A}_G = \{\text{Rad}(G)\}$. \square

Remark 5.14. When the group G is residually finite, there is a shorter way to obtain the conclusion of Theorem 5.13. Indeed, since the G -space $\partial_F G$ is proximal, it is hereditarily minimal [Gla76, Lemma 3.2]. Moreover, it follows from the conclusion of Proposition 2.1 that being hereditarily minimal is inherited from a G -space to its stabilizer URS. Hence, \mathcal{A}_G is a hereditarily minimal URS. Hence, Proposition 4.11 and Theorem 4.12 apply, and the conclusion follows as above.

COROLLARY 5.15. *Let G be a group such that there is a set of laws that detects amenability in G and suppose $\text{Rad}(G)$ is trivial. If G is residually- \mathcal{F} , then G is C^* -simple.*

Proof. The result follows from Theorem 5.13 and the main result of [KK17], which asserts that G is in \mathcal{F} if and only if $G/\text{Rad}(G)$ is C^* -simple. \square

5.2. *Linear groups.* We deduce Corollary 1.5 from the introduction, which asserts that linear groups belong to \mathcal{F} .

Proof of Corollary 1.5. Writing G as the directed limit of its finitely generated subgroups and invoking Lemma 5.6, one sees that without loss of generality, we can assume that G is a finitely generated linear group. By Malcev's theorem, the group G is residually finite. Also by the Tits alternative [Tit72], every amenable subgroup of G is virtually solvable (we are using again that G is finitely generated to have this version of the Tits alternative). Hence, all the assumptions of Theorem 5.13 are verified. The conclusion follows. \square

Acknowledgements. Thanks are due to Uri Bader and Pierre-Emmanuel Caprace. We can trace back that the possibility of using Proposition 2.1 specifically in the space of subgroups to build a URS starting from another one and a semi-continuous map had been originally brought to our attention by them several years ago. We also thank a referee who provided various comments that improved the exposition of the paper. This work had been initiated within the framework of the Labex Milyon (ANR-10-LABX-0070) of Université de Lyon, within the program ‘Investissements d’Avenir’ (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

REFERENCES

- [AG77] J. Auslander and S. Glasner. Distal and highly proximal extensions of minimal flows. *Indiana Univ. Math. J.* **26**(4) (1977), 731–749.
- [BFH00] M. Bestvina, M. Feighn and M. Handel. The Tits alternative for $\text{Out}(F_n)$. I. Dynamics of exponentially-growing automorphisms. *Ann. of Math. (2)* **151**(2) (2000), 517–623.
- [BFH05] M. Bestvina, M. Feighn and M. Handel. The Tits alternative for $\text{Out}(F_n)$. II. A Kolchin type theorem. *Ann. of Math. (2)* **161**(1) (2005), 1–59.
- [BH21] R. Boutonnet and C. Houdayer. Stationary characters on lattices of semisimple Lie groups. *Publ. Math. Inst. Hautes Études Sci.* **133** (2021), 1–46.
- [BK KO17] E. Breuillard, M. Kalantar, M. Kennedy and N. Ozawa. C^* -simplicity and the unique trace property for discrete groups. *Publ. Math. Inst. Hautes Études Sci.* **126** (2017), 35–71.
- [Bou71] N. Bourbaki. *Éléments de mathématique. Topologie générale. Chapitres 1 à 4*. Hermann, Paris, 1971.
- [BS06] U. Bader and Y. Shalom. Factor and normal subgroup theorems for lattices in products of groups. *Invent. Math.* **163**(2) (2006), 415–454.
- [Can11] S. Cantat. Sur les groupes de transformations birationnelles des surfaces. *Ann. of Math. (2)* **174**(1) (2011), 299–340.
- [Cor06] Y. Cornuier. Finitely presented wreath products and double coset decompositions. *Geom. Dedicata* **122** (2006), 89–108.
- [dlH07] P. de la Harpe. On simplicity of reduced C^* -algebras of groups. *Bull. Lond. Math. Soc.* **39**(1) (2007), 1–26.
- [EJZ13] M. Ershov and A. Jaikin-Zapirain. Groups of positive weighted deficiency and their applications. *J. Reine Angew. Math.* **677** (2013), 71–134.
- [Ele18] G. Elek. Uniformly recurrent subgroups and simple C^* -algebras. *J. Funct. Anal.* **274**(6) (2018), 1657–1689.
- [FG23] M. Fraczyk and T. Gelander. Infinite volume and infinite injectivity radius. *Ann. of Math. (2)* **197**(1) (2023), 389–421.
- [Fur73] H. Furstenberg. Boundary theory and stochastic processes on homogeneous spaces. *Harmonic Analysis on Homogeneous Spaces (Proc. Sympos. Pure Math., Vol. XXVI, Williams Coll., Williamstown, Mass., 1972) (Proceedings of Symposia in Pure Mathematics, XXVI)*. Ed. C. C. Moore. American Mathematical Society, Providence, RI, 1973, pp. 193–229.
- [Fur03] A. Furman. On minimal strongly proximal actions of locally compact groups. *Israel J. Math.* **136** (2003), 173–187.
- [Gla75] S. Glasner. Compressibility properties in topological dynamics. *Amer. J. Math.* **97** (1975), 148–171.
- [Gla76] S. Glasner. *Proximal Flows (Lecture Notes in Mathematics, 517)*. Springer-Verlag, Berlin–New York, 1976.
- [Gri11] R. I. Grigorchuk. Some problems of the dynamics of group actions on rooted trees. *Tr. Mat. Inst. Steklova* **273**(Sovremennye Problemy Matematiki) (2011), 72–191.
- [Gro87] M. Gromov. Hyperbolic groups. *Essays in Group Theory (Mathematical Sciences Research Institute Publications, 8)*. Ed. S. M. Gersten. Springer, New York, 1987, pp. 75–263.
- [GW15] E. Glasner and B. Weiss. Uniformly recurrent subgroups. *Recent Trends in Ergodic Theory and Dynamical Systems (Contemporary Mathematics, 631)*. Ed. S. Bhattacharya, T. Das, A. Ghosh and R. Shah. American Mathematical Society, Providence, RI, 2015, pp. 63–75.
- [Hal49] M. Hall. Coset representations in free groups. *Trans. Amer. Math. Soc.* **67** (1949), 421–432.
- [Hor14] C. Horbez. The Tits alternative for the automorphism group of a free product. *Invent. Math.* **3** (2025), 869–901.

- [Ken20] M. Kennedy. An intrinsic characterization of C^* -simplicity. *Ann. Sci. Éc. Norm. Supér. (4)* **53**(5) (2020), 1105–1119.
- [KK17] M. Kalantar and M. Kennedy. Boundaries of reduced C^* -algebras of discrete groups. *J. Reine Angew. Math.* **727** (2017), 247–267.
- [Kur28] C. Kuratowski. Sur les décompositions semi-continues d'espaces métriques compacts. *Fund. Math.* **11**(1) (1928), 169–185 (in French).
- [Lam01] S. Lamy. L'alternative de Tits pour. *J. Algebra* **239**(2) (2001), 413–437.
- [LB17] A. Le Boudec. C^* -simplicity and the amenable radical. *Invent. Math.* **209**(1) (2017), 159–174.
- [LBMB18] A. Le Boudec and N. Matte Bon. Subgroup dynamics and C^* -simplicity of groups of homeomorphisms. *Ann. Sci. Éc. Norm. Supér. (4)* **51**(3) (2018), 557–602.
- [LBMB22] A. Le Boudec and N. Matte Bon. Growth of actions of solvable groups. *Preprint*, 2022, [arXiv:2205.11924](https://arxiv.org/abs/2205.11924).
- [MB17] N. Matte Bon. Full groups of bounded automaton groups. *J. Fractal Geom.* **4**(4) (2017), 425–458.
- [MBT20] N. Matte Bon and T. Tsankov. Realizing uniformly recurrent subgroups. *Ergod. Th. & Dynam. Sys.* **40**(2) (2020), 478–489.
- [Nek20] V. Nekrashevych. Substitutional subshifts and growth of groups. *Preprint*, 2020, [arXiv:2008.04983](https://arxiv.org/abs/2008.04983).
- [Neu37] B. H. Neumann. Some remarks on infinite groups. *J. Lond. Math. Soc. (3)* **12** (1937), 120–127 (in English).
- [Rau20] S. Raum. C^* -simplicity [after Breuillard, Haagerup, Kalantar, Kennedy and Ozawa]. *Astérisque* **422** (2020), 225–252; Séminaire Bourbaki. Vol. 2018/2019. Exposés 1151–1165, Exp. No. 1156.
- [SW05] M. Sageev and D. T. Wise. The Tits alternative for $CAT(0)$ cubical complexes. *Bull. Lond. Math. Soc.* **37**(5) (2005), 706–710.
- [Tit72] J. Tits. Free subgroups in linear groups. *J. Algebra* **20** (1972), 250–270.
- [Ure21] C. Urech. Subgroups of elliptic elements of the Cremona group. *J. Reine Angew. Math.* **770** (2021), 27–57.
- [Wil98] J. S. Wilson. *Profinite Groups (London Mathematical Society Monographs. New Series, 19)*. The Clarendon Press, Oxford University Press, New York, 1998.