

- 21 Alemany S, Arias B, Aguilera M, Villa H, Moya J, Ibáñez MI, et al. Childhood abuse, the BDNF-Val66Met polymorphism and adult psychotic-like experiences. *Br J Psychiatry* 2011; **199**: 38–42.
- 22 National Institute of Mental Health. *Genetics. Initiative. Family Interview for Genetic Studies (FIGS)*. NIMH, 1992.
- 23 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorder (4th edn) (DSM-IV)*. APA, 1994.
- 24 Andreasen NC, Flaum M, Arndt S. The Comprehensive Assessment of Symptoms and History (CASH). An instrument for assessing diagnosis and psychopathology. *Arch Gen Psychiatry* 1992; **49**: 615–23.
- 25 Kendler KS, Lieberman JA, Walsh D. The Structured Interview 559 for Schizotypy (SIS): a preliminary report. *Schizophr Bull* 1989; **15**: 559–71.
- 26 Thomas A, Donnell AJ, Young TR. Factor structure and differential validity of the expanded Brief Psychiatric Rating Scale. *Assessment* 2004; **11**: 177–87.
- 27 van Winkel R. Family-based analysis of genetic variation underlying psychosis-inducing effects of cannabis: sibling analysis and proband follow-up. *Arch Gen Psychiatry* 2011; **68**: 148–57.
- 28 Binder EB, Bradley RG, Liu W, Epstein MP, Deveau TC, Mercer KB, et al. Association of FKBP5 polymorphisms and childhood abuse with risk of posttraumatic stress disorder symptoms in adults. *JAMA* 2008; **299**: 1291–305.
- 29 Xie P, Kranzler HR, Poling J, Stein MB, Anton RF, Farrer LA, et al. Interaction of FKBP5 with childhood adversity on risk for post-traumatic stress disorder. *Neuropsychopharmacology* 2010; **35**: 1684–92.
- 30 Velders FP, Kuningas M, Kumari M, Dekker MJ, Uitterlinden AG, Kirschbaum C, et al. Genetics of cortisol secretion and depressive symptoms: a candidate gene and genome wide association approach. *Psychoneuroendocrinology* 2011; **36**: 1053–61.
- 31 Zimmermann P, Brückl T, Nocon A, Pfister H, Binder EB, Uhr M, et al. Interaction of FKBP5 gene variants and adverse life events in predicting depression onset: results from a 10-year prospective community study. *Am J Psychiatry* 2011; **168**: 1107–16.
- 32 Roy A, Gorodetsky E, Yuan Q, Goldman D, Enoch MA. Interaction of FKBP5, a stress-related gene, with childhood trauma increases the risk for attempting suicide. *Neuropsychopharmacology* 2010; **35**: 1674–83.
- 33 Koenen KC, Uddin M. FKBP5 polymorphisms modify the effects of childhood trauma. *Neuropsychopharmacology* 2010; **35**: 1623–24.
- 34 Read J, Hammersley P, Rudegeair T. Why, when and how to ask about childhood abuse. *Adv Psychiatr Treat* 2007; **13**: 101–10.
- 35 Hartmann J, Wagner KV, Dedic N, Marinescu D, Scharf SH, Wang XD, et al. Fkbp52 heterozygosity alters behavioral, endocrine and neurogenetic parameters under basal and chronic stress conditions in mice. *Psychoneuroendocrinology* 2012; **37**: 2009–12.
- 36 McGowan PO, Sasaki A, D'Alessio AC, Dymov S, Labonté B, Szyf M, et al. Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nat Neurosci* 2009; **12**: 342–48.
- 37 Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, et al. Epigenetic programming by maternal behavior. *Nat Neurosci* 2004; **7**: 847–54.
- 38 Champagne FA, Curley JP. Epigenetic mechanisms mediating the long-term effects of maternal care on development. *Neurosci Biobehav Rev* 2009; **33**: 593–600.
- 39 Touma C, Gassen NC, Herrmann L, Cheung-Flynn J, Büll DR, Ionescu IA, et al. FK506 binding 5 shapes stress responsiveness: modulation of neuroendocrine reactivity and coping behavior. *Biol Psychiatry* 2011; **15**: 928–36.
- 40 Butts KA, Weinberg J, Young AH, Phillips AG. Glucocorticoid receptors in the prefrontal cortex regulate stress-evoked dopamine efflux and aspects of executive function. *PNAS* 2011; **108**: 18459–64.
- 41 Duncan LE, Keller MC. A critical review of the first 10 years of candidate gene-by-environment interaction research in psychiatry. *Am J Psychiatry* 2011; **168**: 1041–9.
- 42 Moffitt TE, Caspi A, Rutter M. Strategy for investigating interactions between measured genes and measured environments. *Arch Gen Psychiatry* 2005; **62**: 473–81.
- 43 Fardo DW, Becker KD, Bertram L, Tanzi RE, Lange C. Recovering unused information in genome-wide association studies: the benefit of analyzing SNPs out of Hardy–Weinberg equilibrium. *Eur J Hum Genet* 2009; **17**: 1676–82.
- 44 Hruschka DJ, Kohrt BA, Worthman CM. Estimating between- and within-individual variation in cortisol levels using multilevel models. *Psychoneuroendocrinology* 2005; **30**: 698–714.

100  
words

## Art Brut

Rachael Hall

Art Brut, translated from the French as 'raw art', is used to describe art occurring outside of the conventional art scene, typically without the process of formal training that renders art 'cooked'. The term was coined by the French artist Jean Dubuffet in the 1940s. A significant contribution to this genre, although not exclusively, is from those people with exceptional artistic abilities who also suffer from serious mental illness. Free from the constraints of formal art culture, Art Brut allows for unbridled creative expression, in turn giving the viewer a direct insight into the artist's perceptions, fantasies, hopes and fears.

The British Journal of Psychiatry (2013)  
202, 268. doi: 10.1192/bjp.bp.112.122317