

CORRIGENDUM

Effect of a cod protein hydrolysate on postprandial glucose metabolism in healthy subjects: a double-blind cross-over trial — CORRIGENDUM

Hanna Fjeldheim Dale^{1,2*}, Caroline Jensen¹, Trygve Hausken^{1,2,3}, Einar Lied⁴, Jan Gunnar Hatlebakk^{1,2,3}, Ingeborg Brønstad^{5,6}, Dag Arne Lihaug Hoff^{7,8} and Gülen Arslan Lied^{1,2,3}

¹Department of Clinical Medicine, Centre for Nutrition, University of Bergen, Bergen, Norway

²Division of Gastroenterology, Department of Medicine, Haukeland University Hospital, Bergen, Norway

³National Centre of Functional Gastrointestinal Disorders, Haukeland University Hospital, Bergen, Norway

⁴Firmenich Bjørge Biomarin AS, Ellingsøy, Ålesund, Norway

⁵Department of Clinical Medicine, University of Bergen, Bergen, Norway

⁶National Centre for Ultrasound in Gastroenterology, Haukeland University Hospital, Bergen, Norway

⁷Division of Gastroenterology, Department of Medicine, Ålesund Hospital, Møre & Romsdal Hospital Trust, Ålesund, Norway

⁸Department of Clinical and Molecular Medicine, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway

Journal of Nutritional Science (2019), vol. 8, e1, page 1 of 2

doi:10.1017/jns.2018.30

doi:10.1017/jns.2018.23, Published online by Cambridge University Press, 28 November 2018

Original text and correction:

ORIGINAL TEXT (page 3, Subjects and methods)

Fig. 1. Flowchart depicting the inclusion process for the study evaluating the effect of a marine protein hydrolysate (MPH) from Atlantic cod (*Gadus morhua*) on postprandial glucose metabolism in healthy individuals aged 40–65 years. Participants were recruited through advertisements on the Internet and posters at Haukeland University Hospital and Ålesund Hospital between October 2017 and February 2018.

Fig. 2. Study protocol for the evaluation of the effect of a marine protein hydrolysate (MPH) from Atlantic cod (*Gadus morhua*) on postprandial glucose metabolism. We included forty-one healthy subjects (age range 40–64 years).

CORRECTION

Fig. 1. Study protocol for the evaluation of the effect of a marine protein hydrolysate (MPH) from Atlantic cod (*Gadus morhua*) on postprandial glucose metabolism. We included forty-one healthy subjects (age range 40–64 years).

Fig. 2. Flowchart depicting the inclusion process for the study evaluating the effect of a marine protein hydrolysate (MPH) from Atlantic cod (*Gadus morhua*) on postprandial glucose metabolism in healthy individuals aged 40–65 years. Participants were recruited through advertisements on the Internet and posters at Haukeland University Hospital and Ålesund Hospital between October 2017 and February 2018.

* Corresponding authors: H. F. Dale, email hanna.dale@outlook.com; C. Jensen, email caroline.j@uib.no

**ORIGINAL TEXT (page 8, Acknowledgements)**

E. L. is Professor Emeritus at the University of Bergen, Bergen, Norway and the managing director of Science of Firmenich Bjørge Biomarin AS, Ellingsøy, Ålesund, Norway. The other authors declare no conflict of interest.

CORRECTION

E. L. is Professor Emeritus at the University of Bergen, Bergen, Norway, and former Scientific Advisor of Firmenich Bjørge Biomarin AS, Ellingsøy, Ålesund, Norway, where he holds a royalty agreement. The other authors declare no conflict of interest.

Reference

Dale H, Jensen C, Hausken T, *et al.* (2018) Effect of a cod protein hydrolysate on postprandial glucose metabolism in healthy subjects: a double-blind cross-over trial. *J Nutr Sci* 7, E33. doi:10.1017/jns.2018.23