

CORRIGENDUM

Comparison of the production rate of bacteria in the rumen of buffalo calves estimated by using labelled *Streptococcus bovis* and mixed ruminal bacterial cells

D. N. VERMA, U. B. SINGH, S. K. SRIVASTAVA AND R. V. N. SRIVASTAVA
J. agric. Sci., Camb. (1976), **87**, 661–664

The authors regret the following error:

Page 663, Table 3, line 1

For Green maize ³⁵S mixed bacterial cells
read Green maize ¹⁴C mixed bacterial cells



Leaf Protein and Other Aspects of Fodder Fractionation

N. W. PIRIE

Recently it has become more acceptable to consider the fractionation of leafy crops into their components instead of using them only for ruminant fodder. In suitable climates, fractionation of leafy crops can yield more edible protein than other forms of agriculture. The fibre residue which is unfit for human consumption is still useful feed for ruminants, and the soluble leaf components can be used in various ways. The author himself has been actively involved in the development of fodder fractionation techniques, and is therefore uniquely placed to describe the methods of separating leaf protein from other leaf components, the practical problems encountered, and how they can be overcome. The production of leaf protein in bulk is shown to be efficient, technically feasible and nutritionally valuable, making fodder fractionation an important agricultural technique for technically unadvanced countries where protein deficiency is most often acute. This interesting and informative account will be of interest to agricultural research workers and students, as well as to those concerned with nutrition and the developing world.

£8.00 net

First Paperback Edition

Crop Physiology

Edited by L. T. EVANS

This book contains comprehensive reviews of the physiological basis of yield in nine of the world's most important crops. In the introductory chapter the contribution of these crops to the world food supply is assessed together with a consideration of aspects of crop evolution and an outline of the history of crop physiology. The following nine chapters on the individual crops are written by international authorities. The final chapter deals with the physiological basis of yield as a foundation for future progress in agronomic techniques and plant breeding for yield. This text will be invaluable to plant breeders, crop physiologists and agronomists, to advanced students in the same fields and others concerned with the problems of yield in plants.

'The timeliness of the book makes it an excellent resource for advanced undergraduate and graduate courses and a valuable reference for crop investigators.'

American Scientist

'The book is extremely readable and well presented and can be recommended not only to crop physiologists but also to agronomists and plant breeders.'

Nature

Also issued in hard covers

£5.50 net

CAMBRIDGE UNIVERSITY PRESS

	PAGE
MAXWELL, T. J., DONEY, J. M., MILNE, J. A., PEART, J. N., RUSSEL, A. J. F., SIBBALD, A. R. and MACDONALD, D. The effect of rearing type and prepartum nutrition on the intake and performance of lactating Greyface ewes at pasture	165
KEATINGE, J. D. H., STEWART, R. H. and GARRETT, M. K. The influence of temperature and soil water potential on the leaf extension rate of perennial ryegrass in Northern Ireland	175
ØRSKOV, E. R., STEWART, C. S. and GREENHALGH, J. F. D. The effect of sodium hydroxide and urea on some storage properties of moist grain	185
PREW, R. D. and DYKE, G. V. Experiments comparing 'break crops' as a preparation for winter wheat followed by spring barley	189
FREE, J. B. and WILLIAMS, INGRID H. The infestation of crops of oil-seed rape (<i>Brassica napus</i> L.) by insect pests	203
VAN KAMPEN, M., MITCHELL, B. W. and SIEGEL, H. S. Thermoneutral zone of chickens as determined by measuring heat production, respiration rate, and electromyographic and electroencephalographic activity in light and dark environments and changing ambient temperatures	219
WURR, D. C. E. The effect of the site of seed potato production on subsequent sprout growth and tuber yield	227
LAMB, C. S. and EADIE, J. The effect of barley supplements on the voluntary intake and digestion of low quality roughages by sheep	235
TOLAND, P. C. Loss of starch and fibre of whole grain in nylon bags suspended in the rumen of steers	243
BOLAND, M. P. and GORDON, I. Effect of timing of ram introduction on fertility in progestagen-PMSG treated anoestrous ewes	247
PAGE, E. R. Aqueous ammonia as a nitrogen fertilizer for summer cauliflowers, compared with ammonium nitrate (broadcast) and urea (broadcast and injected)	251
PHILLIPS, M. S., WILSON, LINDA A. and FORREST, J. M. S. General and specific combining ability of potato parents for resistance to the white potato cyst nematode (<i>Globodera pallida</i>)	255

CONTENTS

Vol. 92 Part 1 February 1979

	PAGE
MITTAL, J. P. and GHOSH, P. K. Comparative semen characteristics of Corriedale, Marwari and Jaisalmeri rams maintained under hot arid conditions	1
BELL, F. R. and SLY, JENNIFER. Sodium appetite and sodium metabolism in ruminants assessed by preference tests	5
FRENCH, B. K. and LEGG, B. J. Rothamsted irrigation 1964-76.	15
SMITH, M. W. and JAMES, P. S. Absorption of methionine by foetal and neonatal lamb colon	39
PRITCHARD, D. T. and BROWN, N. J. Respiration in cropped and fallow soil	45
WILLIAMS, INGRID H. and FREE, J. B. Compensation of oil-seed rape (<i>Brassica napus</i> L.) plants after damage to their buds and pods	53
WILKINS, CAROLYN. The distribution of Mn, Fe, Cu, and Zn in topsoils and herbage of North West Pembrokeshire	61
FOOT, J. Z., SKEDD, E. and MCFARLANE, D. N. Body composition in lactating sheep and its indirect measurement in the live animal using tritiated water	69
JONES, H. G. Visual estimation of plant water status in cereals	83
JONES, R. J., LUDLOW, M. M., TROUGHTON, J. H. and BLUNT, C. G. Estimation of the proportion of C ₃ and C ₄ plant species in the diet of animals from the ratio of natural ¹² C and ¹³ C isotopes in the faeces	91
WALKEY, D. G. A. and INNES, N. L. Resistance to bean common mosaic virus in dwarf beans (<i>Phaseolus vulgaris</i> L.)	101
BARDNER, R. and FLETCHER, K. E. Larvae of the pea and bean weevil, <i>Sitona lineatus</i> , and the yield of field beans	109
OSUJI, P. O. and DEVERS, C. Nitrogen utilization of growing-finishing Barbados Blackbelly lambs	113
DONEY, J. M., PEART, J. N., SMITH, W. F. and LOUDA, F. A consideration of the techniques for estimation of milk yield by suckled sheep and a comparison of estimates obtained by two methods in relation to the effect of breed, level of production and stage of lactation .	123
PEART, J. N., DONEY, J. M. and SMITH, W. F. Lactation pattern in Scottish Blackface and East Friesland × Scottish Blackface cross-bred ewes	133
FREE, J. B. and WILLIAMS, INGRID H. The distribution of insect pests on crops of oil-seed rape (<i>Brassica napus</i> L.) and the damage they cause	139
ALLEN, E. J., BEAN, J. N., GRIFFITH, R. L. and O'BRIEN, P. J. Effects of length of sprouting period on growth and yield of contrasting early potato varieties	151

Continued on inside back cover

© Cambridge University Press 1979

SUBSCRIPTIONS. Two volumes of three parts are published annually. The subscription price of volumes 92 and 93, 1979, is £27.00 net (U.S.A. and Canada US \$66.50) per volume (post free); single parts are available at £11.00 net (U.S.A. and Canada US \$27.50) plus postage. Orders or enquiries may be sent to any bookseller or subscription agent, or to Cambridge University Press, P.O. Box 110, Cambridge CB2 3RL (U.S.A. and Canada, Cambridge University Press, 32 East 57th Street, New York, N.Y. 10022, U.S.A.).

Printed in Great Britain at the University Press, Cambridge