

RADIOCARBON VARIATIONS FROM TASMANIAN CONIFERS: FIRST RESULTS FROM LATE PLEISTOCENE AND HOLOCENE LOGS

MIKE BARBETTI¹, TREVOR BIRD², GEORGE DOLEZAL¹, GILLIAN TAYLOR¹
ROGER FRANCEY³, EDWARD COOK⁴ and MIKE PETERSON⁵

ABSTRACT. Dendrochronological studies have begun on two conifer species in the Stanley River area of western Tasmania. The chronology extends to 273 BC for Huon pine (*Lagarostrobos franklinii*) and to AD 1450 for celery-top pine (*Phyllocladus aspleniifolius*). Apart from living or recently felled trees, sections have been taken from 58 logs preserved in floodplain sediments. Two of these logs have late Pleistocene ages, centered around 13.0 and 12.7 k ¹⁴C yr BP. Four logs are between 8 and 9 ka BP, and one is centered at 7.3 ka BP. The remaining logs have various ages between 6.2 ka BP and the present. ¹⁴C measurements have been performed on decadal samples from the two late Pleistocene logs, providing short (260-yr) records of atmospheric ¹⁴C variations when plotted against individual ring numbers. Decadal measurements on the 7300-yr-old log have been wiggle-matched with ¹⁴C calibration curves from German oak and bristlecone pine. Measurements for the period, AD 1600–1800, show good agreement with northern hemisphere results, and a nearly zero offset between the hemispheres.

INTRODUCTION

Radiocarbon calibration is now well established for most of the Holocene. The differences between tree-ring ages and conventional ¹⁴C ages have been determined for the last 9700 calendar yr by precision ¹⁴C measurements on 10- or 20-ring samples, which are independently dated by dendrochronology (summarized by Stuiver *et al.* 1991). A small part of the difference occurs because ¹⁴C ages are, by international agreement, calculated using a half-life of 5568 yr, which is known to be about 3% too short. Differences apart from this 3% reflect variations in the production rate and in the exchange of ¹⁴C between oceans, atmosphere and biosphere. Most of the Holocene variation is thought to be due to changes in the production rate; the long-term peak-to-trough change is attributed to changes in the Earth's magnetic field strength, which affect the cosmic-ray flux, whereas shorter-term wiggles (with amplitudes of 100 or 200 yr) are attributed to solar modulation of the cosmic-ray flux.

Early Holocene ¹⁴C data from southern Germany (Becker & Kromer 1986) are from a floating tree-ring sequence. They show short-term wiggles, such as those seen in recent millennia, and also a horizontal trend in the calibration curve with nearly constant ¹⁴C ages (9600 BP) over several hundred tree rings.

Southern hemisphere measurements are essential as an independent verification, and are important because the offset from the northern hemisphere (36 yr in recent centuries; Lerman, Mook & Vogel 1970; Vogel *et al.* 1986) may have varied in earlier times due to changes in global carbon fluxes.

We began dendrochronological studies 10 years ago on two conifer species in the Stanley River area of western Tasmania (145°E, 42°S; Francey *et al.* 1984). The chronology for Huon pine (*Lagarostrobos franklinii*) has recently been extended to 273 BC. Living celery-top pine (*Phyllocladus aspleniifolius*) trees are up to 500 yr old. We sampled logs exposed in the river banks and

¹The NWG Macintosh Centre for Quaternary Dating, Madsen Building F09, University of Sydney, Sydney NSW 2006 Australia

²Trades Hall, 219 New Town Road, New Town, Tasmania 7008 Australia

³CSIRO Division of Atmospheric Research, Private Bag No. 1, Mordialloc, Victoria 3195 Australia

⁴Lamont-Doherty Geological Observatory of Columbia University, Palisades, New York 10964 USA

⁵Forestry Commission, Hobart, Tasmania 7001 Australia

excavated in floodplain sediments between 1981 and 1984, and obtained sections from 58 well-preserved logs. The oldest are two celery-top logs with late Pleistocene ages, centered around 13.0 and 12.7 k ¹⁴C yr BP. Four logs are between 8 and 9 ka, and one is centered at 7.3 ka BP. The remaining logs have various ages between 6.2 ka BP and the present, but so far there are no logs between about 3.6 and 2.6 ka.

These discoveries in western Tasmania have given us an opportunity to make selected southern hemisphere comparisons with the 9.7 ka (dendroyear) span of ¹⁴C records obtained from northern hemisphere bristlecone pine and European oak (Stuiver *et al.* 1991).

METHODOLOGY

We polished cross-sections from the logs and split them into consecutive 10-ring samples; for the innermost and outermost parts, where the rings were narrow, we took up to 60 rings per sample. For the time span, AD 1600–1800, we used five-ring samples where possible. We reduced the wood samples to about 0.5 mm particle size in a cutting mill, and prepared holocellulose following the method of Head (1979). We used standard techniques (Gupta & Polach 1985) to prepare benzene samples (4 ml). Stable carbon isotope (¹³C/¹²C) measurements were made on subsamples of the combustion CO₂ at the CSIRO Division of Atmospheric Research.

We made ¹⁴C measurements using Teflon vials in a low-level Wallac Quantulus counter, usually with duplicate measurements in another counter (Wallac Rackbeta) linked to a multichannel analyzer and microcomputer system. A third counter (Packard Tri-Carb with fixed windows) was used for some measurements before February 1987, but most of the data were collected between 1987 and 1989. We calculated conventional ¹⁴C ages using modern standard values derived from measurements of ANU sucrose and NBS oxalic acid, but we assumed an uncertainty of ± 0.1 or ± 0.2 counts min⁻¹ for the standard, considerably larger than the typical Poisson deviation of ± 0.07 counts min⁻¹ associated with an individual standard measurement. We then combined the results from the counters, weighting them inversely by variance; age differences between the counters are indicated by z-statistic values. Measurement uncertainties associated with our combined results typically range from *ca.* ± 30 yr for recent samples to *ca.* ± 60 yr for late Pleistocene samples.

RESULTS

¹⁴C results (Table 1) from the two late Pleistocene logs provide short (260-yr) records of atmospheric ¹⁴C variations when plotted against individual ring numbers (Fig. 1). The variations from sample to sample are not always smooth, and may reflect a combination of experimental uncertainties (trace levels of contamination and random measurement errors). One ¹⁴C result from the center of SRT-462, and two from the outermost rings of SRT-157 (probably sapwood), gave anomalously young results, which are excluded from Figure 1. The overall trend for SRT-462 is fairly flat, and the trend for SRT-157 is downward toward the outer rings. The two logs do not appear to overlap in time.

Results from the 7300-yr-old log SRT-444 (Table 2) can be compared with ¹⁴C calibration curves from German oak and bristlecone pine (Fig. 2). Since SRT-444 is not dated by dendrochronology, we sought to position it by ¹⁴C wiggle-matching. We estimated decadal values for German oak by a combination of averaging (for cases of multiple measurements per decade) and interpolation (for decades without data). Three of the SRT-444 values that did not correspond exactly to decadal spacing were taken as representing the nearest decadal sample. We used these smoothed values and calculated root mean square (rms) differences, moving the zero-point of the SRT-444 data in decadal steps. The minimum rms difference occurred with ring 0 of SRT-444 placed at 8260 den-

TABLE 1. Radiocarbon Dates From Stanley River Celery-Top Pine Logs, SRT-462 and SRT-157

SUA- no.	Tree-ring span years	¹³ C (‰)	Counter 1 (yr BP)	± σ	Counter 2 (yr BP)	± σ	Counter 3 (yr BP)	± σ	Combined (yr BP)	± σ	z-stat	Center ring
<i>SRT-462*</i>												
5142R**	1	-26.5	13,134	53					13,134	53		26
5143†	51	-24.0	12,882	68					12,882	68		63
5144†	76	-23.7	13,105	69	13,165	105			13,123	58	.48	83
5145	91	-23.7	13,149	69	13,121	104			13,140	57	-.22	96
5101**	70	-24.1	13,032	68	12,971	103			13,013	57	-.49	100
5146	101	-23.7	13,039	68	12,837	102			12,977	57	-1.65	106
5147	111	-24.0	13,065	68	13,301	97			13,143	56	1.99	116
5148	121	-23.4	13,174	53					13,174	53		126
5149	131	-23.3	12,852	67	13,018	103			12,901	56	1.35	136
5150	141	-23.4	13,012	53					13,012	53		146
5151	151	-24.2	12,764	67					12,764	67		156
5152	161	-24.3	13,072	69	13,036	104			13,061	57	-.29	166
5153	171	-24.9	13,008	53					13,008	53		176
5154	181	-25.1	12,825	67	12,939	103			12,859	56	.93	186
5155	191	-25.3	12,946	68	12,843	124			12,922	60	-.73	196
5156	201	-24.7	12,906	52					12,906	52		206
5157	211	-24.4	12,947	68	13,054	104			12,979	57	.86	216
5158†	221	-23.5	12,992	68					12,992	68		239
5142 [§]	1	-26.8	12,195	65	12,227	98						
Averages												
										13,000	.22	141

TABLE 1. (Continued)

SUA- no.	Tree-ring span years	¹³ C (‰)	Counter 1 (yr BP)	± σ	Counter 2 (yr BP)	± σ	Counter 3 (yr BP)	± σ	Combined (yr BP)	± σ	z-stat	Center ring
<i>SRT-157*</i>												
5022**	156 205	-23.1					12,390	80				181
5028†	86 110	-26.7					12,870	90				98
5104**	1 60	-25.7	12,747	67	12,821	102			12,769	56	.61	31
5105†	61 85	-25.6	12,734	67					12,734	67		73
5106†	86 110	-25.6	12,940	68	12,815	102			12,902	57	-1.02	98
5107	111 120	-24.4	12,685	67	12,569	92			12,645	54	-1.02	116
5108	121 130	-24.6	12,682	67	12,862	124			12,723	59	1.28	126
5109	131 140	-24.7	12,600	66					12,600	66		136
5110	141 150	-24.7	12,779	67					12,779	67		146
5042	151 160	-23.6	12,593	99					12,593	99		156
5043	161 170	-22.0					12,667	118	12,667	118		166
5044	171 180	-22.7	12,792	83			12,726	113	12,769	67	-47	176
5054	181 190	-23.3	12,692	67	12,507	98			12,633	55	-1.56	186
5136	191 200	-24.9	12,499	66	12,373	90			12,455	53	-1.13	196
5137	200 210	-25.0	12,543	66	12,464	99			12,519	55	-0.66	205
5138	211 220	-25.7	12,507	66	12,622	100			12,542	55	.96	216
5139	221 235	-24.8	12,500	66	12,404	99			12,470	55	-0.81	228
5111‡	236 260	-25.4	11,974	64								
5175‡	236 260	-24.2	10,861	59								
Averages											-38	149

*10-ring samples unless otherwise noted

**50- or 60-ring sample

†25- or 36-ring sample

‡15-ring sample

§rejected ¹⁴C datum

TABLE 2. Radiocarbon Dates From Stanley River Celery-top Pine Log, SRT-444*

SUA- no.	Tree rings span years	¹³ C (‰)	Counter 1		Counter 2		Combined		z-stat	Center ring	Dendro yr BP
			yr BP	± σ	yr BP	± σ	yr BP	± σ			
5088**	307 350	-23.0	7178	48	7115	70	7158	40	-.74	329	7932
5112	301 310	-24.2	7200	48			7200	48		306	7955
5113	291 300	-24.3	7242	48	7189	83	7229	42	-.55	296	7965
5114	281 290	-24.1	7170	48	7186	70	7175	40	.19	286	7975
5115	271 280	-24.3	7130	48	7077	70	7113	40	-.62	276	7985
5116R	261 270	-24.3	7146	48	7154	70	7149	40	.09	266	7995
5117	251 260	-24.6	7003	48	7076	70	7026	40	.86	256	8005
5118	241 250	-24.5	7148	48	7247	71	7179	40	1.16	246	8015
5119	231 240	-24.8	7226	48	7087	70	7182	40	-1.64	236	8025
5120	221 230	-24.7	7221	48	7371	71	7268	40	1.75	226	8035
5121	211 220	-24.4	7253	48	7270	66	7259	39	.21	216	8045
5122	201 210	-24.9	7306	48	7385	66	7333	39	.97	206	8055
5123	191 200	-24.5	7340	48	7336	84	7339	42	-.04	196	8065
5124	181 190	-24.6	7398	49	7497	72	7429	41	1.14	186	8075
5125	171 180	-24.4	7363	49	7425	71	7383	40	.72	176	8085
5126	161 170	-25.3	7399	49	7484	72	7426	41	.98	166	8095
5127	151 160	-25.3	7337	48	7476	66	7385	39	1.70	156	8105
5128	141 150	-25.0	7240	48	7360	71	7278	40	1.40	146	8115
5129	131 140	-25.6	7398	49	7460	66	7420	39	.75	136	8125
5130	121 130	-25.4	7383	49	7363	66	7376	39	-.24	126	8135
5131	111 120	-25.1	7297	48	7398	71	7329	40	1.18	116	8145
5132	101 110	-25.0	7367	49	7479	72	7402	41	1.29	106	8155
5133†	76 100	-24.9	7317	48	7344	71	7325	40	.32	88	8172
5134†	51 75	-25.6	7389	49	7598	72	7455	41	2.40	63	8197
5135**	1 50	-25.2	7455	49	7541	72	7482	41	.99	26	8235
5116†	261 270	-24.4	6262	48	6146	67					
Averages											
7292											
.59											
193											

*10-ring samples unless otherwise noted

**44- or 50-ring sample

†25-ring sample

‡rejected radiocarbon datum




Fig. 1. ¹⁴C ages from Stanley River celery-top pine logs, SRT-462 (A) and SRT-157 (B). The data are combined results from measurements in two liquid scintillation counters.

droyears BP. At that position, the SRT-444 data were an average of 14 yr younger than the oak data, with an rms difference of 61 yr. We also examined the robustness of the fit by adding or subtracting an arbitrary 50 or 100 yr to all the SRT-444 ¹⁴C data, but did not find a better fit. The bristlecone pine data covered only part of the region of interest, and was insufficient to allow a formal fitting procedure.

Results for the period, AD 1600–1800 (SRT-31B and SRT-225 in Table 3), indicated small systematic differences between the counters. The results from Counter 3 (measured 1986/87) were generally older than those from Counter 1, whereas the results from Counter 2 (measured 1987/88)




Fig. 2A. Rms differences between the ^{14}C ages measured for SRT-444 and a smoothed (decadal) representation of the German oak data. The minimum difference occurs when ring 0 of SRT-444 is placed at 8260 yr BP. Adding or subtracting an arbitrary 50 or 100 yr to all the SRT-444 ^{14}C data gives a poorer fit between the data sets.




Fig. 2B. ^{14}C ages from Stanley River celery-top pine log, SRT-444 (—); combined results from two counters. The measurements are compared with data from the unified German oak series with ring 0 placed at 7230 BC, *i.e.*, 9179 BP (Kromer *et al.* 1986; —) and a smoothed representation of the German oak data (---). SRT-444 is shown with ring 0 placed at 8260 yr BP, which gives the best match to the German oak data. Bristlecone pine data from the USA (Linick *et al.* 1986; |||) are also shown.

were younger. The results from Counter 1 alone are therefore used in Figure 3, where they show good agreement with the high-precision decadal data of Stuiver and Becker (1986). There are no significant differences in phase or amplitude of the variations. However, our data are, on average, slightly younger than the corresponding northern hemisphere data, and not older by *ca.* 36 yr, as would be expected from other comparisons of northern and southern hemisphere wood (Lerman, Mook & Vogel 1970; Vogel *et al.* 1986). Since the northern and southern hemisphere measurements were made in different laboratories, we considered the possibility of an error in the modern standard determination in our laboratory. Thus, two of our pretreated wood samples were sent to Seattle for measurement; those results coincide with northern hemisphere data (Fig. 3).

DISCUSSION AND CONCLUSION

Comparisons of conventional ^{14}C ages with results from other dating methods provide data on ^{14}C calibration for the late Pleistocene. Uranium-thorium data have been published by Bard *et al.* (1990); varve data by Stuiver (1971) and Stuiver *et al.* (1986); and summaries of thermoluminescence data were given by Barbetti (1980) and Aitken (1987). These data are summarized in Figure 4.

TABLE 3. Radiocarbon Dates From Stanley River Huon Pine Trees, SRT-31B, SRT-40 and SRT-225*

SUA-no.	Tree ring span yr AD	¹³ C (‰)	Counter 1 yr BP	Counter 2 yr BP	Counter 3 yr BP	Combined yr BP	z-stat	Dendro yr AD
			± σ	± σ	± σ	± σ	± σ	
<i>SRT-31B</i>								
5047	1610	1614	342	50	405	371	.85	1612
5048	1615	1619	399	50	414	406	.20	1617
5049	1620	1624	329	50	388	354	.77	1622
5050	1625	1629	311	50	376	339	.86	1627
5051	1630	1634	313	50	384	344	.94	1632
5052	1635	1639	273	50	264	269	-.12	1637
5056	1640	1644	320	35	303	316	-.24	1642
5057	1645	1649	258	35	180	236	-1.20	1647
5058	1650	1654	258	33	274	262	.26	1652
5059	1655	1659	166	34	167	166	.02	1657
5060	1660	1664	158	34	197	169	.60	1662
5061	1665	1669	144	34	154	147	.15	1667
5062**	1670	1694	120	34	126	122	.10	1682
5063†	1695	1704	72	34	10	57	-.89	1700
5064†	1705	1714	88	34	21	69	-1.04	1710
5065†	1715	1724	142	34	87	129	-.79	1720
5066†	1725	1734	102	34	29	82	-1.13	1730
5067†	1735	1744	158	34	116	146	-.65	1740
5068†	1745	1754	191	34	153	182	-.54	1750
5069†	1755	1764	158	34	143	154	-.23	1760
5070†	1765	1774	163	34	69	141	-1.35	1770
5071	1775	1779	118	34	104	114	-.22	1777
5072	1780	1784	175	34	63	144	-1.73	1782
5073	1785	1789	209	34	75	172	-2.07	1787
5074	1790	1794	230	34	145	206	-1.31	1792
5075	1795	1799	174	34	144	166	-.46	1797
5076	1800	1804	99	34	120	105	.32	1802

TABLE 3. (Continued)

SUA- no.	Tree ring span yr AD	¹³ C (‰)	Counter 1 yr BP	± σ	Counter 2 yr BP	± σ	Counter 3 yr BP	± σ	Combined yr BP	± σ	z-stat	Dendro yr AD
<i>SRT-40</i>												
5077†	1350	1399	-21.5	566	35	524	55	554	30	-0.64	1375	
<i>SRT-225</i>												
5085†	1601	1610	-24.6	358	35	297	55	340	30	-0.94	1606	
5086†	1611	1620	-24.7	294	35	359	55	313	30	1.00	1616	
<i>SRT-31B</i> (Stuiver, personal communication 1992)												
†	1660	1664		208	14						1662	
	1765	1774		174	15						1770	

*5-ring sample unless otherwise noted

**25-ring sample

†10-ring sample

*50-ring sample




Fig. 3. ^{14}C ages from Stanley River Huon pine trees, SRT-31B and SRT-225, for the period AD 1600–1800; Counter 1 data only. The results are compared with the high-precision decadal curve of Stuiver and Becker (1986). Two measurements were made on our samples by Minze Stuiver (\diamond).




Fig. 4. Comparison of conventional ^{14}C ages with other dating methods for part of the late Pleistocene and early Holocene. The solid line would be the ideal relationship if conventional ^{14}C ages were always equal to true ages, and the dashed line shows the deviation expected for the 5568-yr half-life. The U-Th data are from Bard *et al.* (1990), the varve data from Stuiver (1971) and Stuiver *et al.* (1986), and the TL data from Barbetti (1980) and Aitken (1987). ^{14}C data from southern Germany (Becker & Kromer 1986) are from a floating Preboreal tree-ring sequence, shown here on the assumption that the ^{14}C ages are, on average, 1 ka too young (*i.e.*, ring 0 for the Preboreal pine is tentatively placed at 9031 BC or 10,980 BP). The Stanley River data from SRT-462 and SRT-157 are shown here on the assumption that ^{14}C ages are 2 ka too young.

Virtually all data indicate that ^{14}C ages are younger than true ages during the late Pleistocene, with probable differences ranging from 2.6 ka at 15 ka BP to 0.9 ka at 10 ka BP (Stuiver *et al.* 1991).

^{14}C data from the Stanley River and from German Preboreal pine (Fig. 4) show trend changes on a time scale of 1 or 2 centuries, such as those seen throughout the Holocene. The Preboreal data also show a plateau in the calibration curve with nearly constant ^{14}C ages (9600 BP) over several centuries in early Holocene time.

Tasmanian ^{14}C data from SRT-444 have been wiggle-matched with German oak data. With ring 0 of the oak data at 7230 BC (Kromer *et al.* 1986), our match places ring 0 of SRT-444 at 8260 dendroyears BP. With that match, our data appear to be slightly, but not significantly, younger than the northern hemisphere data.

Our data from the southern hemisphere for the period, AD 1600–1800, also indicate little or no offset in ^{14}C concentration when compared with northern hemisphere data. These minimal apparent offsets, at around 8.1 k dendroyears BP, and again, in recent centuries, would provide an important constraint on global carbon models if confirmed by further measurements.

Detailed intercomparisons with other laboratories are being planned, and it should be possible eventually to estimate precisely the offset between the northern and southern hemispheres during early and mid-Holocene time.

ACKNOWLEDGMENTS

We thank the Forestry Commission, Tasmania, the Hydro Electric Commission of Tasmania, and Corinna Sawmills for logistical support in the field, and the Australian Research Council and the University of Sydney for financial support. We also thank Minze Stuiver for comparative measurements on two of our wood samples.

REFERENCES

- Aitken, M. J. 1987 Archaeometrical dating: Rapporteur review. In Aurenche, O., Evin, J. and Hours, F., eds., *Chronologies in the Near East. BAR International Series* 379: 207–218.
- Barbetti, M. 1980 Geomagnetic strength over the last 50,000 years and changes in atmospheric ^{14}C concentration: Emerging trends. In Stuiver, M. and Kra, R. S., eds., *Proceedings of the 10th International ^{14}C Conference. Radiocarbon* 22(2): 192–199.
- Bard, E., Hamelin, B., Fairbanks, R. G. and Zindler, A. 1990 Calibration of the radiocarbon timescale over the past 30,000 years using mass-spectrometric U-Th ages from Barbados corals. *Nature* 345: 405–410.
- Becker, B. and Kromer, B. 1986 Extension of the Holocene dendrochronology by the Preboreal pine series, 8800 to 10,100 BP. In Stuiver, M. and Kra, R. S., eds., *Proceedings of the 12th International ^{14}C Conference. Radiocarbon* 28(2B): 961–967.
- Francey, R. J., Barbetti, M., Bird, T., Beardsmore, D., Coupland, W., Dolezal, J. E., Farquhar, G. D., Flynn, R. G., Fraser, P. J., Gifford, R. M., Goodman, H. S., Kunda, B., McPhail, S., Nanson, G., Pearman, G. I., Richards, N. G., Sharkey, T. D., Temple, R. B. and Weir, B. 1984 Isotopes in tree rings – Stanley River Collections 1981/82. CSIRO Division of Atmospheric Research, Aspendale, Victoria. *Technical Paper* 4: 86 p.
- Gupta, S. K. and Polach, H. A. 1985 *Radiocarbon Dating Practices at ANU*. Handbook, Research School of Pacific Studies, Canberra: 173 p.
- Head, J. (ms.) 1979 Structure and chemical properties of fresh and degraded wood. M.Sc. thesis, Australian National University, Canberra: 103 p.
- Kromer, B., Rhein, M., Bruns, M., Schoch-Fischer, H., Münnich, K. O., Stuiver, M. and Becker, B. 1986 Radiocarbon calibration data for the 6th to the 8th millennia BC. In Stuiver, M. and Kra, R. S., eds., *Proceedings of the 12th International ^{14}C Conference. Radiocarbon* 28(2B): 954–960.
- Lerman, J. C., Mook, W. G. and Vogel, J. C. 1970 ^{14}C in tree rings from different localities. In Olsson, I. U., ed., *Radiocarbon Variations and Absolute Chronology*. *Proceedings of the 12th Nobel Symposium*. New York, John Wiley & Sons: 257–299.
- Linick, T. W., Long, A., Damon, P. E. and Ferguson, C. W. 1986 High-precision radiocarbon dating of bristlecone pine from 6554 to 5350 BC. In Stuiver, M. and Kra, R. S., eds., *Proceedings of the 12th*

- International ¹⁴C Conference. *Radiocarbon* 28(2B): 943–953.
- Stuiver, M. 1971 Evidence for the variation of atmospheric C¹⁴ content in the late Quaternary. In Turekian, K. K., ed., *The Late Cenozoic Glacial Ages*. New Haven, Connecticut, Yale University Press: 57–70.
- Stuiver, M. and Becker, B. 1986 High-precision decadal calibration of the radiocarbon time scale, AD 1950–2500 BC. In Stuiver, M. and Kra, R. S., eds., Proceedings of the 12th International ¹⁴C Conference. *Radiocarbon* 28(2B): 863–910.
- Stuiver, M., Braziunas, T. F., Becker, B. and Kromer, B. 1991 Climatic, solar, oceanic and geomagnetic influences on late-glacial and Holocene atmospheric ¹⁴C/¹²C change. *Quaternary Research* 35: 1–24.
- Stuiver, M., Kromer, B., Becker, B. and Ferguson, C. W. 1986 Radiocarbon age calibration back to 13,300 years BP and the ¹⁴C age matching of the German oak and US bristlecone pine chronologies. In Stuiver, M. and Kra, R. S., eds., Proceedings of the 12th International ¹⁴C Conference. *Radiocarbon* 28(2B): 969–979.
- Stuiver, M. and Pearson, G. W. 1986 High-precision calibration of the radiocarbon time-scale, AD 1950–500 BC. In Stuiver, M. and Kra, R. S., eds., Proceedings of the 12th International ¹⁴C Conference. *Radiocarbon* 28(2B): 805–838.
- Vogel, J. C., Fuls, A., Visser, E. and Becker, B. 1986 Radiocarbon fluctuations during the third millennium BC. In Stuiver, M. and Kra, R. S., eds., Proceedings of the 12th International ¹⁴C Conference. *Radiocarbon* 28(2B): 935–938.