

# BECOME A LIFE MEMBER AND YOU'LL JOIN PRESTIGIOUS COMPANY



Become a Life Member of Fauna & Flora International to show an extraordinary commitment to an organisation that has been at the forefront of conservation for over 100 years.

Since its inception in 1903 as the world's first international conservation organisation Fauna & Flora International has been working tirelessly for the preservation of plants and animals across the world. For over 100 years Life Members have provided the support and commitment that has helped us to achieve so much, not least key achievements such as:

- Reintroducing the Arabian oryx to the wild
- Establishing the International Gorilla Conservation Programme
- Establishing Vietnam's first locally managed conservation organisation

Collaboration is key to our approach, and wherever possible we work with other global conservation organisations to ensure we are effective and efficient. That's why in 2015 we moved into the David Attenborough Building of the Cambridge Conservation Initiative with several other global conservation groups. Now we are asking you to be part of our wider collaborative work by becoming a Life Member.

As a Life Member you will receive Oryx—The International Journal of Conservation and our annual magazine Fauna & Flora, and you will also be invited to special events, where you can network with some of the world's leading conservationists. You will be joining a select group of supporters who have shown an extraordinary commitment to international conservation.

By joining Fauna & Flora International as a Life Member with a one-off payment of £1,500 you will be making a genuine difference to our conservation work and will forever be part of our global conservation organisation.



Gary Morrisroe/FFI

"I have been a member of Fauna & Flora International since the 1950s... investment in the work of FFI is truly an investment in the future of our planet"

Sir David Attenborough

To join as a Life Member, you can:

Call us on +44 (0)1223 749 019

Email us at [members@fauna-flora.org](mailto:members@fauna-flora.org)

Visit [www.fauna-flora.org/life-membership](http://www.fauna-flora.org/life-membership)



- 321 How can we increase capacity for species conservation in the post-2020 Global Biodiversity Framework? *G. Maggs, H.D. Slater & P.J.K. McGowan*
- 323 Briefly
- 329 Conservation news
- Ungulate conservation**
- 336 Swayne's hartebeest in Ethiopia: population estimate, genetic variability and competition with livestock, *M. Tamrat, A. Atickem, Ø. Flagstad, M. Fischer, C. Roos, P. Evangelista et al.*
- 345 Estimating forest antelope population densities using distance sampling with camera traps, *R. Amin, H. Klair, T. Wacher, C. Ndjassi, A. Fowler, D. Olson & T. Bruce*
- 352 A systematic survey of online trade: trade in Saiga antelope horn on Russian-language websites, *D.L. Roberts, K. Mun & E.J. Milner-Gulland*
- 360 Probable extirpation of the hog deer from China: implications for conservation, *C. Ding, J. Liu, C. Li & Z. Jiang*
- 367 Genetic evidence indicates the occurrence of the Endangered Kashmir musk deer *Moschus cupreus* in Uttarakhand, India, *A. Kumar, B. Singh, S. Sahoo, K.B. Gautam & S.K. Gupta*
- 373 Tapirs in trouble: estimating Baird's tapir densities in the Sierra Madre de Chiapas, Mexico, *M. Rivero, J.A. de la Torre, G. Camacho, E.J. Naranjo, M.W. Tobler, C.A. Jordan et al.*
- 383 Free-ranging dogs as a potential threat to Iranian mammals, *D. Nayeri, A. Mohammadi, A.T. Qashqaei, A.T. Vanak & M.E. Gompper*
- 390 Intact forests of the Hkakabo Razi Landscape are a hotspot of bat diversity in South-east Asia, *P.J.J. Bates, P. Soisook, S.S.L. Oo, M. Suarez-Rubio, A. Pimsai, A. Dejtaradol & S.C. Renner*
- 396 Female wanted for the world's rarest turtle: prioritizing areas where *Rafetus swinhoei* may persist in the wild, *T.V. Pham, O. Le Duc, C. Bordes, B. Leprince, C. Ducotterd, T. Zuklin et al.*
- 404 The global magnitude and implications of legal and illegal wildlife trade in China, *Y. Jiao & T.M. Lee*
- 412 Uncovering prevalence of pangolin consumption using a technique for investigating sensitive behaviour, *A. Olmedo, D. Verissimo, E.J. Milner-Gulland, A. Hinsley, H.T.T. Dao & D.W.S. Challender*
- 421 Occupancy patterns of prey species in a biological corridor and inferences for tiger population connectivity between national parks in Bhutan, *L. Letro, K. Fischer, D. Duba & T. Tandim*
- 429 Biological richness of Gunung Slamet, Central Java, and the need for its protection, *C. Devenish, A.R. Junaid, Andriansyah, R. Saryanthi, S.(B.) van Balen, F. Kaprawi et al.*
- 439 Status of a second viable population of the Critically Endangered Delacour's langur in Viet Nam, *A.T. Nguyen, H. Trinh-Dinh, M. Le & N. Manh Ha*
- 442 Intermediate levels of wood extraction may facilitate coexistence of an endemic arboreal marsupial and Indigenous communities, *H. González-Ancín, M. Spínola, J.M. Mora-Benavides, J.C. Sáenz, A. Paillacar & F.E. Fontúrbel*
- 451 Clinging to survival: Critically Endangered Chapman's pygmy chameleon *Rhampholeon chapmanorum* persists in shrinking forest patches, *K.A. Tolley, C.R. Tilbury, J.M. da Silva, G. Brown, Y. Chapeta & C.V. Anderson*
- 457 Illegal settlement in the Babile Elephant Sanctuary is threatening the resident elephant population, *E. Neil & E. Greengrass*
- 465 Field surveys can improve predictions of habitat suitability for reintroductions: a swift fox case study, *Z. Paraskevopoulou, H. Shamon, M. Songer, G. Ruxton & W.J. McShea*
- 475 Detecting tropical wildlife declines through camera-trap monitoring: an evaluation of the Tropical Ecology Assessment and Monitoring protocol—CORRIGENDUM, *L. Beaudrot, J. Ahumada, T.G. O'Brien & P.A. Jansen*
- 476 A systematic survey of online trade: trade in Saiga antelope horn on Russian-language websites—CORRIGENDUM, *D.L. Roberts, K. Mun & E.J. Milner-Gulland*
- 477 Biological richness of Gunung Slamet, Central Java, and the need for its protection—CORRIGENDUM, *C. Devenish, A.R. Junaid, Andriansyah, R. Saryanthi, S.(B.) van Balen, F. Kaprawi et al.*
- Book reviews*
- 478 Extinctions: Living and Dying in the Margin of Error
- 478 All Asian Primates
- 479 Climate Ghosts: Migratory Species in the Anthropocene
- 480 Grants & opportunities
- Cover* Swayne's hartebeest *Alcelaphus bucelaphus swaynei*, once widely distributed in the Horn of Africa, is now limited to two relict populations in the Ethiopian Rift Valley, in Senkele Swayne's Hartebeest Sanctuary and Maze National Park. Investigation of the size and genetic diversity of these two remaining populations indicates a positive population trend in both protected areas, but suggests that spatial overlap with livestock could lead to future population decline as a result of resource competition and disease transmission. Genetic diversity is higher in the larger population in Maze National Park. A translocation of individuals from Senkele Swayne's Hartebeest Sanctuary to other protected areas within the subspecies' former range could reduce the vulnerability of this population. For further details, see pp. 336–344. (Photograph © NaturesMomentsUK/Shutterstock)